



OPEN ACCESS

Changes in admission thresholds in English emergency departments

Steven Wyatt,¹ Kieran Child,¹ Andrew Hood,¹ Matthew Cooke,² Mohammed A Mohammed³

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/emmermed-2016-206213>).

¹Strategy Unit, NHS Midlands and Lancashire Commissioning Support Unit, West Bromwich, UK

²Warwick Medical School, University of Warwick, Coventry, UK

³Faculty of Health Studies, University of Bradford, Bradford, UK

Correspondence to

Steven Wyatt, Strategy Unit - NHS Midlands and Lancashire Commissioning Support Unit Kingston House 438-450 High Street West Bromwich B70 9LD UK; swyatt@nhs.net

Received 16 July 2016

Revised 22 June 2017

Accepted 28 June 2017

ABSTRACT

Background The most common route to a hospital bed in an emergency is via an Emergency Department (ED). Many recent initiatives and interventions have the objective of reducing the number of unnecessary emergency admissions. We aimed to assess whether ED admission thresholds had changed over time taking account of the casemix of patients arriving at ED.

Methods We conducted a retrospective cross-sectional analysis of more than 20 million attendances at 47 consultant-led EDs in England between April 2010 and March 2015. We used mixed-effects logistic regression to estimate the odds of a patient being admitted to hospital and the impact of a range of potential explanatory variables. Models were developed and validated for four attendance subgroups: ambulance-conveyed children, walk-in children, ambulance-conveyed adults and walk-in adults.

Results 23.8% of attendances were for children aged under 18 years, 49.7% were female and 30.0% were conveyed by ambulance. The number of ED attendances increased by 1.8% per annum between April 2010–March 2011 (year 1) and April 2014–March 2015 (year 5). The proportion of these attendances that were admitted to hospital changed negligibly between year 1 (27.0%) and year 5 (27.5%). However, after adjusting for patient and attendance characteristics, the odds of admission over the 5-year period had reduced by 15.2% (95% CI 13.4% to 17.0%) for ambulance-conveyed children, 22.6% (95% CI 21.7% to 23.5%) for walk-in children, 20.9% (95% CI 20.4% to 21.5%) for ambulance conveyed adults and 22.9% (95% CI 22.4% to 23.5%) for walk-in adults.

Conclusions The casemix-adjusted odds of admission via ED to NHS hospitals in England have decreased since April 2010. EDs are admitting a similar proportion of patients to hospital despite increases in the complexity and acuity of presenting patients. Without these threshold changes, the number of emergency admissions would have been 11.9% higher than was the case in year 5.

INTRODUCTION

Rising demand for emergency hospital admissions is a feature of many healthcare systems.¹ There were 5.6 million emergency admissions to NHS hospitals in England during April 2014–March 2015, an increase of 27% over 10 years.² These admissions account for almost 70% of hospital bed days. Some emergency admissions are preventable or could be avoided if alternatives were available, and as such represent an unwarranted cost for commissioners.^{3,4}

Key messages

What is already known on this subject?

- The most common route to a hospital bed in an emergency is via an ED.
- Many recent initiatives and interventions have the objective of reducing the number of unnecessary emergency admissions.
- Several studies have identified patient and attendance characteristics that are associated with increased risk of admission.

What this study adds?

- The casemix-adjusted odds of admission via ED to NHS hospitals in England have decreased since April 2010.
- The number of attendances that have a low probability of admission has reduced since April 2010.

Many recent NHS policy initiatives and commissioning interventions have been designed to avoid unnecessary emergency admissions.^{5–7} Clinical decision units in EDs, ambulatory care units and acute frailty units have expanded in recent years, with one aim being to reduce admissions. Since April 2010 hospital trusts have been paid at a marginal rate (30%) for emergency admissions above historical levels.⁸ This policy was introduced to ‘incentivise lower rates of emergency admissions’ and ‘stimulate acute providers to work with other parties in the local health economy to reduce the demand for emergency care’.⁸ Others have suggested that some emergency admissions may be supply-induced or associated with ED performance targets.^{9,10}

The most common route to a hospital bed in an emergency is via an ED.² Admissions via ED represent an increasing proportion of all emergency admissions.¹¹ Approximately one-fifth of patients attending EDs in English NHS hospitals are admitted to a hospital bed.¹² English NHS hospitals are expected to triage, treat and discharge or admit patients attending ED within 4 hours.¹³ The standard was originally set at 98% of patients but was lowered in 2010 to 95% to allow greater time for complex investigations.¹⁴ Performance against this standard has been deteriorating since 2010 and has fallen below 95% since 2013.¹⁵

Previous studies have identified a range of casemix variables (eg, age, presenting condition, arrival mode) and attendance characteristics (eg, time of year, day of week) that are associated with



CrossMark

To cite: Wyatt S, Child K, Hood A, et al. *Emerg Med J* Published Online First: [please include Day Month Year]. doi:10.1136/emmermed-2016-206213



an increased risk of admission for a patient in ED.^{16–20} We use a routine English administrative data set to assess changes in the casemix-adjusted odds of admission via ED between April 2010–March 2011 (year 1) and April 2014–March 2015 (year 5) to provide insight into whether ED admission thresholds have changed and the impact of admission avoidance policies and interventions.

METHODS

Study design, setting and population

We conducted a retrospective cross-sectional analysis of 20 224 711 attendances at 47 consultant-led EDs in England between April 2010 and March 2015. Given the importance of diagnosis as a casemix variable, the NHS organisations included in the study were those where at least 70% of ED attendances had a valid primary diagnosis code and at least 95% had a valid disposal code within each year of the study period. Follow-up attendances were excluded along with attendances for patients who were dead on arrival, who died in the department or who left the department having refused treatment or before being treated.

Variables and data sources

An anonymised extract of the Accident and Emergency Hospital Episode Statistics (HESAE) was obtained from the NHS Health and Social Care Information Centre for the 5-year period from April 2010 to March 2015.

The outcome variable was defined with reference to the disposal code in HESAE, taking a value of 1 when the patient was admitted to a hospital bed with the same health-care provider and a value of 0 in all other circumstances (eg, discharge, referral or transfer to another service). Admissions may include short-stay admissions to assessment units outside of the ED.

Potential explanatory variables were identified with reference to previous studies and included patient characteristics (age, gender, ethnicity, deprivation, primary diagnosis, prior ED attendances and admissions), arrival year, time of year, part of week and time of day of arrival, arrival mode (by ambulance or other means) and the ED of attendance.^{16–20}

A patient's age, gender, ethnicity and primary diagnosis, and the date, time and ED of the attendance, are routinely recorded in HESAE.¹² Patient ethnicities were assembled into six groups: white, Asian/Asian British, black/black British, mixed parentage, other ethnic groups and not known/not stated. Socioeconomic status was measured using the 2004 Index of Multiple Deprivation (IMD) rank assigned to the lower super output area in which the patient lived.²¹ IMD ranks were grouped into quintiles. A patient's prior A&E activity was assigned to three levels: none, attended ED at least once but not admitted via ED, and attended and admitted at least once. Prior A&E activity levels were assigned for two time periods: the 28 days before attendance and between 29 and 365 days before attendance. The 39 two-digit primary A&E diagnosis classification codes in HESAE were used to define the patient's diagnosis. Arrival year (1=April 2010–March 2011; 2=April 2011–March 2012; 3=April 2012–March 2013; 4=April 2013–March 2014; and 5=April 2014–March 2015), time of year (November–February and March–October), part of week (weekday and weekend) and time of day (08:00–22:00 and 22:00–08:00) were derived from the arrival date and time fields in HESAE.

We excluded from the analysis 2.5% of attendances that did not have a valid age, gender, IMD quintile or disposal code.

Statistical methods

We used mixed-effects logistic regression to estimate the association between the odds of a patient being admitted to hospital and the impact of each of the potential explanatory variables. Initial univariate analysis indicated markedly different unadjusted odds of admission for children and adults and between patients conveyed by ambulance and those who arrived by some other means (henceforth referred to as walk-ins). These differences remained substantial having adjusted for other covariates. Given that the records available to build models were plentiful, to minimise the reliance on interaction terms and to increase the model accuracy, we developed four separate models for ambulance-conveyed children, walk-in children, ambulance-conveyed adults and walk-in adults.

All candidate predictor variables were included in the model on the basis of adequately strong univariate association with the outcome variable or because inclusion improved the fit of the multivariate model. The hospital trust (provider) of the ED was included as a random effect to reflect clustering of attendances within hospital trusts. Final model fit was measured using the C-statistic (area under the receiver operating characteristic curve), calibration plots and the Hosmer-Lemeshow goodness-of-fit test.

The impact of adding interaction terms was tested as a sensitivity analysis. All two-way interaction terms between the fixed effects (other than arrival year) were tested individually and in combination, for their impact on the coefficients for the arrival year variable. In addition, the coefficients of interaction terms between arrival year and each of the other fixed effects were assessed.

To further explore changes in admission thresholds over time, we applied the casemix-adjusted odds of admission from year 1 to attendances in years 2–5.

Data processing was conducted in Microsoft SQL Server 2012 and analysis in R V.3.2.3 statistical software package.

RESULTS

Description of ED attendances

The analysis included 20.2 million ED attendances at 47 hospitals trusts (online supplementary file 1) between 1 April 2010 and 31 March 2015 (figure 1).

There were 4.81 million attendances (28%) for patients aged 0–17 years, and of these 605 000 were conveyed to ED by ambulance and 4.21 million by walk-in. There were 15.4 million (72%) aged 18+ years, and of these 5.47 million were conveyed to ED by ambulance and 9.94 million by walk-in.

The number of attendances increased by an average of 1.8% per year from 3.89 million in year 1 to 4.18 million in year 5 (table 1). The most common primary diagnoses were dislocation/fracture/joint injury/amputation, gastrointestinal conditions, sprain/ligament injury, laceration, respiratory conditions and contusion/abrasion, each with more than 1 million attendances during the study period (online supplementary file 2, table B1). Thirty-two per cent of attendances occurred during the winter (November–February), 29% on the weekend (similar to the rate of activity during the week) and 19% overnight (22:00–08:00). Six per cent of patients attending ED had attended in the previous 28 days but had not been admitted; a further 4% had attended ED and been admitted (online supplementary file 2, table B2).

Recording of patients' ethnicity improved substantially, increasing from 30.5% in year 1 to 90.1% in year 5.




Figure 1 Selection of study population.

Model description and model fit

In each of the four models, age group, gender, deprivation, ethnicity, diagnosis, arrival year, time of year, part of week and time of day of attendance, activity in the previous month, and year of attendance were included in the model as fixed effects. The hospital trust (provider) of the ED was included as a random effect.

A full description of the four models, including ORs for the fixed effects and SD for the random effect, can be found in online supplementary file 3, tables C1–5.

The C-statistic was 0.73 for the child ambulance model, 0.81 for the child walk-in model, 0.76 for the adult ambulance model and 0.84 for the adult walk-in model. Calibration plots are provided in figure 2.

Odds of admission by arrival year

The unadjusted and adjusted odds of admission by arrival year are shown in table 2. For children conveyed by ambulance, the unadjusted odds of admission were 9.7% higher in year 5 than in year 1 (OR 1.097, 95% CI 1.079 to 1.116). However, having adjusted for the other covariates in the model, the odds of admission are shown to be 15.2% lower in year 5 than in year 1 (OR 0.848, 95% CI 0.830 to 0.866). The adjusted odds of admission were also considerably lower in year 5 than in year 1 for child walk-in attendances (OR 0.774, 95% CI 0.765 to 0.783), adult ambulance-conveyed attendances (OR 0.791, 95% CI 0.785 to 0.796) and adult walk-in attendances (OR 0.771, 95% CI 0.765 to 0.776). In each of the four models, the adjusted odds of admission decreased monotonically with arrival year. Effects

sizes are shown in online supplementary file 4 as average relative risks and average absolute risk reductions using the approach described by Grant.²²

Changes in risk profile of attendances

Of all cases, 27.5% were admitted in year 5. The models suggest that 30.8% of cases in year 5 would have been admitted if they had experienced the casemix-adjusted odds of admission that applied in year 1.

The proportion of low admission risk ($p \leq 0.2$) attendances fell from 52.0% in year 1 to 46.3% in year 5 (see figure 2). Correspondingly there were increases in the proportion of high admission risk ($p > 0.6$) attendances over the same period, from 15.9% to 19.7% (figure 3).

Sensitivity analyses: inclusion of interaction terms

The inclusion of many of the two-way interaction terms between the fixed effects improved the model fit, but did not materially affect the relationship between the outcome variable and arrival year. In most cases the coefficients of the interaction terms between arrival year and other fixed effects were small (ORs less than 0.5 or greater than 2), with larger coefficients found only with some low volume diagnoses variables.

DISCUSSION

Key findings

Having adjusted for other factors that have previously been shown to be associated with the odds of admission, the four models for child ambulance, child walk-in, adult ambulance and adult walk-in ED attendances independently suggest that the odds of admission have decreased over time.^{16–20} Patients attending in year 5 are substantially less likely to be admitted than patients attending in year 1 with similar characteristics and at similar times of the year, week and day. Ambulance-conveyed children are 15.2% less likely to be admitted in year 5 than in year 1. The reduction is more marked for other groups: a 22.6% reduction for child walk-in attendances, 20.9% for ambulance-conveyed adults attending and 22.9% for adult walk-in attendances.

While there are a number of potential explanations for this finding, we note that these changes coincide with the development of ambulatory emergency care, the wider use of fast track clinics, increases in the number of emergency medicine consultants and the development of the acute medicine specialty.²³ Over the same period there have been increases in the rate of investigations and treatments delivered in ED and in the average time between patients being seen and being discharged from ED.^{24 25} This was part of the rationale for changing the 4-hour standard from 98% of patients to 95% in 2010.¹⁴

We estimate that there would have been 137 000 (11.9%) additional emergency admissions in the selected hospitals in year 5, if the casemix-adjusted odds of admission from year 1 had applied. Estimating the cost to commissioners (or forgone income for providers) of these avoided admissions is not trivial since the average cost of the avoided admissions is likely to be lower than the average cost of all admissions. However, applying the lower decile tariff for emergency admissions provides a conservative savings estimate of £65 million in year 5.

Reductions in average length of stay in hospital and the increase in the number of patients discharged on the same day as admission that have been reported elsewhere should be seen as more remarkable given the increased average acuity of patients admitted via ED.²

Table 1 Patient characteristics of attendances

Number of attendances		April 2010–March 2011	April 2011–March 2012	April 2012–March 2013	April 2013–March 2014	April 2014–March 2015
Total attendances		3 890 957	3 994 016	4 077 353	4 078 857	4 183 528
Subgroup	Child via ambulance	122 594	117 443	119 278	125 503	119 907
	Child walk-in	841 525	847 953	836 137	831 359	851 091
	Adult via ambulance	1 027 281	1 057 863	1 102 920	1 125 592	159 010
	Adult walk-in	1 899 557	1 970 757	2 019 018	1 996 403	2 053 520
Age (years)	0	102 857	102 231	110 715	104 462	107 525
	1–4	278 615	284 443	293 566	296 468	299 783
	5–12	321 507	324 100	312 127	318 298	329 357
	13–17	261 140	254 622	239 007	237 634	234 333
	18–34	935 292	953 570	965 533	964 577	976 519
	35–54	836 176	853 970	865 978	859 363	869 102
	55–74	618 819	646 466	678 733	683 209	712 571
	75+	536 551	574 614	611 694	614 846	654 338
Gender	Male	1 989 629	2 033 730	2 036 712	2 038 930	2 079 707
	Female	1 901 328	1 960 286	2 040 641	2 039 927	2 103 821
Deprivation (IMD04)	Quintile 1 (most deprived)	1 103 751	1 092 746	1 112 747	1 121 666	1 154 306
	Quintile 2	769 280	790 969	809 231	808 259	828 239
	Quintile 3	711 936	745 523	762 722	764 983	784 924
	Quintile 4	687 724	716 119	730 576	727 656	743 337
	Quintile 5	618 266	648 659	662 077	656 293	672 722
Ethnicity	White	1 109 765	2 175 454	3 072 385	3 193 024	3 418 286
	Asian/Asian British	12 188	29 272	47 526	49 146	54 708
	Black/Black British	30 287	97 499	153 756	169 082	188 384
	Mixed parentage	10 953	23 403	35 062	38 656	44 118
	Other ethnic group	21 708	39 044	61 212	62 209	63 549
	Not known/Not stated	2 706 056	1 629 344	707 412	566 740	414 483

IMD, Index of Multiple Deprivation.

In addition we found that the number of attendances with a low risk admission ($p < 0.2$) fell by 1.1% per annum, whereas the number of all other attendances grew by 4.7% per annum. This finding fits with reported increases in general practitioner consultation rates, walk-in centre/minor injuries units attendances, ambulance ‘hear and treat’ and ‘see and treat’ cases.^{12 26 27}

If low admission risk attendances had increased at the same rate as all other attendances, then we estimate that there would have been an additional 564 000 low admission risk attendances in year 5. If we conservatively assume that these attendances would have

been charged at the lowest available tariff, then the avoided cost to the commissioner can be estimated at £32 million in year 5.

Relation to existing literature

The effects of casemix variables on the risk of admissions that we found in this study reflect those found in previous published studies, notably an increased risk of admission with ambulance conveyance, old age, attendance during the daytime and during the week, and with recent prior admissions.^{16–18 28} There is considerable interest in the number of deaths in hospitals in England during the weekend, and one paper has suggested that the ‘weekend-effect’ might be explained by a lower probability of admission at the weekend.²⁸ This paper also finds that the casemix-adjusted odds of admission are lower on the weekend (see online supplementary file 3, table C3).

Limitations

The models are built on a subset of ED attendances. While some exclusions are justified from a design perspective (eg, excluding follow-up attendances), others are based on the availability of coded covariates. Limiting the study population to those hospitals where at least 70% of attendances have a valid diagnosis code and 95% have a valid disposal code in each year of the study introduces the potential for selection bias and raises questions about the extent to which the model can be generalised to all EDs. We note however that no significant interactions were found between the design variable representing ‘diagnosis not classified’ and arrival year. Exclusions for other data quality




Figure 2 Per cent of ED attendances by modelled risk of admission and year.

Table 2 Odds of admission by year

Model	Year	Unadjusted		Adjusted	
		OR	(95% CI)	OR	(95% CI)
Child ambulance	April 2010–March 2011 (ref)	1.000	–	1.000	–
	April 2011–March 2012	1.054	(1.037 to 1.072)	0.937	(0.919 to 0.955)
	April 2012–March 2013	1.129	(1.111 to 1.148)	0.919	(0.901 to 0.938)
	April 2013–March 2014	1.085	(1.068 to 1.104)	0.868	(0.850 to 0.886)
	April 2014–March 2015	1.097	(1.079 to 1.116)	0.848	(0.830 to 0.866)
Child walk-in	April 2010–March 2011 (ref)	1.000	–	1.000	–
	April 2011–March 2012	0.890	(0.882 to 0.899)	0.862	(0.853 to 0.871)
	April 2012–March 2013	0.976	(0.967 to 0.985)	0.849	(0.839 to 0.858)
	April 2013–March 2014	0.932	(0.923 to 0.940)	0.821	(0.812 to 0.831)
	April 2014–March 2015	0.918	(0.909 to 0.926)	0.774	(0.765 to 0.783)
Adult ambulance	April 2010–March 2011 (ref)	1.000	–	1.000	–
	April 2011–March 2012	1.008	(1.002 to 1.013)	0.919	(0.913 to 0.925)
	April 2012–March 2013	1.007	(1.002 to 1.013)	0.838	(0.832 to 0.844)
	April 2013–March 2014	0.968	(0.963 to 0.974)	0.806	(0.800 to 0.811)
	April 2014–March 2015	1.008	(1.002 to 1.013)	0.791	(0.785 to 0.796)
Adult walk-in	April 2010–March 2011 (ref)	1.000	–	1.000	–
	April 2011–March 2012	0.993	(0.988 to 0.999)	0.918	(0.912 to 0.923)
	April 2012–March 2013	1.024	(1.019 to 1.030)	0.837	(0.832 to 0.843)
	April 2013–March 2014	0.989	(0.984 to 0.995)	0.791	(0.786 to 0.797)
	April 2014–March 2015	1.012	(1.007 to 1.018)	0.771	(0.765 to 0.776)

reasons (eg, missing data on gender, lower super output areas) represent only 2.5% of attendances.

We are not aware of any fiscal incentives that might induce providers to systematically misclassify patients' diagnoses in ED in favour of those diagnoses with the greatest risk of admission. Note that unlike inpatient admissions, a patient's diagnosis in ED does not alter the charge for the attendance or influence high profile quality metrics such as the hospital standardised mortality ratio. Substantial changes in the frequency of diagnoses were largely confined to growth in low volumes diagnoses and between years 1 and 2, whereas we find that the casemix-adjusted odds of admission reduced in all years. Diagnoses of septicæmia grew in each year in line with interventions to improve the identification of this condition in ED.

Previous literature has indicated that other factors not included in our models may influence the odds of admission, notably the relative business of the ED, bed occupancy and the distance travelled to ED.^{11 19 20 29} While the precise estimate of the casemix-adjusted odds of admission by arrival year would no doubt change with the inclusion of these variables, it seems unlikely that this would have a material impact on the headline findings.

Although information about the investigations and treatments delivered in ED was available in the HESAE data sets, these were not included in the model because these variables represent the ED response to presenting patients rather than the intrinsic characteristics of patients as they present. Indeed, increases in the number of investigations and treatments in ED may be one of the means by which admission thresholds have been increased.

Age is used as a casemix adjustment variable. If age did not confer a constant risk (of admission) over time, then this might explain some or all of the observed reductions in casemix-adjusted odds of admission. This is indeed feasible if we consider that life expectancy is known to be increasing, and for older adults age can be thought of as a proxy for the latent variable, proximity to death. However, tests for interactions between the age and arrival year variables did not support this hypothesis.

We considered whether the observed results may be explained in part by the considerable improvements in recording of a patient's ethnicity. However the arrival year coefficients were not materially altered by limiting the models to cases with a valid ethnicity or by introducing an interaction term between ethnicity and arrival year.

The outcome variable for the model is admission directly from ED. In some cases, patients may be referred elsewhere for immediate attention (eg, to a fracture clinic) and as a result of this contact be admitted to hospital on the same day. An increase in the frequency of these indirect routes to admissions may explain some or all of the observed reduction in the odds of (direct) admission. We note however that the number of patients attending ED who are referred to A&E clinic, fracture clinic or other outpatient clinics has not changed substantially over the study period (11.0% in year 1 to 10.4% in year 5).

Implications for policy and research

This analysis suggests activity has changed in line with the long-standing UK health policy to ensure that patients access ED and are admitted to hospital only when clinically necessary. It should be noted, however, that this study did not attempt to assess the consequence of these changes on patient experience or outcomes.

Against a backdrop of increasing average acuity of ED attendances without increases in hospital beds (see online supplementary file 1, table A2), policy makers need to regularly review the trade-off between two goals: the narrow process-based target of managing 95% of ED cases within 4 hours and the wider goal of eliminating avoidable or unnecessary hospital admissions.

Additional work would be required to determine whether this approach could be used to compare the casemix-adjusted odds of admission between individual providers.

Further research to explore whether the reduction in odds of admission over time holds when one considers admissions that follow indirectly from an ED attendance (eg, from ED to urgent




Figure 3 Calibration plots.

outpatient clinic to admission) would provide another pathway for consideration. This might equally be explored by evaluating the effectiveness of these urgent outpatient clinics in preventing admissions.

CONCLUSION

Whereas trends in admissions imply that admission avoidance strategies are not working, casemix-adjusted odds of admission reveal a different story. Providers, commissioners and policy makers may benefit from calculating and tracking the casemix of patients presenting at ED and the casemix-adjusted odds of admission as a means of evaluating the impact of interventions to reduce hospital admissions.

Acknowledgements The authors would like to thank Alison Turner, Abeda Mulla, Paul Seamer and Peter Spilsbury for reviewing the draft manuscript.

Contributors SW and AH conceived the work. KC conducted and AH supervised the preliminary data analysis. SW conducted the final data analysis and wrote the first draft of the paper. MAM provided methodological and statistical advice. MC provided clinical and operational interpretation of the findings. All authors commented on draft versions and approved the final version of the article.

Competing interests None declared.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement All data used are published.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2017. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- 1 Wittenberg R, Sharpin L, McCormick B, *et al*. Understanding emergency admissions of older people, Centre for Health Service Economics & Organisation, December 2014. see. . (last checked 6 December 2016) <https://www.chseo.org.uk/downloads/report6-emergencyadmissions.pdf>.
- 2 Health & Social Care Information Centre. Hospital Episode Statistics – Admitted patient Care, England – 2014-15 January 2015 See. . (last checked 4 July 2016) <http://www.hscic.gov.uk/catalogue/PUB19124/hosp-epis-stat-admi-summ-rep-2014-15-rep.pdf>.
- 3 Purdy S. Avoiding Hospital Admissions: what does the research evidence say? . See (last checked 30 May 2016) <http://www.kingsfund.org.uk/publications/avoiding-hospital-admissions>.
- 4 Blunt I, Bardsley M, Grove A, *et al*. Classifying emergency 30-day readmissions in England using routine hospital data 2004-2010: what is the scope for reduction? *Emerg Med J* 2015;32.
- 5 Department of Health, Department of Communities and Local Government. Better Care Fund Policy Framework See. 2014. (last checked 18 November 2015) https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/381848/BCF.pdf.
- 6 NHS England. Enhanced service specification – Avoiding unplanned admissions: practice case finding and patient review for vulnerable people 2015/16 See. 2015. (last checked 18 November 2015) <https://www.england.nhs.uk/commissioning/wp-content/uploads/sites/12/2015/03/avoid-unplanned-admissions-03-15.pdf>.
- 7 Lewis G, Curry N, Bardsley M. Choosing a predictive risk model: a guide for commissioners in England November 2011 See (last checked 18 November 2015) http://www.nuffieldtrust.org.uk/sites/files/nuffield/publication/choosing_predictive_risk_model_guide_for_commissioners_nov11.pdf.
- 8 NHS England and Monitor. 2014/15 National tariff payment system See. 2014. (last checked 15 February 2016) https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/300547/2014-15_National_Tariff_Payment_System_-_Revised_26_Feb_14.pdf.
- 9 Purdy S, Griffin T, Salisbury C, *et al*. Emergency respiratory admissions: influence of practice, population and hospital factors. *J Health Serv Res Policy* 2011;16:133–40.
- 10 Information Centre for Health and Social Care (2009) Further analysis of the published 2007–08 A&E HES Data (Experimental Statistics). Leeds: Information Centre. . (last checked 22 June 2016) See <http://www.hscic.gov.uk/catalogue/PUB02547/acci-emer-pati-jour-2007-2008-rep.pdf>.

- 11 Cowling TE, Soljak MA, Bell D, *et al*. Emergency hospital admissions via accident and emergency departments in England: time trend, conceptual framework and policy implications. *J R Soc Med* 2014;107:432–8.
- 12 Health & Social Care Information Centre. Hospital Episodes Statistics – Accident and Emergency Attendances in England 2013–14. See. 2015. (last checked 18 November 2015) <http://www.hscic.gov.uk/catalogue/PUB16728/acci-emer-atte-eng-2013-14-rep.pdf>.
- 13 England NHS. The Handbook to the NHS Constitution for England, See. 2017. (last checked 21 June 2016) https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/474450/NHS_Constitution_Handbook_v2.pdf.
- 14 Department of Health, President of the College of Emergency Medicine. Letter from Andrew Lansley, Secretary of State for Health to John Heyworth, See. 2010;21. (last checked 21 June 2016) https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/213836/dh_116917.pdf.
- 15 NHS England. A&E Activity and Emergency Admissions statistics, England Time Series see. . (last checked 21 June 2016) <https://www.england.nhs.uk/statistics/wp-content/uploads/sites/2/2015/08/Quarterly-time-series-2004-05-onwards-with-Annual-updated-06-05-2016-Q4-2016.xls>.
- 16 Kim SW, Li JY, Hakendorf P, *et al*. Predicting admission of patients by their presentation to the emergency department. *Emergency Medicine Australasia* 2014;26:361–7.
- 17 Sun Y, Heng BH, Tay SY, *et al*. Predicting hospital admissions at emergency department triage using routine administrative data. *Acad Emerg Med* 2011;18(8):844–50.
- 18 Cameron A, Rodgers K, Ireland A, *et al*. A simple tool to predict admission at the time of triage. *Emerg Med J* 2015;32:174–9.
- 19 Blom MC, Jonsson F, Landin-Olsson M, *et al*. The probability of patients being admitted from the emergency department is negatively correlated to in-hospital bed occupancy - a registry study. *Int J Emerg Med* 2014;7:8.
- 20 Smith A, Dolan D, Mahmoud O, *et al*. Does a Busy Emergency Department have a Lower threshold for referral to the Acute Medical team? The University of Manchester.
- 21 Office of the Deputy Prime Minister. The English Indices of Deprivation 2004 (revised), See. 2004. (last checked 15 February 2016) <http://webarchive.nationalarchives.gov.uk/20120919132719/www.communities.gov.uk/documents/communities/pdf/131209.pdf>.
- 22 Grant RL. Converting an odds ratio to a range of plausible relative risks for better communication of research findings. *BMJ* 2014;348:f7450.
- 23 Health and Social Care Information Centre. NHS Workforce Statistics - Provisional statistics, April 2016. see. 2016. (last checked 9th January 2017). <http://content.digital.nhs.uk/searchcatalogue?productid=20703&topics=2%2fWorkforce%2fStaff+numbers%2fHospital+and+community+health+service+staff&sort=Relevance&size=10&page=2#top>.
- 24 England NHS. A&E Activity and Emergency Admissions statistics, NHS and independent sector organisations in England (June 2016), See. . (last checked 30 May 2016). <https://www.england.nhs.uk/statistics/wp-content/uploads/sites/2/2015/04/2015.06.28-AE-TimeseriesBaG87.xls>.
- 25 The Health Foundation and Nuffield Trust, Quality Watch: a&e Waiting Times, see. . (last checked 30 May 2016) <http://www.qualitywatch.org.uk/indicator/ae-waiting-times>.
- 26 Hobbs FDR, Bankhead C, Mukhtar T, *et al*. Clinical workload in UK primary care: a retrospective analysis of 100 million consultations in England, 2007?14. *The Lancet* 2016;387:2323–30.
- 27 England NHS. Ambulance Quality Indicators: system Indicators Time Series See. 2016. (last checked 30 May 2016) <https://www.england.nhs.uk/statistics/wp-content/uploads/sites/2/2013/04/AmbSYS-TimeSeries-Interactive-Web-File-March-2016-with-revisions.xlsx>.
- 28 Meacock R, Anselmi L, Kristensen SR, *et al*. Higher mortality rates amongst emergency patients admitted to hospital at weekends reflect a lower probability of admission. *J Health Serv Res Policy* 2017;22.
- 29 Rudge GM, Mohammed MA, Fillingham SC, *et al*. The combined influence of distance and neighbourhood deprivation on Emergency Department attendance in a large English population: a retrospective database study. *PLoS One* 2013;8. (last checked 24 June 2016).



Changes in admission thresholds in English emergency departments

Steven Wyatt, Kieran Child, Andrew Hood, Matthew Cooke and Mohammed A Mohammed

Emerg Med J published online September 12, 2017

Updated information and services can be found at:

<http://emj.bmj.com/content/early/2017/09/12/emmermed-2016-206213>

These include:

References

This article cites 8 articles, 2 of which you can access for free at: <http://emj.bmj.com/content/early/2017/09/12/emmermed-2016-206213#BIBL>

Open Access

This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Topic Collections

Articles on similar topics can be found in the following collections

[Open access](#) (88)

Notes

To request permissions go to:

<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:

<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:

<http://group.bmj.com/subscribe/>