Acute carbon monoxide intoxication during pregnancy. One case report and review of the literature

J L Greingor, J M Tosi, S Ruhlmann, M Aussedat

Acute carbon monoxide (CO) poisoning during pregnancy is an uncommon event, but failing to recognise maternal CO intoxication can have a dramatic effect on the fetus. To limit the toxic effects of CO, the use of hyperbaric oxygen is usually recommended. We report the case of one patient admitted for acute CO poisoning during pregnancy.

Case report

A 31 year old (gravida 4, para 3) at 21 weeks gestational age was exposed to CO. CO was produced by a defective water heater. She did not lose consciousness and her vital signs were: regular pulse rate of 90 beats per minute, blood pressure of 100/80 mm Hg and respirations of 20/min. Physical examination was unremarkable. She complained of dizziness and palpitation. She was promptly removed from the source of CO. The patient received 9 l/min supplemental oxygen by face mask.

A level of 24.6% carboxyhaemoglobin was measured in the expired air. High flow oxygen therapy was started immediately. The patient was then treated with 100% O₂ at 2.5 atmospheres absolute for 90 minutes. Treatment with hyperbaric oxygen was started in the 30 minutes after the end of exposure. Sonography confirmed the presence of fetal heart tone of 140/min.

She was discharged in good health one day after exposure. The patient delivered at term a healthy male infant weighing 3800 g.

Discussion

CO intoxication is the most frequently reported poisoning in Western developed countries. In England and Wales acute CO poisoning was responsible for 877 deaths in 1996.¹ In France and in the USA, CO is the leading cause of death by poisoning.^{2,3} Many situations result in the production of CO. Signs are variable and non-specific and may easily be confused with those of other disorders. This explains the difficulty of a diagnosis of acute CO intoxication. Hardy estimates that nearly one third of all cases may go undiagnosed.⁴

Breslau and Freund were responsible for the first published report of acute CO intoxication in pregnant women. Breslau reported the intoxication of two pregnant women admitted to a Zurich hospital after a gas explosion.⁵ One woman had an apparently normal child at delivery, but the other suffered a stillbirth: the death occurring one day after exposure. Freund reported a death in utero during the seventh month of gestation.⁶ The frequency of acute CO poisoning during pregnancy is difficult to estimate. In France, Elkharrat *et al*

found that 4.6% of hospitalised victims of CO poisoning were pregnant women.⁷ In 1994, in the Nord Pas de Calais region, where intoxication is frequent, it was observed that 5.4% of the victims were pregnant women.⁸ In USA, the frequency was as high as 8.5%.⁹

Inhalation of CO ends in tissue hypoxia by inhibiting transport, delivery and utilisation of oxygen. It interferes with cellular metabolism not only by binding to haemoglobin and myoglobin but also by the inhibition of cytochrome a₃ and cytochrome P-450. Recent studies suggest other mechanisms of CO mediated toxicity. Hypoxia may be followed by reoxygenation and reperfusion injury.¹⁰ CO exposure has been shown to cause lipid peroxidation,¹¹ and the production of free oxygen radicals.¹² CO toxicity is not only mediated by hypoxia but also by reperfusion injury.

CO passes through the placenta either by passive diffusion,¹³ or by the facilitated mechanism of a carrier.¹⁴ Placental CO diffusion capacity increases with gestational age and in proportion to fetal weight.¹⁵ This is attributable to the rate of placental blood flow and maternal haemoglobin concentration.¹⁶ The increased area of placental exchange and the decreased distance between maternal and fetal blood are probably partially responsible for this change. Slow dissociation of CO from maternal haemoglobin accounts for the delay in the release of CO and for the accumulation of CO in the fetus.¹⁷ Acute CO toxicity in pregnancy causes fetal tissue hypoxia through two mechanisms: decreased levels of release of maternal oxygen and the transplacental passage of CO. As maternal carboxyhaemoglobin rises and the blood oxygen content decreases, oxygen release by haemoglobin is diminished, and oxygen transport across the placenta decreases dramatically.¹⁶ The fetal umbilical vein oxygen decreases and tissue hypoxia occurs.¹⁸ The fetus is unable to increase cardiac output to compensate this decrease in oxygen.¹⁹ The CO has not yet crossed the placenta yet the fetus already suffers from hypoxia. Maternal carboxyhaemoglobin dissociation creates a pressure gradient between maternal and fetal blood. CO crosses the placenta and combines with fetal haemoglobin. Fetal carboxyhaemoglobin levels are approximately 10%–15% higher than maternal levels.¹⁹ Elimination is slower in the fetus.^{18,20} Maximal concentrations of CO in fetal blood are found after about four hours.¹⁹

Hyperbaric oxygen not only accelerates the dissociation of CO from haemoglobin and from the mitochondrial respiratory chain but also increases the level of dissolved oxygen in

CHR Metz, Metz, France

Correspondence to:
Dr Greingor, 35 rue de la
folie, 57050 Metz, France
(JGreingor@aol.com)

Accepted for publication
26 September 2000

Table 1 Clinical grading of central nervous system severity for acute carbon monoxide poisoning victims³⁰

Grade	Symptoms
Grade 1	oriented for time, place and person
Grade 2	headache, dizziness, nausea disorientation for time, place or person confusion or abnormal behaviour
Grade 3	headache, nausea, dizziness disorientation, loss of recent memory, muscle weakness or incoordination
Grade 4	if intubated: appropriate responses to simple commands not intubated: inappropriate responses to simple commands
Grade 5	intubated: requiring sedation comatose responding only to pain not responding to any stimulus

blood and thus prevents further injury. Its action on free radical production has recently been emphasised.

Oxygen shortens the half life of carboxyhaemoglobin, which is four to five hours at atmospheric oxygen levels. Hyperbaric oxygen reduces this half life to 20 minutes. Fetal carboxyhaemoglobin half life in hyperbaric conditions is unknown. Reduction of maternal carboxyhaemoglobin probably greatly contributes to the prompt reduction in fetal levels. Approximately 98.5% of oxygen binds to haemoglobin and only 1.5% is dissolved in normobaric, normoxic conditions. Plasma volume increases by about 50% during pregnancy, theoretically permitting the dissolved oxygen levels to rise. At rest, the tissues extract 5 to 6 ml of oxygen per decilitre.²¹ With a 100% administered oxygen at 3 ATA, dissolved oxygen content is approximately 6 ml per decilitre of blood, which satisfies tissue needs²² without any contribution from haemoglobin bound oxygen. Hyperbaric oxygen reduces the production of free radicals,²³ whereas normobaric oxygen is unable to reduce free radical concentration.²⁴

The level of fetal morbidity and mortality in acute carbon monoxide poisoning is significant. The teratogenicity of CO is controversial. Norman *et al*, from a review of the literature, have concluded that anatomical malformations are more frequent when intoxication occurs during the first stage of gestation²⁵; malformations were reported in 6 of 12 cases. Skeletal abnormalities are frequent: malformations of the limbs are common. When exposure happens in the later stages of pregnancy, anatomical malformations are rare: one case of 48.²⁵ Courtens proposes the systematic screening for acute exposure to CO in early pregnancy in every reported case of malformation syndrome.²⁶ The fetal brain seems to be more sensitive to CO during the late gestational stage.²⁷ However, at any stage of pregnancy, functional changes in psychomotor and mental development can appear. The abnormalities described in the literature are similar to those observed after other types of hypoxic injury to the central nervous system in the prenatal period. Since Breslau and Freund, several reported cases of fetal death have been published. Maternal death certainly leads to fetal death. In cases of maternal coma, fetal death is probable. However, maternal wellbeing is misleadingly reassuring. Fetal death can still occur

even in the presence of apparent maternal wellbeing.^{14 15 19 28} The severity of damage depends on the degree and duration of hypoxia or asphyxia. Death or permanent neurological damage can follow a single episode of hypoxia.

Fetal prognosis is difficult to estimate. The level of carboxyhaemoglobin has been proposed as a criterion but maternal symptoms seem to be the best indicators of fetal prognosis. The carboxyhaemoglobin level is only valuable in the confirmation of exposure to CO and has little correlation with the gravity of the outcome. Maternal carboxyhaemoglobin levels do not accurately reflect fetal haemoglobin or tissue levels.¹⁷ Kopelman *et al* report the case of a pregnant woman who had been exposed to exhaust fumes with a carboxyhaemoglobin level (after two hours) of 6.9%.²⁹ Her symptoms were headache, dyspnoea and dizziness. Six days later, at term, the infant presented signs of multiorgan injury resulting from in utero hypoxia. The fetus had been more severely affected than the mother. According to Norman *et al*, the maternal symptoms predict the wellbeing of the fetus.²⁵ When the mother suffers a period of unconsciousness or coma, the fetal outcome is poor, but when the severity of maternal toxicity is graded as mild the fetal outcome is generally good. On the other hand, if the mother dies, fetal death inevitably follows. Koren *et al*, using a classification by grades (table 1), observe that grades 1–2 have a good fetal outcome.³⁰ Grades 4–5 have a seriously adverse outcome and according to this author termination of pregnancy should be envisaged in these cases.

The treatment of acute CO poisoning during pregnancy begins by removing the victim from the environmental source of CO. Then 100% normobaric oxygen should be administered immediately and hyperbaric oxygen should be considered. The safety of hyperbaric oxygen administration for the fetus is still to be established. Some published studies suggest that hyperbaric oxygen administration entails a risk of abortion and fetal defects.^{31 32} In animal studies hyperbaric oxygen administration may be responsible for fetal reabsorption, abortion and teratogenesis. Prolonged administration of hyperbaric oxygen, as used in non-human studies, is clearly teratogenic.³³ However, Elkharrat *et al* maintain there is no evidence of fetal or obstetric morbidity if the following protocol is observed: a two hour session of hyperbaric oxygen administered at 2 ATA.⁷ In Koren's study, two cases of grade 4–5 intoxication resulted in a normal outcome after treatment with hyperbaric oxygen. Sporadic cases reveal no adverse effects on the fetus after hyperbaric oxygen administration.^{32 34 35}

It is admitted that fetal malformation and hypotrophy risks are, after hyperbaric oxygen, as significant as that of the general population.⁷ It has been suggested that exposure to hyperbaric oxygen during the first stage of pregnancy can induce abortion or premature labour.³² The only reported case of premature labour related to hyperbaric oxygen administration was probably fortuitous according to the authors.³²

Conclusion

Acute CO poisoning during pregnancy is comparatively uncommon, yet can result in fetal death and functional alterations or anatomical malformations in survivors. The level of maternal CO haemoglobin is a poor indicator of fetal toxicity. Despite maternal wellbeing, fetal morbidity or mortality can still occur. Hyperbaric oxygen seems to be the treatment of choice and all pregnant women suffering CO intoxication should be referred for hyperbaric oxygen therapy. It should be instituted as rapidly as possible, yet its efficacy is still theoretically possible long after initial exposure. In human protocol, the hyperoxic period is short and well tolerated by the fetus.

Funding: none.

Conflicts of interest: none.

Contributors

J L Greingor is guarantor of the paper.

- Office for National Statistics. *Mortality statistics cause: England and Wales*. London: HMSO, 1996. Series DH2 No23.
- Conseil Supérieur d'Hygiène Publique de France. *Diagnostic systématique de l'intoxication par le monoxyde de carbone* [Systematic detection of carbon monoxide intoxication] [In French]. *Revue du SAMU* 1994;3:125.
- Cobb N, Etzel RA. Unintentional carbon monoxide-related deaths in the United States, 1979 through 1988. *JAMA* 1991;266:659–63.
- Hardy KR, Thom SR. Pathophysiology and treatment of carbon monoxide poisoning. *J Toxicol Clin Toxicol* 1994;32:613–29.
- Breslau F. Intoxication zweier Schwangeren mit Halbleuchtgas Tod und vorzeitige Geburt eines Kindes. *Mösch Geburt Frauenkr* 1859;13:449–56.
- Freund MB. Accidental death of the 7 month foetus as a result of slight intoxication of the mother with carbon monoxide gas. *Mösch Geburt Frauenkr* 1859;14:31–3.
- Elkharat D, Raphael JC, Korach JM, et al. Acute carbon monoxide intoxication and hyperbaric oxygen in pregnancy. *Intens Care Med* 1991;17:289–92.
- Wattel F, Mathieu D, Nevier R, et al. L'intoxication au monoxyde de carbone [Carbon monoxide poisoning] [In French]. *Presse Med* 1996;25:1425–9.
- Hyperbaric Center Advisory Committee Emergency Medical Service, City of New York. A registry for carbon monoxide poisoning in New York city. *J Toxicol Clin Toxicol* 1988;26:419–41.
- Zhang J, Piantadosi CA. Mitochondrial oxidative stress after carbon monoxide hypoxia in the rat brain. *J Clin Invest* 1992;90:1193–9.
- Thom SR. Deshydrogenase conversion to oxidase and lipid peroxidation in brain after carbon monoxide poisoning. *J Appl Physiol* 1992;73:1584–9.
- Thom SR. Carbon monoxide mediated brain lipid peroxidation in the rat. *J Appl Physiol* 1990;68:997–1003.
- Ginsberg MD, Myers RE. Fetal brain damage following maternal carbon monoxide intoxication: an experimental study. *Acta Obstet Gynecol Scand* 1974;53:309–17.
- Caravati EM, Adams CJ, Joyce SM, et al. Fetal toxicity associated with maternal carbon monoxide poisoning. *Ann Emerg Med* 1988;17:714–17.
- Cramer CR. Fetal death due to accidental maternal carbon monoxide poisoning. *J Toxicol Clin Toxicol* 1982;19:297–301.
- Christensen P, Gronlund J, Carter AM. Placental gas exchange in the guinea pig: fetal blood gas tensions following the maternal oxygen capacity with carbon monoxide. *J Dev Physiol* 1986;8:1–9.
- Curtis GW, Algeri EJ, McBay AJ, et al. The transplacental diffusion of carbon monoxide. *Arch Pathol Lab Med* 1955;59:677–90.
- Margulies JL. Acute carbon monoxide poisoning during pregnancy. *Am J Emerg Med* 1986;4:516–19.
- Longo LD. The biological effects of carbon monoxide on the pregnant woman, fetus and newborn infant. *Am J Obstet Gynecol* 1977;129:69–103.
- Hill EP, Hill JR, Power GG, et al. Carbon monoxide exchanges between the human fetus and mother: a mathematical model. *Am J Physiol* 1977;232:H311–23.
- Kety SS, Schmidt CF. The effects of altered arterial tensions of carbon dioxide and oxygen cerebral blood flow and cerebral oxygen consumption of normal young men. *J Clin Invest* 1948;27:484–92.
- Lambertsen CJ, Kough RH, Cooper DY, et al. Oxygen toxicity : effects in man of oxygen inhalation at 1 and 3.5 atmospheres upon blood gas transport, cerebral circulation and cerebral metabolism. *J Appl Physiol* 1953;5:471–86.
- Thom SR. Antagonism of carbon monoxide-mediated brain lipid peroxidation by hyperbaric oxygen. *Toxicol Appl Pharmacol* 1990;105:340–4.
- Brown S, Piantadosi C. Recovery of energy metabolism in the rat brain after carbon monoxide poisoning. *J Clin Invest* 1992;89:666–72.
- Norman CA, Halton DM. Is carbon monoxide a workplace teratogen ? A review and evaluation of the literature. *Ann Occup Hyg* 1990;4:335–47.
- Courten W, Hennequin Y, Blum D, et al. Charge association in a neonate exposed in utero to carbon monoxide. *Birth Defects* 1996;1:407–12.
- Okeda R, Matsuo T, Kuroiwa T, et al. Experimental study on pathogenesis of the fetal brain damage by acute carbon monoxide intoxication of the pregnant mother. *Acta Neuropathol* 1986;69:244–52.
- Muller GL, Graham S. Intrauterine death of the foetus due to accidental carbon monoxide poisoning. *N Engl J Med* 1955;25:1075–8.
- Kopelman AE, Plaut TA. Fetal compromise caused by maternal carbon monoxide poisoning. *J Perinatol* 1998;18:74–7.
- Koren G, Sharav T, Pastuszak A, et al. A multicenter prospective study of fetal outcome following accidental carbon monoxide poisoning in pregnancy. *Reprod Toxicol* 1991;5:397–403.
- Grote W, Wagner WD. Malformations in rabbit embryos after hyperbaric oxygenation. *Klin Wochenschr* 1973;51:248–50.
- Hollander DI, Nagey DA, Welch R, et al. Hyperbaric oxygen therapy for the treatment of acute carbon monoxide poisoning in pregnancy. A case report. *J Reprod Med* 1987;32:615–17.
- Miller PD, Telford IR, Haas GR. Effect of hyperbaric oxygen on cardiogenesis in the rat. *Biol Neonate* 1971;17:44–52.
- Van Hoessen KB, Camporesi EM, Moon RE, et al. Should hyperbaric oxygen be used to treat the pregnant patient for acute carbon monoxide poisoning ? A case report and literature review. *JAMA* 1989;7:1039–43.
- Ledingham IM, McBride TI, Jennett WB, et al. Fetal brain damage associated with cardiomyopathy of pregnancy with notes on caesarean section in a hyperbaric chamber. *BMJ* 1968;4:285–7.