CHILDHOOD SEIZURE GUIDELINE

An evidence and consensus based guideline for the management of a child after a seizure

K Armon, T Stephenson, R MacFaul, P Hemingway, U Werneke, S Smith

Emerg Med J 2003;20:13–20

Structured in the format recommended by Hayward *et al*¹ for guideline reports.

Objective: An evidence and consensus based guideline for the management of the child who presents to hospital having had a seizure. It does not deal with the child who is still seizing. The guideline is intended for use by junior doctors, and was developed for this common problem (5% of all paediatric medical attenders) where variation in practice occurs.

Options: Assessment, investigations (biochemistry, lumbar puncture, serum anticonvulsant levels, EEG in particular), and/or admission are examined.

Outcomes: The guideline aims to direct junior doctors in recognising those children who are at higher risk of serious intracranial pathology including infection, and conversely to recognise those children at low risk who are safe to go home.

Evidence: A systematic review of the literature was performed. Articles were identified using the electronic data bases Medline (from 1966 to June 1998), Embase (from 1980 to June 1998) and Cochrane (to June 1998), and selected if they investigated the specified clinical question. Personal reviews were excluded. Selected articles were appraised, graded, and synthesised qualitatively. Statements of recommendation were made.

Consensus: An anonymous, postal Delphi consensus development was used. A national panel of 30 medical and nursing staff regularly caring for these children were asked to grade their agreement with the statements generated. They were sent the relevant original publications, the appraisals, and literature review. On the second and third rounds they were asked whether they wished to re-grade their agreement in the light of other panellists' responses. Consensus was defined as 83% of panellists agreeing with the statement.

Recommendations in brief: For afebrile seizures all children should have their blood pressure recorded, but no other investigations are routine although a seizing or somnolent child should have blood glucose measured; all children under 1 year should be admitted. For seizures with fever, clinical signs indicating the need to treat as meningitis are given. Children should be admitted if they are under 18 months old, have had a complex seizure, or after pretreatment with antibiotics.

Validation: The guideline has undergone implementation and evaluation in a paediatric accident and emergency department, the results of which will be published separately. Only one alteration was made to the guideline as a result of this validation process, which is included here.

See end of article for authors' affiliations

Correspondence to:
Dr K Armon, The Old
Parsonage, Norwich Road,
Long Stratton, Norwich
NR15 2PX, UK;
armon@ukonline.co.uk

Accepted for publication
10 May 2002

Seizures are common in children. Three per cent of all children 6 months to 5 years have a febrile seizure, and 5% of all medical attendances to the accident and emergency department are attributable to seizure.² Depending on the hospital attended and the clinician seen, about 70% of these children are admitted and undergo varying degrees of investigation³ and length of stay in hospital. The Royal College of Paediatrics and Child Health recommends that priority for guideline development be given to aspects of health care that are common, expensive or serious, and where variations in practice occur.⁴ Thus the management of childhood seizure is appropriate for guideline development.

Potential benefits (improved health outcome, consistency of care, patient information) and harms (priorities and recommendations flawed) of clinical guidelines have been well documented.⁵ We developed the guideline with the following goals: (1) to improve the process and outcome of care for children attending our accident and emergency department with a seizure; (2) to promote consistency of care so that patients with similar clinical problems would be managed in the same way; (3) to inform, educate, and improve the clinical decision making of the junior clinicians who see most of these children initially.²

SCOPE OF THE GUIDELINE

The guideline deals with children who have had a seizure as a presenting problem, rather than a defined diagnosis as staff

need guidance on determining diagnosis before deciding on management.⁶ The emergency management of the seizing child is not covered as it is dealt with clearly in accessible texts.⁷ We present a summary version of the full guideline, which can be obtained from the corresponding author electronically, and to which reference should be made for clarification or further information. The development group assumes that healthcare professionals will use general medical knowledge and clinical judgement in applying the recommendations in this document to the management of individual patients.

METHOD

We chose an algorithm format, where the recommended course of action at each stage depends critically on the available information.⁸ This has been shown to be successful in education.⁹

The guideline concentrates on the management of the child up to the point at which the decision is made to admit to hospital or discharge home. As such, the primary options considered are the important features on history and examination directing the clinician to a particular cause and the investigations required for diagnosis and initial management. Key clinical questions concerning the management of the child presenting with seizure drove our search strategy. Specific

questions, type of question and specific search terms (as mesh headings and text words) were defined.

We used electronic databases (Cochrane library (to March 1998); Medline (1966 to March 1998); Embase (1980 to March 1998), confined to humans aged 0–16 and English language), and citations in references found. The references generated were sifted for relevance to the clinical questions by their titles and abstracts. Inclusion criteria were: articles that investigated the clinical questions identified; scientific literature reviews; review or clinical guideline written by a national body; large well designed clinical trials (RCT, matched case control, cohort). Non-experimental, descriptive, or opinion based studies were excluded. Some retrospective cohort studies were included in the absence of stronger evidence and were graded accordingly. Articles were appraised using a pre-designed proforma and graded using a standard scale (appendix 1). Statements of recommended management were made based on this literature.

The statements were subject to consensus development by means of a postal Delphi process. Panellists selected were drawn from the United Kingdom, represented practice in both urban and rural settings and were clinicians who would be involved in management of a child after presentation at hospital. Eighty medical and nursing staff were invited, of whom 30 took part (panel composition in appendix 2). They rated their level of agreement on a 1 to 9 Likert scale and were asked to comment. Consensus agreement was defined as 83% of panellists rating 7 to 9. Two rounds of feedback and further rating were made. Those statements that gained consensus are included in the guideline.

THE GUIDELINE

The final guideline is in the form of an algorithm (flow diagram or decision tree) and is shown in figure 1, with integral tables 1 to 5. Each box is numbered, and key decision points are allocated a letter, with recommendations explained in the text. A brief discussion concerning the evidence base is given, followed by the consensus guideline statements in italics, with their level of evidence and grade of recommendation.

Throughout, the word “admit” is defined as follows: any admission to a paediatric facility with paediatric trained staff for observation, further investigation, and management regardless of the expected length of stay.

Children are managed differently depending on whether they are thought to have had a febrile or an afebrile seizure. The upper part of the algorithm deals with afebrile seizures, whereas the middle and bottom parts deal with febrile seizures. Crucial to this division is the definition of febrile seizure.

A Definition of fever in febrile seizure

There is no compelling evidence in the literature concerning the level of fever required to diagnose febrile seizure. Definition of fever depends on the method of measurement and these guidelines should be adapted to a locally preferred method. An axillary temperature of $>38^{\circ}\text{C}$ was proposed in the first Delphi round^{10–12} (Vb,D) but consensus was not achieved. In the second round the Likert scale was modified such that actual temperatures (between 37.0°C and 38.6°C in increments of 0.1 of a degree) were given. Panellists were asked to “place one cross only in the temperature box that corresponds to the lowest recorded axillary temperature that would lead you to make a preliminary diagnosis of seizure with fever”. The median and modal values were 37.8°C with a range of 37.7°C to 38.0°C and interquartile range all within box 37.8°C .

During the validation study it was clear that a simple temperature cut off was not practical and therefore the statement was modified to allow diagnosis of “febrile seizure” if the history and examination was indicative. This decision was in keeping with the definition developed at a consensus conference at the National Institutes of Health in 1980.

RECOMMENDATION on defining febrile seizure:

- *The temperature level to define fever with seizure is above 37.8°C .*
- *Modification during validation: “However, if the clinical history and examination is indicative of a febrile seizure (seizure associated with fever but without evidence of intracranial infection or a defined cause) the patient should be managed as such”.*

(level of evidence Vb, Delphi panel based, no consensus reached, precise wording decided during validation study, grade D recommendation)

AFEBRILE SEIZURES

B Determining the aetiology of a child presenting with an afebrile seizure

RECOMMENDATION on most likely causes of an afebrile seizure:

Table 1 (based on Smith *et al*¹³ observed diagnoses in children presenting to an A&E department over a one year period with a first seizure) for differential aetiology of an afebrile seizure.

(Based on Level Vb evidence and Delphi consensus, grade D recommendation)

C Diagnostic tests for the child presenting with an afebrile seizure

Studies on afebrile seizures have found no routine investigations to be of consistent value (in children^{13 14} and adults^{15 16}). Unusual historical or examination features should direct investigations. Studies on the positive yield of EEG were done in adults only, where it is not a useful diagnostic test after a first simple seizure.^{15 16} Children should have systemic hypertension presenting as a seizure excluded by sphygmomanometry.

RECOMMENDATION on investigations for afebrile seizure:

- *All children presenting with an afebrile seizure should have their blood pressure measured at the time of presentation.*
- *A finger prick blood glucose should be performed if a child is still seizing or not fully alert.*
- *No other investigations (that is, other than blood pressure and blood sugar) are routinely indicated in a child over 1 year of age with a simple afebrile seizure, who does not fulfil the criteria for observation or admission.*
- *There is no need for an EEG after a first simple afebrile seizure.*

(The above based on Level Vb evidence and Delphi consensus, grade D recommendation)

D Need for admission in children with afebrile seizures

Seizures in early infancy are often “symptomatic” (that is, secondary to underlying pathology) and therefore investigation and observation are essential¹⁷ (Vb,D). No other literature was found investigating this issue. Initial statements were derived from text book recommendation, and modified by consensus.

RECOMMENDATION concerning admission in afebrile seizure:

After a first afebrile seizure, children conforming to the stated criteria (table 2) should be admitted to an acute paediatric facility for observation and further investigation.

(Level Vb evidence and Delphi consensus, grade D recommendation)

FEBRILE SEIZURES

E Febrile seizure, risk of meningitis, and clinical features

In children admitted to hospital there are several studies of variable validity looking at the probability of meningitis (table 6). The signs included in the statement below were found to indicate an increased risk of meningitis in a child with seizure and fever^{18 19} (III, C):

Figure 1 Seizure guideline algorithm.

Table 1 List of causes of an afebrile seizure in a child presenting to the accident and emergency department

Type of seizure (% of non-febrile seizures)	Cause
Isolated seizure (25%)	No cause found
Epilepsy (30%)	Idiopathic, complex partial, Rolandic, etc.
Symptomatic seizure (30%)	CNS structural abnormalities (congenital, prenatal or perinatal, shunt obstruction)
	Intracranial infection (bacterial/viral, diffuse/localised)
	Ingestion (deliberate, accidental)
	Trauma (head injury, non accidental injury)
	Tumour
	Intracranial haemorrhage
	Hypertension
	Metabolic (low glucose, calcium, magnesium, high and low sodium, amino and organic acidurias, etc)
	Anoxic (breath holding attacks, respiratory pathology)
	Neurocutaneous syndromes
	CNS degenerative diseases
Neonatal/early infant seizure (<3 months) (15%)	In addition to the above causes:
	Hypoxic ischaemic encephalopathy (from birth)
	CNS infections (acute and congenital)
	Fifth day fits
	Drug withdrawal
	Pyridoxine dependency
Other diagnoses (not true seizures)	Reflex anoxic "seizures" syncope, arrhythmias, pseudoseizures

Table 2 Criteria for admission of a child with a first afebrile seizure to an acute paediatric facility

Category	Criteria/signs
Age	Less than 1 year
Neurology	Glasgow coma scale (or equivalent) <15 (>1 hour post fit)
	New neurological signs
Raised intracranial pressure	Papilloedema, tense fontanelle
Generally unwell	Irritable, disinterested, vomiting
Meningism	Kernig's positive, photophobia, neck stiffness
Complex seizure	Prolonged (>15 minutes), focal, recurrent
Signs of aspiration	Respiratory distress, need for oxygen, chest signs.
High parent or carer anxiety	Parents/carers do not feel happy to take the child home

Table 3 Common causes of fever in children presenting with fever and seizure

Cause for fever
Viral infection (for example, upper respiratory tract infection, non-specific viral illness, roseola, chicken pox and other exanthema, etc)
Otitis media
Tonsillitis
Urinary tract infection
Gastroenteritis
Lower respiratory tract infection
Meningitis
Post-immunisation
Post-ictal fever (only likely after generalised seizure of >10 min)

NB Viral infection, otitis media, and tonsillitis account for 85%–90% with the others making up 10%–15% of all causes.

RECOMMENDATION on clinical features indicating meningitis:

- A child who presents with seizure and fever and has any of the following on history or examination should be treated as having meningitis until proven otherwise: drowsy pre-seizure, neck stiffness, petechial rash, bulging fontanelle, a Glasgow Coma Scale of <15 (more than one hour post seizure).

(Level III evidence and Delphi consensus, grade C recommendation)

F Complex febrile seizures

The literature suggests that complex febrile convulsions (prolonged seizure (>15 min), multiple seizures in 24 hours and focal features) increase the risk of further seizures and epilepsy^{20–23} (III,C) and are also predictive of CNS

infection^{18 19} (III,C)^{24 25} (Vb, D). The risk of bacterial meningitis in children presenting with fever and seizure is about 3%^{18 19 25–27} (III,C) and in a complex seizure, 9%.

After the first Delphi round it was agreed that children with complex seizures should be admitted to hospital. Two alternative statements were then given suggesting that children with a complex seizure should either have a lumbar puncture on admission or be reviewed in two hours. The second option gained consensus.

RECOMMENDATION in children with complex febrile seizures

- A child presenting with a complex febrile seizure (defined above) with no clinical signs of meningitis (section E) should be observed closely and reviewed within two hours by a paediatrician of at least registrar level to decide on need for lumbar puncture.

Table 4 Contraindications to lumbar puncture

Category	Criteria/signs
Drowsiness or impaired consciousness	Falling conscious level, Glasgow coma scale of <13
Signs of septicaemic shock	Poor perfusion, low BP, tachycardia
Clinical diagnosis of invasive meningococcal disease	Rapid onset illness, typical haemorrhagic rash
Signs of raised intracranial pressure	Papilloedema, coma, abnormal posturing, abnormal pupillary responses, high BP, low pulse
Focal neurological signs	On clinical examination of cranial and peripheral nerves
Bleeding tendency	Thrombocytopenic rash or excessive bruising—check platelets and clotting screen
	Known haemophilic—consult haematologist

Table 5 Prognosis of febrile seizures

Risk	Percentage
Population risk of febrile seizure	2.7 to 3.1
Risk of recurrence of febrile seizure after first seizure	27 to 32
Risk of epilepsy after simple febrile seizures	1.5 to 2.4
Risk of epilepsy after complex febrile seizures (prolonged >15 minutes, focal, multiple in 24 hours)	4.1 to 6.3

(Implied by Level III evidence. Delphi consensus, grade C recommendation)

RECOMMENDATION on contraindications to lumbar puncture:

Table 4 shows the contraindications to LP.^{28 29}

(Level Vb evidence and Delphi consensus, grade D recommendation)

G Age and risk of meningitis in febrile seizures

The population risk of meningitis (with or without seizure) based on figures from 1980 to 1990³⁰ is highest in those aged 1 to 11 months (115/100 000 versus 28.5/100 000 for 1–5 years and 2.8/100 000 for 5–16 years). This rate has decreased since the introduction of Hib vaccine, but remains highest in the under 1 year olds. The statement that children under 1 years of age (often extended to 18 months) can have meningitis without displaying any of the classic signs seems to be based on clinical experience and some early published case reports but is not born out by formal studies.^{18 19 24}

Consensus views from the RCPCH (then BPA)¹¹ and The American Academy of Pediatrics³¹ (AAP) suggest lumbar puncture under 18 months (probably) and almost certainly under 12 months. No view on admission is given.

RECOMMENDATION on age for admission and lumbar puncture in febrile seizure:

- All children under 12 months with a first simple febrile seizure should be admitted.
- Children under 12 months with a simple febrile seizure should have a lumbar puncture unless an experienced paediatrician (at least paediatric registrar) has decided not to lumbar puncture and will review within two hours.
- Children 12 to 18 months with a simple febrile seizure should be admitted to an acute paediatric facility for a period of observation (at least two hours).
- Maintain a low threshold for an lumbar puncture in children 12 to 18 months with a simple febrile seizure.

(Level Va evidence and Delphi consensus, grade D recommendation)

H Antibiotics and meningitis risk in febrile seizures

There is little evidence to support the statement that prior antibiotic treatment masks the signs of meningitis.^{18 19 24} However, the AAP³¹ suggest that lumbar puncture be strongly considered in these children. Three options were given to the panel after the first round of comments (i) admit and observe, (ii) lumbar puncture, or (iii) discharge if source of infection is evident and does not require hospital treatment. The first gained consensus.

RECOMMENDATION on management of the child who has already received antibiotics:

- Those children with a simple febrile seizure, >1 year of age and with no serious historical or examination findings indicating meningitis who have had prior antibiotic treatment should be admitted to an acute paediatric facility for a period of observation (at least two hours)

(Level Va evidence and Delphi consensus, grade D recommendation)

I Causes of fever in children presenting with febrile seizure

RECOMMENDATION for differential diagnosis of fever in febrile seizure:

Table 3 gained consensus and was based on literature summarised in table 7.^{13 26 32 33}

(Level Vb evidence and Delphi consensus, grade D recommendation)

J Management of the child with febrile seizure and no focus of infection

There was no published evidence on this issue. The need for a good urine sample collected without contamination³⁴ was agreed in the first round.

RECOMMENDATION on management of child with no focus of infection:

- A child who has had a simple febrile seizure where no source for infection has been found clinically, should have a urine sample (clean catch, SPA or catheter specimen) taken for microscopy and culture.

After first round comments, two statements were given as alternative management plans concerning admission or discharge, and both achieved consensus, at the same level of agreement.

- Children with no focus for infection can be admitted for a short period of observation (minimum two hours) OR
- Children with no focus for infection can be discharged home if the child looks well, parents/carers have ready access to health care if required and they are happy with this decision.

(All the above based on Delphi consensus only, no published evidence)

K Prognosis after a first febrile seizure

RECOMMENDATION for prognosis of febrile seizure:

Information concerning prognosis is given in table 5 of the guideline^{21 35} and is available to inform the clinician's discussion with parents.

(Level III evidence and Delphi consensus, grade C recommendation)

Table 6 Published reports of the probability of bacterial meningitis in children with fever and a seizure

Date	Author	Number in study	Number with lumbar puncture	Number with bacterial meningitis	Number with viral meningitis
1980	Lorber ²⁶	452	304 (67%)	14 (3.1%)	1
1981	Jaffe ²⁵	562	All*	6 (1%)	17 (3%)
1983	Jaffe ¹⁸	254	241 (95%)	13 (5.1%) (In 2 CSF was sterile)	?2
1990	McIntyre ²⁷	307	154 (50%)	2 (0.6%) (1 diagnosed later on repeat lumbar puncture)	3
1992	Offringa ¹⁹	309	171 (55%)	21 (6.7%) (In 5 cases the CSF was sterile)	2

*Policy to lumbar puncture all cases, number of unsuccessful lumbar punctures not stated.

Table 7 Identified or attributed causes of fever in children presenting with a febrile seizure (%)

	Rutter (n=328)	Green/MacFaul (n=199)	Smith (n=116)	Lorber (n=452)
Viral infection (URTI and other)	45.7	64	32.7	57.9
Unknown	20.4	NA	20	23.6
Otitis media	8.5	18.5	28.4	9.5
Urinary tract infection	1.2	4	2.5	NR
Gastroenteritis	3.4	4	2.5	NR
Chest infection	3.9	5	0.8	?3
Tonsillitis	12	(included in URTI)	10.3	NR
Meningitis	1.8*	NA	NA	3.3
Other (chicken pox, mumps, exanthema, post-triple immunisation)	2.7	4	2.5	2.4

NR, not recorded; NA, not applicable. *These cases are reported as they were detected at a later stage, after the initial diagnosis of febrile seizure without CNS infection had been made.

L Follow up for the child with febrile seizure

There is no published evidence concerning the need for follow up of children discharged following a simple febrile seizure.

RECOMMENDATION on follow up for the child with febrile seizure:

- The Delphi panel agreed that an information sheet should be supplied (available from corresponding author) and follow up does not need to be arranged.
- Parents of children sent home from A&E with a diagnosis of first simple febrile seizure should be encouraged to contact their own GP or community nurse specialist if they feel they need further information or care.

(No published evidence, Delphi consensus).

RESEARCH QUESTIONS

During the development of this guideline several areas were identified where good research evidence was lacking. The following questions were identified.

- 1 Is measuring the anticonvulsant level routinely justified in a child who presents with seizure and who is already taking anticonvulsant medication? If not, in which children and on which anticonvulsants should blood levels be measured?
- 2 What is the true population risk for meningitis in a child presenting with fever and seizure?
- 3 In children who present with a first simple febrile seizure, is discharge to general practitioner care as good as in patient care for limitation of risks, cost effectiveness, patient and parent satisfaction and education?
- 4 In a child presenting with a febrile seizure and a cause for fever is suspected on clinical examination, is it cost effective to take a routine urine sample? In what age group and by what method?

DISCUSSION

A comprehensive guideline is presented for the management of the child who has had a seizure based on best evidence and formal consensus using a Delphi panel. Management for those presenting with an afebrile seizure differs from those with a seizure associated with fever. Consensus was not achieved in

Key recommendations

For afebrile seizures

- All children should have blood pressure recorded and a blood glucose taken if still seizing or not fully alert. Other investigations should be directed by history and clinical findings.
- Children under 1 year of age should be admitted to hospital for further observation and investigation.
- There is no need for an EEG in a first uncomplicated seizure.
- An outpatient appointment should be offered.

For a seizure with fever

- Admit and treat as meningitis children who were drowsy at home before the seizure, have a GCS<15 more than one hour after seizure, have neck stiffness, petechiae, or a bulging fontanelle.
- Admit children with complex seizures, those less than 18 months, and those who have had antibiotics. The child should have an experienced paediatric review (at least registrar) within two hours concerning need for lumbar puncture.
- Consider admission in those in which no focus for infection was found.
- Others may go home with adequate parental information, reassurance, and ready access to health care.

defining a temperature level below which afebrile seizure could be diagnosed. However, panelists' responses were tightly grouped from 37.7°C to 38.0°C. The reason for the lack of consensus may be that clinicians take into account many other factors in the history and examination when deciding on a diagnosis of "febrile seizure". The guideline therefore recommends that the cut off level be decided locally. During the validation study it was clear that clinicians wanted the freedom to diagnose "febrile seizure" if the history and examination were highly suggestive even if the temperature did not reach the required level. For this reason the guideline suggests that clinicians should use their clinical judgement.

Evidence from randomised trials was not available for constructing the original management statements. The highest

level of evidence was level III on which only two statements (of 42) were directly based and seven were based by implication. The highest level of evidence (grade I) is that of a systematic review of randomised controlled trials using meta-analysis, and grade II is a good single randomised controlled trial with enough power to detect a difference. However this type of evidence only applies to questions concerning treatment, which are in the minority in a presenting problem based guideline. Different types of studies answer questions concerning population risk and diagnostic tests (symptoms and signs and laboratory investigations). A multicentre randomised controlled trial could be designed for practical management issues such as need for admission and follow up, but measurable outcomes are difficult to define. Thus for many practical clinical questions good evidence is lacking and is unlikely to be available in the near future, which highlights the need for good consensus methods.

Delphi consensus enabled a guideline to be produced where good published evidence is sparse. The guideline reflects a consensus of peer practice in that country at that time. Clinicians expressed reservations in some areas despite consensus being achieved. For example, the panel agreed that a urine sample should be taken from a child with febrile seizure in whom another source of infection is suspected. Despite this consensus, 14 of the panellists made written comments that expressed reservations about such a policy. It is possible that the "safest" options achieve consensus using this process but clinicians may act differently when they are seeing an individual child and can take other information into account.

There is little research into the method of Delphi consensus development, including the influence of different panels of individuals on outcome. While the panellists in this study were "self selected" in that they agreed to take part, it is difficult to conceive of an alternative to this.

As there was no level I or II evidence, these recommendations should be treated as a guide only. They will need adaptation for each local unit, and clinicians must exercise their own judgement as to whether the guideline applies to the individual patient seen.

The guideline has been validated in a paediatric A&E department and we intend to publish the results of the

implementation study. Formal assessment of the costs of development of a presenting problem based guideline using Delphi consensus was not undertaken. We intend to evaluate costs during the development of the next guideline using the same methodology.

ACKNOWLEDGEMENTS

The following Delphi panelists for contributing a great deal of time and effort: H Allen (paediatric SpR), J H Baumer (paediatric consultant), R Bell (paediatric consultant), D Beverley (paediatric consultant), A Chambers (paediatric nurse), C Cramp (paediatric consultant), J Dawling (paediatric SpR), H Dixon (Paed. SpR), S Edees (paediatric consultant), M Everard (paediatric consultant), A Gregory (paediatric SpR), P Hardy (paediatric A&E SpR), S Hartland (paediatric A&E nurse), H Huynh (paediatric SpR), K Jackson (paediatric nurse), J Jenkins (paediatric consultant), B Lloyd (paediatric consultant), I Mecrow (paediatric consultant), J Moorcraft (paediatric consultant), R Morton (paediatric consultant), R Newton (paediatric neurologist), J Nixon (paediatric A&E nurse), K Palmer (paediatric consultant), A Raffles (paediatric consultant), C Simpson (paediatric nurse), C Smith (paediatric SpR), E Szondy (paediatric nurse), T Tinklin (paediatric consultant), C Upton (paediatric consultant), R Watkins (paediatric SpR), L Williams (paediatric A&E consultant). The authors thank Jeanette Taylor-Meek for her efficient and effective administration of the Delphi process

Contributors

Kate Armon planned and coordinated the development of the study, developed core ideas, performed the literature review, appraised the articles found, generated initial guideline statements, coordinated the Delphi process, collated and analysed the data, produced the final guideline, and wrote the paper. Terence Stephenson had the original idea for the research and oversaw its development, participated in literature appraisal, development of guideline statements, and interpretation of data, discussed core ideas, edited the paper, and will act as guarantor. Roddy MacFaul initiated the research, had original ideas and discussed its development, participated in developing guideline statements, interpretation of data, and edited the paper. Pippa Hemingway discussed core ideas and development of research and edited the paper. Ursula Werneke had original ideas concerning the Delphi process, discussed the development of the research, and edited the paper. Stephanie Smith discussed core ideas and development of the research and participated in the Delphi process.

Appendix 1 Levels of evidence and grade of recommendation

Level	Strength of evidence (Adapted from Muir Gray ⁶⁴)	Grade	Grade of recommendation (Cook <i>et al</i> ⁶⁷)
I	Strong evidence from at least one systematic review of multiple well designed randomised controlled trials	A	Supported by level I evidence and therefore highly recommended.
II	Strong evidence from at least one properly designed randomised controlled trial of appropriate size.	B	Supported by level II evidence, and therefore recommended
III	Evidence from well designed trials without randomisation, single group pre-post, cohort, time series or matched case-control studies	C	Supported by level III, evidence. Several potential clinical actions might be considered appropriate.
IV	Evidence from well designed non-experimental studies from more than one centre or research group	D	Supported by level IV and V evidence. The consensus route would have to be adopted.
Va	Opinions of respected authorities	D	
Vb	Clinical evidence, descriptive studies or reports of expert committees	D	

Appendix 2 Delphi panellists

Clinician	Number
Paediatric district general hospital consultant	11
Paediatric teaching hospital consultant	3
Paediatric specialist registrar/SHO post membership	4
Paediatric neurologist	1
Paediatric nurse	2
A&E nurse	2
A&E trainee	1
Paediatric A&E nurse	1
Paediatric A&E consultant	1

Authors' affiliations

K Armon, T Stephenson, P Hemingway, Academic Division of Child Health, Nottingham, UK

R MacFaul, Pinderfields General Hospital, Wakefield, UK

U Werneke, Maudsley Hospital, London, UK

S Smith, Queen's Medical Centre, Nottingham, UK

Funding: Children Nationwide Medical Research Fund.

Conflicts of interest: none.

REFERENCES

- 1 **Hayward R**, Wilson M, Tunis S, *et al*. More informative abstracts of articles describing clinical practice guidelines. *Ann Intern Med* 1993;**118**:731–7.
- 2 **Armon K**, Stephenson TJ, Gabriel V, *et al*. Determining the common medical presenting problems to an accident and emergency department. *Arch Dis Child* 2001;**84**:390–2.
- 3 **Sweeney A**, Gibbs J, Monteil F, *et al*. The management of febrile seizures in the Mersey Region. *Dev Med Child Neurol* 1996;**38**:578–84.
- 4 **Royal College of Paediatrics and Child Health**. *Standards for development of clinical guidelines in paediatrics and child health: role of the Royal College of Paediatrics and Child Health. Report of the quality of practice committee*. 1st edn. London: Royal College of Paediatrics and Child Health. 1998:19.
- 5 **Woolf SH**, Grol R, Hutchinson A, *et al*. Potential benefits, limitations, and harms of clinical guidelines. *BMJ* 1999;**318**:527–30.
- 6 **Hopkins A**. Some reservations about clinical guidelines. *Arch Dis Child* 1995;**72**:70–5.
- 7 **ALSG**. *Advanced paediatric life support; the practical approach*. 2nd edn. London: BMJ Publishing Group, 1997:306.
- 8 **Grimshaw JN**, Russell IT. Achieving health gain through clinical guidelines I: developing scientifically valid guidelines. *Qual Health Care* 1993;**2**:243–8.
- 9 **Margolis CZ**, Cook CD, Barak N, *et al*. Clinical algorithms teach paediatric decision making more effectively than prose. *Med Care* 1989;**27**:576–92.
- 10 **Sahib El-Radhi A**, Carrol JE. *Fever in paediatric practice*. Oxford: Blackwell Scientific, 1994.
- 11 **Addy DP**, Hopkins AP. Guidelines for the management of convulsions with fever. *BMJ* 1991;**303**:634–6.
- 12 **Baraff LJ**, Bass JW, Fleisher GR, *et al*. Practice guideline for the management of infants and children 0 to 36 months of age with fever without source. Agency for Health Care Policy and Research. *Ann Emerg Med* 1993;**22**:1198–210.
- 13 **Smith RA**, Martland T, Lowry MF. Children with seizures presenting to accident and emergency. *J Accid Emerg Med* 1996;**13**:54–8.
- 14 **Nypaver MM**, Reynolds SL, Tanz RR, *et al*. Emergency department laboratory evaluation of children with seizures: Dogma or dilemma? *Pediatr Emerg Care*. 1992;**8**:13–16.
- 15 **Turnbull TL**, Vanden Hoek TL, Howes DS. Utility of laboratory studies in the emergency department patient with a new-onset seizure. *Ann Emerg Med* 1990;**19**:373–7.
- 16 **Eisner RF**, Turnbull TL, Howes DS, *et al*. Efficacy of a 'standard' seizure work up in the emergency department. *Ann Emerg Med* 1986;**15**:69–75.
- 17 **Chevrie JJ**, Aicardi J. Convulsive disorders in the first year of life: neurological and mental outcome and mortality. *Epilepsia*, 1978;**19**:67–74.
- 18 **Joffe A**, McCormick M, DeAngelis C. Which children with febrile seizures need lumbar puncture? A decision analysis approach. *American Journal of Diseases of Children* 1983;**137**:1153–6.
- 19 **Offringa M**, Beishuizen A, Derksen-Lubsen G, *et al*. Seizures and fever: can we rule out meningitis on clinical grounds alone? *Clin Pediatr* 1992;**31**:514–22.
- 20 **Nelson KB**, Ellenburg JH. Predictors of epilepsy in children who have experienced febrile seizures. *N Engl J Med* 1976;**295**:1029–33.
- 21 **Annegers JF**, Hauser WA, Shirts SB, *et al*. Factors prognostic of unprovoked seizures after febrile convulsions. *N Engl J Med* 1987;**316**:493–8.
- 22 **Berg AT**, Shinnar S, Hauser WA, *et al*. A prospective study of recurrent febrile seizures. *N Engl J Med* 1992;**327**:1122–7.
- 23 **Verity CM**, Golding J. Risk of epilepsy after febrile convulsions: a national cohort study. *BMJ* 1991;**303**:1373–6.
- 24 **Green SM**, Rothrock SG, Clem KJ, *et al*. Can seizures be the sole manifestation of meningitis in febrile children? *Pediatrics*, 1993;**92**:527–34.
- 25 **Jaffe M**, Bar-Joseph G, Tirosh E. Fever and convulsions—indications for laboratory investigations. *Pediatrics* 1981;**67**:729–31.
- 26 **Lorber J**, Sunderland R. Lumbar puncture in children with convulsions associated with fever. *Lancet* 1980;*ii*:785–6.
- 27 **McIntyre PB**, Gray SV, Vance JC. Unsuspected bacterial infections in febrile convulsions. *Med J Aust* 1990;**152**:183–6.
- 28 **Slack J**. Deaths from meningococcal infection in England and Wales in 1978. *J R Coll Physicians Lond* 1982;**16**:40–4.
- 29 **Wylie P**, Stevens D, Drake W, *et al*. Epidemiology and clinical management of meningococcal disease in West Gloucestershire: retrospective population based study. *BMJ* 1997;**315**:774–9.
- 30 **Fortnum HM**, Davis AC. Epidemiology of bacterial meningitis. *Arch Dis Child* 1993;**68**:763–7.
- 31 **Bergman DA**, Baltz RD, Cooley JR, *et al*. Practice parameter: the neurodiagnostic evaluation of the child with a first simple febrile seizure. *Pediatrics* 1996;**97**:769–72.
- 32 **Green AL**, MacFaul R. Duration of admission for febrile convulsions. *Arch Dis Child* 1985;**60**:1182–4.
- 33 **Rutter N**, Smales OR. Role of routine investigations in children presenting with their first febrile convulsion. *Arch Dis Child* 1977;**52**:188–91.
- 34 **Stephenson T**, O'Callahan. *Pocket paediatrics*. 1st edn. London: Churchill Livingstone, 1992:394.
- 35 **Ellenberg JH**, Nelson KB. Sample selection and the natural history of disease. Studies of febrile seizures. *JAMA* 1980;**243**:1337–40.
- 36 **Muir Gray JA**. *Evidence-based healthcare*. 1st edn. London: Churchill Livingstone, 1997.
- 37 **Cook D**, Guyatt G, Laupacis A, *et al*. Rules of evidence and clinical recommendations on the use of antithrombotic agents. *Chest* 1992;**102**:305–11S.