EDITORIAL

This autumn should see the launch of the national 'Save a life' campaign, here in the UK. The British Broadcasting Corporation has recorded a special series of short television programmes to be broadcast at peak viewing times. Their aim is to provide a brief introduction to 'first-aid' and then to refer the viewer to a local centre for a 2-hour demonstration of the essential life-saving techniques. The fact that such an undertaking has even been considered is a major step forward in public education. There is no doubt that the mortality from acute myocardial infarction, in particular, can be significantly reduced when the public become involved with early resuscitation (Cummins & Eisenberg, 1985) and all of us involved in the care of the acutely ill must give the campaign all the support we can.

However, the fact that this campaign *needs* to be considered must beg some rather delicate questions. There are long-established voluntary-aid societies who enjoy a high degree of public esteem. If they have run first-aid courses for years, then why is there a problem? There can be only two answers: either not enough people have attended or those who have have been taught the wrong things. Is the unattractiveness for the public in the subject to be taught, the way it is taught or the way it is perceived it might be taught? The response to the campaign may shed a little light on this.

The coming together of such seemingly disparate bodies as the St John Ambulance, the Casualty Surgeons Association and a television producer, already seems to have catalysed an early reaction. It has, at last, been accepted that most people will attend only once and then only for a short time. Teachers must capitalise on this brief opportunity and ensure that only those techniques that will actually save life are acquired. Time spent on anything else is time wasted.

It is, of course, not only the voluntary-aid societies that need to audit their performance in this field. What has the medical profession as a whole been doing? Not a lot. Recent adverse publicity about the inadequacies of junior doctors in resuscitation may have shocked the public. It should not have shocked the doctors. For too long the teaching of even the most basic of life-saving skills has been left for the learner to acquire almost by osmosis. Unfortunately, information has passed from a concentrated pool of isolated opinion to a dilute lake of patient-contact across a semipermeable membrane of indifference.

First-aid should be taught only by those who practise it—now. The involvement of specialists who daily resuscitate the critically ill should, at last, remove spurious debate based on idle theory and replace it with the results of (recent) personal experience. Perhaps then such irrelevancies as the 'Holger-Neilsen' and 'Silvester' methods (St John Ambulance *et al.*, 1982) will disappear from the front line of first-aid teaching, and less time will be spent in such fatuous arguments as whether the fingers should or shouldn't be crossed in chest massage.

Science has shown that the quality of the resuscitative attempts is far less important than the fact that an attempt was made. Perhaps, at last, we will heed this lesson and

more time will be spent in ensuring that as many people as possible can at least do something.

REFERENCES

- Cummins R. O. & Eisenberg M. S. (1985) Cardiopulmonary resuscitation—American style. *British Medical Journal* **291**, 1401–3.
- St John Ambulance, St Andrew's Ambulance Association & British Red Cross Society (1982) *First-Aid Manual*, pp. 30–2. London, Dorling Kindersley Ltd.

A. D. REDMOND

Editor