GUEST EDITORIAL

Malaria

In excess of 18000 cases of malaria were reported to the Malarial Reference Laboratory over the decade ending in 1986, an incidence rise of 51% occurring in that time. Cases of *Plasmodium falciparum* malaria (with a case mortality ratio of 1.3%) increased more rapidly than those due to *P. vivax* malaria. Vivax is hardly ever fatal but can produce severe disease with notable systemic toxicity, and occasional falciparum-like manifestations such as blackwater fever and cerebral symptoms. There are three aspects of malaria which are important to address:

- (a) appropriate chemo-prophylaxis and other preventative measures,
- (b) early diagnosis, and
- (c) appropriate management of confirmed or suspected cases.

General preventative measures such as use of insect repellents, mosquito nets, wearing of long-sleeved shirts and long-legged trousers are coming into vogue once again. Chemo-prophylaxis has become progressively more complex with the increasing incidence of drug resistance. Chloroquine resistant *P. falciparum* malaria has now spread from East Africa westwards, cases being currently reported from as far west as The Gambia. In addition, resistant malaria is reported from India and Pakistan. This trend is further complicated by the not insignificant frequency of toxic side effects from anti-malarials which include severe skin reactions (Fansidar), agranulocytosis (Malo-prim, amodiaquine) and hepatitis (amodiaquine). Fatalities associated with the use of these drugs are of the order of 1:10000–1:20000. Doxycycline, mefloquine and metakelfin are three relatively recent additions to the prophylactic armamentarium but resistance to some of these such as mefloquine is of great concern. There are additional problems arising from the need for prophylaxis during pregnancy, in neonates and immune incompetent individuals. A sporozoite vaccine is advanced in trials.

Several studies have substantiated the concern that specialists have regarding the poor standards of antimalarial chemo-prophylaxis: in the UK 93% of patients with vivax malaria took no prophylaxis; only 48% of people who had sought pre-travel advice from a malarial reference authority complied with the advice given; a substantial number of general practitioners had important defects in their understanding of malarial prophylaxis. In view of the complex and changing situation, advice from an infectious diseases or tropical medicine unit should always be sought if there is any question or uncertainty about antimalarial prophylaxis.

Malaria should always be excluded in *any* ill patient who has been resident in or passing through (and that includes transit stops at an airport) a malarial area within the last 5 years or so, even in the absence of fever and even if there is a history of apparently appropriate antimalarial drug prophylaxis. Malaria is also more likely to be the cause of illness in a patient who returns from an endemic area of viral haemorrhagic fever. It

should not be forgotten that congenital malaria can occur from mothers who had last travelled to malarial areas up to 1–2 years previously. The neonate may present with failure to thrive, jaundice, hepatosplenomegaly or fever.

Symptoms of malaria can be non-specific or can mimic other illnesses. Because of the multi-system nature of the disease, *P. falciparum* malaria has been mistaken for viral illness, meningo-encephalitis, pneumonia, hepatitis, gastro-entero-colitis and pyelonephritis. Failure to diagnose correctly can prove fatal. The hypnozoite liver form of vivax or ovale malaria, and dormant blood forms of falciparum mean that some patients occasionally have an incubation period of up to 2 years, although most present within a few weeks or months of arrival in the UK. Remember also that relapse may occur for several years after an initial attack of vivax or ovale. *Plasmodium malariae* can occur for decades afterwards. The well-described cold, hot and sweating stages are not always present, particularly in *P. falciparum* malaria, and fever may be absent on occasions. The mainstay of diagnosis is to examine at least three serial thick and thin blood films. These should be taken over a period of 24–48 h and examined by a competent trained person. It should be stressed that inability to detect parasites in the peripheral blood film does not exclude the diagnosis of malaria. In *P. falciparum* malaria, overwhelming infection can occur with scanty falciparum peripheral blood parasitaemia due to the deep organ requirement for schizogony of this species. If in doubt, treatment for severe *P. falciparum* malaria should always be instigated in the light of the travel history and clinical findings, for example in the confused or disorientated patient from Central Africa in whom other diagnoses have been provisionally excluded. As a corollary, proven malaria should not be assumed always to be the only cause of fever in a particular patient: multiple infections do occur. Mixed malarial infections also occur. There have been notable advances in serological diagnosis, including indirect fluorescent antibody tests, ELISA and immunoradiometric assay. Genetically engineered DNA gene probe techniques are also available for the diagnosis but, as yet, these are not practicably applicable in the UK setting where examination of the blood film is still more sensitive and specific.

There are two broad issues in management: the patient should receive appropriate specific drug therapy and appropriate additional supportive therapy. Oral chloroquine in a dose of 1500–1800 mg given over 48 h will cover most infections seen in the UK. Intravenous chloroquine is extremely hazardous and is rarely necessary. Quinine should always be given to: (a) a severely ill patient with cerebral signs or deteriorating despite other therapy, (b) a patient with proven *P. falciparum* malaria acquired from an area known to be associated with chloroquine resistance, and (c) a patient with peripheral parasitaemia in excess of 10%.

If quinine is indicated and the clinical condition permits, it can be given orally (10 mg per kg every 8 h for 3 days—this rapidly reduces parasitaemia. Fansidar (three tablets) given immediately after the final dose of quinine will effect a radical cure. For the patient who is more ill, or is vomiting, IV quinine in the same dosage is indicated, switching to the oral route as soon as clinical improvement permits. For patients who come from Thailand, an initial dose of 20 mg per kg is recommended; occasionally quinine resistance has been described: quinidine is indicated in such cases. Quinine and quinidine are toxic drugs—apart from cinchonism, tinnitus, vertigo, tremor, blurred vision etc.—cardiotoxicity with hypotension and fatal arrhythmias is possible and

cardiac monitoring is indicated when they are given intravenously. Hypoglycaemia is a definite complication of malarial treatment. It is related partly to the stimulatory action of quinine on beta islet cells, producing hyperinsulinaemia; to the increased metabolic requirements of the parasites and, finally, to impairment of hepatic gluconeogenesis. It is particularly common in pregnancy, the severely ill, African children and adults. The overall incidence is between 8 and 42%. Hypoglycaemia can be mistaken for a worsening of malaria *per se* but is, in theory, reversible by IV dextrose. The hypoglycaemia may prove refractory or may actually be worsened by IV dextrose.

There are a number of other practical complications. Pulmonary oedema, possibly as a consequence of increased pulmonary capillary permeability, may prove fatal, particularly with over-enthusiastic administration of fluids. Prevention is the aim, with judicious fluid balance, and maintenance of adequate cardiac output using a low right heart pressure. Definitive treatment is IPPV + PEEP as for ARDS. Secondary bacterial infection, particularly with gram negative organisms occurs, for example, secondary to catheterization or directly through damaged gut and demands careful surveillance, specific diagnosis and appropriate antimicrobial therapy, together with plasma expanders, inotropic agents and physiotherapy. Acute tubular necrosis requires dialysis. Since 1982, exogenous corticosteroids have been shown to be deleterious in cerebral malaria. The normal CSF opening pressure and lack of CT head scan evidence of cerebral oedema in many patients with cerebral malaria adds further support to the approach which contra-indicates the use of steroids in patients with cerebral malaria. In the presence of high parasitaemia (greater than 50% in the peripheral blood film) or greater than 10% accompanied by significant major organ complications, exchange blood transfusion between quinine infusions using fresh whole blood (usually between 5 and 10 units) and under CVP monitoring, is an effective and rapid means of reducing the peripheral parasite count. It may reverse coma and prove life saving. Severe haemolysis (including blackwater fever), coagulopathies such as DIC and other haematological abnormalities, particularly thrombocytopaenia sometimes occur. Convulsions, pneumonia, myocarditis, hypotension and splenic rupture should also be considered in the severely ill.

Malaria which occurs in two particular groups demands a slightly different emphasis. It is often said that malaria in a semi-immune person (with a history of repeated infections) will respond to lower doses of a drug regime which would be ineffective in a non-immune. It is probably wise not to adopt this approach in a patient who has recently arrived in the UK, since immunity can diminish rapidly within 12 months after departure and ineffective drugs may select resistant organisms. Malaria in pregnancy is usually more severe. There is a higher incidence of abortion, premature labour, maternal anaemia and congenital infection. In addition, there are fears (some unfounded) that the use of prophylactic drugs may prove harmful to the foetus. The traditional advice that travel to a malarial area during pregnancy should be avoided is still correct. There should be no hesitation in using quinine in treating malaria in pregnancy despite the theoretical risk of precipitating labour.

Follow-up of patients recovered from malaria is not usually necessary but may be indicated in a few instances. For example, after the initial treatment of malaria additional treatment is necessary in the case of vivax and ovale to eliminate hepatic parasites which may later produce relapse. Primaquine, 15 mg daily for a fortnight (or

25 mg daily for patients from South-East Asia or the Pacific due to resistant organisms) is given after the initial treatment. If G6PDH level are low, Primaquine is given in a smaller dose for a longer period and the blood screened for haemolysis. Primaquine is sometimes erroneously given to patients with *Plasmodium malariae* infections—however, there are no hypnozoite forms. Persistence is due to erythrocytic forms and each relapse, which can occur up to decades later, should be treated with chloroquine. An additional complication of *Plasmodium malariae* is immune-complex-mediated nephrotic syndrome which is unresponsive to antimalarials but occasionally responds to immune suppressive treatment. Finally, a few people develop a chronic abnormal immune response to any one of the Plasmodium species producing symptomatic splenomegaly, debility and haemolysis. Long-term antimalarial treatment is needed together with splenectomy.

In summary, malaria should always be considered in any ill patient with a history of foreign travel through or within a malarial area. Following a clinical diagnosis, possibly supported by blood films, treatment should be prompt. Oral and parenteral quinine should be available in every district general hospital and given if the clinical picture is compatible with falciparum malaria. Because of the complexity of management and the possibility of concomitant foreign acquired infection, management ideally should be within an infectious diseases unit with facilities for intensive care. If this is not possible, minimum specialist input should be telephoned advice from the nearest infectious diseases or tropical medicine consultant.

Advice on malaria can be obtained by telephoning the following hospitals and asking for the infectious diseases consultant:

ABERDEEN	City Hospital	(0224) 681818
AIRDRIE	Monklands General Hospital	(023 64) 69344
BIRMINGHAM	East Birmingham District General Hospital	(021) 772 4311
BRADFORD	Leeds Road Hospital	(0274) 29661
BRISTOL	Ham Green Hospital	(0275) 81 2661
CAMBRIDGE	Addenbrookes	(0223) 45151
COVENTRY	Walsgrave General Hospital	(0203) 613232
DUNDEE	Kings Cross Hospital	(0382) 82541
EDINBURGH	City Hospital	(031) 447 1001
GLASGOW	Ruchill Hospital	(041) 946 7120
HUMBERSIDE (NORTH)	Castle Hill, Cottingham	(0482) 659331
IRVINE	Ayrshire Central	(0294) 2491
LEEDS	Seacroft	(0532) 648164
LEICESTER	Grobby Road	(0533) 874141
LIVERPOOL	Fazackerley Hospital	(051) 525 5980
	Liverpool School of Tropical Medicine	(051) 708 9393
LONDON	Coppetts Wood Hospital	(01) 883 9792
	Hither Green Hospital	(01) 698 4611
	Northwick Park Hospital	(01) 864 5311
	St George's Hospital	(01) 672 1255
	London School of Tropical Medicine	(01) 580 2127
MANCHESTER	Monsall Hospital	(061) 205 2393
MIDDLESBROUGH	Middlesbrough General	(0642) 83133
NEWCASTLE ON TYNE	Newcastle General	(091) 27 38811

NOTTINGHAM	City Hospital	(0602) 608111
OXFORD	Radcliffe	(0865) 49891
SHEFFIELD	Lodge Moor	(0742) 306555
STOKE ON TRENT	Bucknall Hospital	(0782) 25191
WINDYGATES, FIFE	Cameron Hospital	(0592) 2472

FURTHER REFERENCES

- Phillips-Howard P. A., Bradley D. J., Blaise M. & Hearn M. (1988) Malaria in Britain 1977–1986. *British Medical Journal* **296**, 245–248.
- Strickland G. T. (ed.) (1986) *Clinics in Tropical Medicine and Communicable Diseases: Malaria*, Vol. I, No. 1. W. B. Saunders & Company, London.
- WHO Malarial Action Programme (1986) Severe and complicated malaria. *Transcripts of the Royal Society of Tropical Medicine and Hygiene* **80**, S 1–50.

M. E. ELLIS
Regional Department of Infectious Diseases and Tropical Medicine,
Monsall Hospital,
University of Manchester School of Medicine,
Newton Heath,
Manchester, England