

development of the journal club. Anecdotally we found that participants did not feel that their reading time had significantly increased. However, their reading time was more focused, critical, and across a much wider range of journals. Overall the members of the group consider the journal club to be a success, and an effective use of the limited amount of time available for private study in a busy emergency department.

We have not assessed the impact of the journal club on clinical practice and patient care within the department as no data was collected prior to the development of the journal club system. Anecdotally lessons learned from BETs and structured journal scans have been implemented within the department.

Emergency medicine has close links with a large number of other specialties. It is important that evidence used in the emergency department is shared with in hospital specialties and on call teams. We believe that the construction and dissemination of the BETs can aid this interspecialty liaison and improve clinical practice.

Conclusion

Traditional methods of instituting evidence based medicine are not applicable to the emergency department setting. A structured journal club was designed to teach critical appraisal, keep abreast of developments in the field of emergency medicine, and institute evidence based medicine in this setting.

A proactive approach to evidence based medicine is necessary in emergency medicine. The use of a journal club to critically appraise journals relevant to emergency medicine and to construct and disseminate BETs, is one mechanism by which this can be achieved.

- 1 Sackett DL, Haynes RB, Guyatt GH, et al. *Clinical epidemiology—a basic science for clinical medicine*. London: Little Brown, 1991: 187.

- 2 Davidoff F, Haynes B, Sackett D, et al. Evidence-based medicine. A new journal to help doctors identify the information they need. *BMJ* 1995;310:1085–6.
- 3 Greenhalgh T. “Is my practice evidence based?” *BMJ* 1996; 313:957–8.
- 4 Board of the Faculty of Accident and Emergency Medicine. *Guidelines for specialist training in accident and emergency medicine*. London: Faculty of Accident and Emergency Medicine, 1996.
- 5 Crombie IK. *The pocket guide to critical appraisal*. London: BMJ Publishing Group, 1996.
- 6 Greenhalgh T. *How to read a paper: the basics of evidence based medicine*. London: BMJ Publishing Group, 1997.
- 7 Sackett DL, Richardson WS, Rosenberg W, et al. *Evidence based medicine—how to practice and teach EMB*. London: Churchill-Livingstone, 1996.
- 8 Carley S, Mackway-Jones K. Are British hospitals ready for the next major incident? Analysis of hospital major incident plans. *BMJ* 1996;313:1242–3.
- 9 Johnstone AJ, Zuberi SH, Scobie WG. Skull fractures in children: a population study. *J Accid Emerg Med* 1996;13: 386–9.
- 10 Feero S, Hedges JR, Simmons E. Does out-of-hospital EMS time affect trauma survival? *Am J Emerg Med* 1995;13: 133–5.
- 11 Schrader H, Obelieniene D, Bovim G, et al. Natural evolution of the late whiplash syndrome outside the medicolegal context. *Lancet* 1996;347:1207–11.
- 12 Abbaszadegan H, Jonsson U. Regional anaesthesia preferable for Colles’ fracture. Controlled comparison with local anaesthesia. *Acta Orthop Scand* 1990;61:348–9.
- 13 Browne GJ, Penna AS, Phung X, et al. Randomised trial of intravenous salbutamol in early management of acute severe asthma in children. *Lancet* 1997;349:301–5.
- 14 Du X, Cruikshank K, McNamee R, et al. Case-control study of stroke and the quality of hypertension control in north west England. *BMJ* 1997;314:272–6.
- 15 Stueland DT, Lee BC, Nordstrom DL, et al. A population based case-control study of agricultural injuries in children. *Inj Prev* 1996;2:192–6.
- 16 Johnson G. Hyperextension soft tissue injuries of the cervical spine—a review. *J Accid Emerg Med* 1996;13:3–8.
- 17 Hammacher ER, van der Werken Chr. Acute neck sprain: “whiplash” reappraised. *Injury* 1996;27:463–6.
- 18 Stevens A, Raftery J, eds. US task force on preventative health care. *Health care needs assessment: the epidemiologically based needs assessment reviews*. Second series. Oxford: Radcliffe Medical Press, 1997.
- 19 Mackway-Jones K, Carley SD, Morton RJ, et al. The best evidence topic report: a modified CAT for summarising the available evidence in emergency medicine. *J Accid Emerg Med* 1998;15:222–6.
- 20 Burstein JL, Hollander JE, Barlas D. Enhancing the value of journal club: use of a structured review instrument. *Am J Emerg Med* 1996;14:561–3.
- 21 Krough CL. A checklist system for critical review of medical literature. *Med Educ* 1985;19:392–5.
- 22 Markett RJ. A research methods and statistics journal club for residents. *Acad Med* 1989;April:223–4.

The best evidence topic report: a modified CAT for summarising the available evidence in emergency medicine

K Mackway-Jones, S D Carley, R J Morton, S Donnan

In order to achieve the optimal care for patients in the emergency department it is essential that their management is based upon the best available evidence. While the concept of evidence based medicine is widely accepted across many hospital and community specialties, emergency medicine has lagged behind in formalising the approach to evidence based practice.

One commonly used approach is that of the critically appraised topic (CAT).^{1,2} This was designed to be a one page summary of the evidence related to a particular clinical question. This established method would appear to be a

reasonable starting point for developing a clinically based approach to examining the evidence in emergency medicine. Unfortunately problems have been identified. CATs work best when based on papers that stand up to rigorous critical appraisal. In emergency medicine the evidence that does exist is frequently not of high quality, and consequently the critical appraisal process would tend to discard many papers because of either methodological flaws or poor design. The experience of the critical appraisal journal club³ was that an absolute requirement for high quality evidence meant

Department of
Emergency Medicine,
Manchester Royal
Infirmary, Oxford
Road, Manchester
M13 9WL

K Mackway-Jones
S D Carley
R J Morton

Department of Public
Health and
Epidemiology,
University of
Manchester,
Manchester
S Donnan

Correspondence to:
Kevin Mackway-Jones
(e-mail: kmackway@
fs1.mci.man.ac.uk).

Accepted for publication
16 April 1998

that large numbers of clinical questions in emergency medicine could not be answered at all. It became apparent during the searches that evidence of lower quality was available, and that this could be assimilated to give the best available answer at the present time.

This realisation that it was necessary to use the best available evidence led to the development of the best evidence topic report (BET) discussed below.

Design

BETs are constructed in four stages, based on the principles underlying all evidence based medicine. These stages are summarised in box 1 and are discussed below.

Box 1

- Asking the right question
- Searching for the evidence
- Appraising the evidence
- Summarising the evidence

ASKING THE RIGHT QUESTIONS

If a BET is to have real meaning to practising clinicians it is important that each topic selected is rooted in the practice of emergency medicine. In order to ensure that this is the case a clinical scenario is developed to illustrate the topic of interest. This scenario will be familiar to the person preparing the report, and will represent a real clinical problem in emergency medicine.

In order to ensure that the question is well defined and answerable a three part question is prepared from the clinical scenario. This process is well described as a tool for evidence based medicine.²⁻⁴ The three part question has the structure shown in box 2.

Box 2

- Patient characteristic
- Intervention(s) or defining question
- Relevant outcome(s)

It is vitally important to define the three part question well as this helps ensure that an appropriate search strategy can be formulated. An example of a general question of interest and a derived three part question are shown in box 3.

Box 3

General question

Is a haematoma block the best way to reduce a Colles' fracture?

Derived three part question

In [elderly patients with a Colles' fracture]
Is [a haematoma block better than a Bier's block]

At [reducing pain during manipulation and decreasing the need for remanipulation]

Table 1 Maximally sensitive randomised controlled trial filter

1	Randomised controlled trial.pt.
2	Controlled clinical trial.pt.
3	Randomised controlled trials.sh.
4	Random allocation.sh.
5	Double-blind method.sh.
6	Single-blind method.sh.
7	or/1-6
8	Animal.sh not human.sh.
9	7 not 8
10	Clinical trial.pt.
11	Exp clinical trials
12	(clin\$ adj25 trial\$.ti,ab.
13	((single or double or treble or triple) adj25 (blind\$ or mas\$)).ti,ab.
14	Placebos.sh.
15	Placebo\$.ti,ab.
16	Random\$.ti,ab.
17	Research design.sh.
18	or/10-17
19	18 not 8
20	19 not 9
21	Comparative study.sh.
22	Exp evaluation studies/
23	Follow up studies.sh.
24	Prospective studies.sh.
25	(control\$ or prospectiv\$ or volunteer\$).ti,ab.
26	or/21-25
27	26 not 8
28	26 not (9 or 20)
29	9 or 20 or 28

SEARCHING FOR EVIDENCE

In order to ensure that as much evidence as possible is included in the BET, it is essential that search strategies are sound and explicit. Two sources are explored in the construction of a BET. First the Medline database is searched using a strategy constructed as outlined below. Secondly the Best Evidence CD-ROM⁵ can be searched to see if there are any relevant systematic reviews.

Medline

There are three parts to the Medline search. First it is necessary to use search terms to produce a subset of the Medline database that contains all the papers relevant to the subject defined by the three part question. Second evidence based or methodological filters are applied if appropriate. Finally the titles and abstracts of the "hits" are scanned to see which papers are actually relevant to the question posed. Techniques for searching the Medline database have been reviewed recently.⁶

The subject search

In order to achieve the aims of the BET (finding and summarising the best evidence) it is important that the subject search has a high sensitivity (that is it has a high likelihood of retrieving all relevant items) so that important evidence is not overlooked. It may be necessary to use a number of different search strategies (that is use different search terms and combinations of terms) as well as using search terms that cover a slightly broader question. Greater sensitivity is achieved by using a combination of free text and text words together with medical subject headings (MeSH). A particularly useful feature enabling increased search sensitivity is the explosion command; this maps an individual MeSH term on to other related MeSH terms. Since some 50% of the articles




Figure 1 Overall search strategy for BETs (exp = explode; MeSH = medical subject headings; RCT = randomised controlled trial).

on Medline are misclassified by subject heading, this ability to search for related terms is invaluable.

If the highly sensitive search yields too many "hits" to be manageable, then it may be necessary to increase the specificity (the proportion of documents which are relevant) by refining the search. In general techniques for increasing specificity are the opposite of those for increasing sensitivity. Thus the question needs to be better defined and more specific terms need to be used in free text. Other techniques for increasing specificity include combining terms using the boolean operator *and* to combine different aspects of the question, and use of the Medline LIMIT command to set various general limits of the search such as language, species, or publication type.

As has been mentioned above the proper construction of a three part question holds the key to the search. A properly constructed question will allow a sensitive search to be carried out in a particular area of interest, so that all relevant papers relating to that area can be gathered and appraised.

Table 2 Type and strength of evidence

I	Strong evidence from at least one published systematic review of multiple well designed randomised controlled trials
II	Strong evidence from at least one published properly designed randomised controlled trial of appropriate size and in an appropriate clinical setting
III	Evidence from published well designed trials without randomisation, single group pre-post, cohort, time series, or matched case-control studies
IV	Evidence from well designed non-experimental studies from more than one centre or research group
V	Opinions of respected authorities, based on clinical evidence, descriptive studies, or reports of expert consensus committees

Applying filters

If a properly sensitive search carried out in a relevant area still yields a large number of hits then filters can be used. These are search strategies that have been designed to perform particular tasks such as finding randomised controlled trials or research overviews. A number of these are available in the public domain and have been published⁶ or can be found at appropriate web sites.⁷

The vast majority of questions of interest to emergency physicians at the present time (and therefore those most likely to appear as BETs) involve either treatments (interventions) or diagnosis. An appropriate filter for intervention studies in the context of BETs is a maximally sensitive randomised controlled trial filter; one such is shown in table 1.⁶

Diagnostic questions require a different approach since randomised controlled trials are not in general an appropriate method of investigation. In such cases an appropriate diagnostic evidence based medicine filter can be used. Diagnostic evidence filters and other evidence based filters appropriate to different types of questions (such as epidemiological or aetiological ones) are available from the Centre for Evidence Based Medicine and other appropriate sources.⁷

Making the searches explicit meant that they can easily be repeated to update the BET at a later date.

Scanning titles and abstracts

Once the subject search has been completed and any necessary filters have been applied a number of papers will fulfil the search criteria. There is no guarantee that these will all be relevant to the question posed and a final scan of titles and, if there is any doubt, abstracts will be necessary. Ideally a maximum of 50 papers will be left for scanning in this way, but on some occasions more than this number will remain despite increasing the specificity of the search. It is better to have a higher number of papers to scan than to miss relevant papers by over refining the search.

At the end of the scanning process a number of papers will be discarded as "irrelevant to the question". All the other papers remaining will need to be critically appraised. This process is described below.

The overall search strategy used in the construction of a BET is summarised in fig 1.

APPRAISING THE EVIDENCE

Evidence based medicine is not about dismissing all evidence that has not been gained by randomised trials and meta-analyses.⁸ Some types of question are not best answered by these approaches and appropriate evidence needs to be found for these. Furthermore for many questions in emergency medicine there is no appropriate research that stands up to rigorous critical appraisal. BETs are designed to find the best evidence that exists to answer a particular question. If high level evidence does exist then the question posed can be answered with some certainty. If high level evidence does not exist then best evidence available can be




Figure 2 Critical appraisal strategy.

summarised, and the failure to find good evidence can be used as a springboard for appropriate research.

The first critical appraisal filter to be applied to papers that are found by the search is methodological. The question underlying the methodological filter is: “what is the appropriate design of research to answer this question?”, and, if more than one approach is

possible the second question is: “what is the relative merit of each of the possible methods?” In the case of questions concerning interventions the hierarchy of evidence shown in table 2 is generally accepted⁷ and is used in constructing BETs. Although other hierarchies have been proposed⁹ the concepts underlying them all are the same.

The highest available level of evidence is used to construct the BET—thus level I evidence will be used if it exists; if there is no level I evidence then level II will be sought and so on.

Once papers containing the highest level of evidence available have been identified, they are critically appraised. The critical appraisal methods used are standard and have been published in a number of places.^{2 10 11} Key questions include methodological and analytical soundness.

After critical appraisal a paper may be found to be so flawed as to be unusable and will thus be discarded as of “insufficient quality for inclusion”. If a paper is not fatally flawed or if, despite being flawed, it represents the best evidence that is available it will be included in the BET. Comments about study weaknesses will be included in the last column of the table summarising the evidence.

The overall strategy for critical appraisal used in the construction of the BET is summarised in fig 2.

SUMMARISING THE EVIDENCE

In order to keep a record of appraised topics a summary of the evidence is made in the form of a BET. The first of these BETs appear in this month’s journal. The reports follow a standardised format.¹² This includes the title, the clinical scenario, the derived three part question, the detailed search strategy, and the results of the search. The number of “hits” is followed by the number of papers which were not relevant to the question and the number which were of insufficient quality for inclusion (after critical appraisal). Those papers which are included in the BET are reported in table format. The table includes the author, date, and country of the research, the patients groups the study type (and the level of evidence), the outcomes that were investigated,

Table 3 Template for a BET

Title					
Report by:					
Search checked by:					
Clinical scenario					
Three part question					
Search strategy					
Search outcome					
Author, date, and country	Patient group	Study type (level of evidence)	Outcomes	Key results	Study weaknesses
Comment					
Clinical bottom line					
References					

the key results for each outcome and the study weaknesses that were identified. A further analysis of the results of the best papers may be given if the results allow this. Free text comment about the search and its outcome is then given, followed by the clinical bottom line (the answer to the original question). The references of the relevant papers are given in full.

The template for the BET is shown in table 3.

Each BET is attributed to a main author and also a subsidiary author who is responsible for checking the search strategy and outcome.

NEGATIVE BET REPORTS

It is foreseen that a significant number of topics will fail to produce any evidence to answer the clinical question posed. These "negative" BETs are indicative of areas for further research, and a list of the questions for which no answer at all can be found should be circulated alongside "positive" reports.

Discussion

BETs are an attempt to formalise evidence based practice in the emergency department. Their very name indicates the uncertain nature of much of the "evidence" in this area. The hope is that by basing BETs on real problems in emergency medicine, real questions will be answered with the best evidence available. This process is essential as real questions about patient care arise on a daily basis in the emergency department, yet there is rarely time to search for evidence at the time of a patient presentation. It is therefore essential to be proactive in the approach to evidence based practice. Emergency medicine is still a relatively young specialty and there is a paucity of high quality evidence for many aspects of practice, BETs will therefore not be confined to questions regarding new or novel practices and will examine the evidence for well established practices as well.

The BET does have its weaknesses. While it reports the best available evidence obtained

through literature searching on Medline and other information resources, it does not examine unpublished literature in the way that systematic reviews do. It is unlikely that systematic reviews will become available for many of the more minor complaints that present to the emergency department for some time. Even if they did, as the BETs will show, it is unlikely that much of the evidence would pass the methodological filters used in systematic reviews.

BETs contain the best evidence that can be practically obtained by busy practising clinicians and should be used to inform best practice. They can be disseminated around members of journal clubs, training schemes, and can be published in peer reviewed journals or web sites. It is hoped that BETs will demonstrate both the strengths and the weaknesses of the evidence base on which the practice of emergency medicine is based.

- 1 Sauve S, Lee HN, Meade MD, et al. The critically appraised topic: a practical approach to learning critical appraisal. *Ann R Coll Phys Surg Canada* 1995;28:396-8.
- 2 Sackett DL, Richardson WS, Rosenberg W, et al. *Evidence based medicine: how to practice and teach EBM*. New York: Churchill Livingstone, 1997.
- 3 Carley SD, Mackway-Jones K, Jones A, et al. Moving towards evidence based emergency medicine: use of a structured critical appraisal journal club. *J Accid Emerg Med* 1998;15:220-2.
- 4 Richardson WS, Wilson MC, Nishikawa J, et al. The well built clinical question: a key to evidence-based decisions (editorial). *ACP J Club* 1995;123:A12-13.
- 5 Best evidence CD-ROM. London: BMJ Publishing Group, 1997.
- 6 Greenhalgh T. The Medline database. *BMJ* 1997;315:180-3.
- 7 Centre for Evidence Based Medicine <http://cebmr2.ox.ac.uk>
- 8 Sackett DL, Rosenberg WMC, Gray JAM, et al. Evidence based medicine: what it is and what it isn't. *BMJ* 1996;312:71-2.
- 9 Stevens A, Raftery J, eds. US task force on preventative health care. *Health care needs assessment: the epidemiologically based needs assessment reviews*. Second series. Abingdon: Radcliffe Medical Press, 1997.
- 10 Crombie IK. *The pocket guide to critical appraisal*. London: BMJ Publishing Group, 1996.
- 11 Greenhalgh T. *How to read a paper: the basics of evidence based medicine*. London: BMJ Publishing Group, 1997.
- 12 Mackway-Jones K, ed. Towards evidence based emergency medicine: best BETs from the Manchester Royal Infirmary. *J Accid Emerg Med* 1998;15:227-30.