

CASE REPORTS

A case of extreme hypercapnia: implications for the prehospital and accident and emergency department management of acutely dyspnoeic patients

L Urwin, R Murphy, C Robertson, A Pollok

Emerg Med J 2004;21:119-120

A 64 year old woman was brought by ambulance to the accident and emergency department. She had been referred by her GP because of increasing dyspnoea, cyanosis, and lethargy over the previous four days. On arrival of the ambulance crew at her home she was noted to be tachycardic and tachypnoeic (respiratory rate 36/min) with a GCS of 5 (E 3, M 1, V 1). She was given oxygen at 6 l/min via a Duo mask, and transferred to hospital.

The patient arrived at the accident and emergency department 18 minutes later. In transit, there had been a clinical deterioration. The GCS was now 3 and the respiratory rate 4/min. Oxygen saturation, as measured by a pulse oximeter was 99%. The patient was intubated and positive pressure ventilation started. Arterial blood gas measurements taken at the time of intubation were consistent with acute on chronic respiratory failure (fig 1).

Further investigations included a chest radiograph, which showed hyperinflation of the lungs and a 12-lead ECG, which showed right axis deviation and right atrial hypertrophy.

Treatment with nebulised bronchodilators, intravenous hydrocortisone, and antibiotics was started empirically with a diagnosis of carbon dioxide narcosis attributable to oxygen therapy in a patient with previously undiagnosed chronic obstructive pulmonary disease. Repeat arterial blood gas analysis was performed seven minutes after intubation and showed: P_{aO_2} 55.2 kPa, P_{aCO_2} 21.74 kPa, H^+ 101.0 nmol/L, HCO_3^- 40.3 mmol/L.

The patient was transferred to the intensive care unit where mechanical ventilation was continued. Further history revealed that she had smoked 20 cigarettes/day for 40 years but had previously been able to walk as far as her local shops without becoming short of breath. She had not seen her GP nor attended hospital for the previous 16 years.

The following day the patient was much improved. Over a period of two days she was weaned from the ventilator and extubated. She was discharged from hospital to home nine days after admission. At discharge, on room air, arterial blood gas pressures were as follows: P_{aO_2} 7.52 kPa, P_{aCO_2} 7.61 kPa, H^+ 37.0 nmol/L, HCO_3^- 36.7 mmol/L.

DISCUSSION

Prehospital, the emergency treatment of patients with dyspnoea must often be undertaken without the benefit of a definitive diagnosis. In the United States between 9% and 15% of patients with acute dyspnoea have chronic obstructive pulmonary disease.¹ In many situations it is possible to identify these patients and modify treatment accordingly. This necessitates some access to the medical history however and on occasion, as in this case, this is not available.

Unfortunately when such a history is unavailable, no out of hospital monitoring technique exists to identify patients with, or at risk of, carbon dioxide retention. Pulse oximetry is used by many ambulance services and can be an extremely

useful non-invasive technique to aid the assessment of peripheral oxygen saturation. In situations of poor perfusion, movement and abnormal haemoglobin, however, this technique may not reliably reflect P_{aO_2} values. More importantly, and as shown in our case, there is no definite relation between S_{aO_2} values measured by pulse oximetry and P_{aCO_2} values although it has been shown that the more oxygenated

Acc. No	0001	B.P.	745 mmHg
Arterial sample			
Patient info			
pH/blood gases			
Measured at	37°C		
cH	134.0 nmol/L		
pCO ₂	31.05 KPa		
pO ₂	18.0 KPa		
Electolytes			
Na ⁺	131 mmol/L		
K ⁺	4.9 mmol/L		
Ca ⁺⁺	1.20 mmol/L		
Calculations			
HCO ₃ ⁻	43.4 mmol/L		
TCO ₂	50.5 mmol/L		
BE _b	3.5 mmol/L		
BE _{Ecf}	9.8 mmol/L		
SBC	27.5 mmol/L		
sO _{2c}	94.2%		

Figure 1 Arterial blood gas measurements.

patients with chronic obstructive pulmonary disease become the higher the incidence of respiratory acidosis.²

Clearly, in patients known to have chronic obstructive pulmonary disease it is important to control oxygen therapy to ensure the correction of dangerous levels of hypoxia while minimising the risk of hypercapnia. Despite the fact that our patient deteriorated with the administration of high concentration oxygen, this case should not be used to imply that all dyspnoeic patients should be given lower concentrations of oxygen. In many situations such as acute pulmonary oedema, anaphylaxis, acute asthma, etc, high concentrations of oxygen are essential and even in some patients with acute exacerbations of chronic obstructive pulmonary disease low concentration oxygen will not provide adequate relief of hypoxia.³

This case highlights a number of useful learning points:

- Firstly, not all patients with severe chronic obstructive pulmonary disease are identified in the community. Consequently when they become increasingly dyspnoeic there is a risk that inappropriately high concentrations of oxygen may be administered.
- Secondly, the well known risks of inducing hypercapnia and carbon dioxide narcosis cannot reliably be judged by currently available out of hospital monitoring techniques.

- Thirdly, extreme levels of hypercapnia, provided they are treated promptly and appropriately, are compatible with full recovery. To our knowledge, the PaCO_2 value of 31.05 kPa with which this patient presented, is the highest recorded in the human literature.

Authors' affiliations

I Urwin, R Murphy, C Robertson, Department of Accident and Emergency Medicine, Edinburgh Royal Infirmary, Edinburgh, UK
A Pollok, Department of Intensive Care Medicine, Edinburgh Royal Infirmary

Correspondence to: Mr R Murphy, Department of Accident and Emergency Medicine, The Royal Infirmary of Edinburgh, Old Dalkeith Road, Little France, Edinburgh EH16 4SU, UK; rossmurphy@doctors.org.uk

Accepted for publication 26 March 2003

REFERENCES

- 1 **Michelson E**, Holbrah S. Evaluation of the patient with shortness of breath: an evidence based approach. *Emerg Med Clin North Am* 1999;17:221–37.
- 2 **Plant PK**, Owen JL, Elliott MW. One year period prevalence study of respiratory acidosis in acute exacerbations of COPD: implications for the provision of non-invasive ventilation and oxygen administration. *Thorax* 2000;55:550–4.
- 3 **Hutchison DCS**, Flenley DC, Donald KW. Controlled oxygen therapy in respiratory failure. *BMJ* 1964;2:1159–66.

Hypokalaemic thyrotoxic periodic paralysis in an Asian man in the United Kingdom

R Sinharay

Emerg Med J 2004;21:120–121

A large number of ethnic Chinese and other oriental populations are living in the West because of the modern day migration of people. Hypokalaemic periodic paralysis attributable to thyrotoxicosis is a common presentation in an Asian emergency department. It is uncommon in the white communities. There is a difference in the genetic type in the different racial groups. Thyrotoxic features are often masked or absent. Life may be threatened because of severe hypokalaemia and therefore a prompt diagnosis of this condition in the certain ethnic group presenting with weakness and hypokalaemia is essential. Thyroid function studies are mandatory in these cases.

In an Asian population, thyrotoxic periodic paralysis (TPP) is most commonly associated with hypokalaemia.^{1–4} As oriental races are now frequently found in the West, TPP should be included in the differential diagnosis of muscle weakness with hypokalaemia in an emergency department.

CASE REPORT

In October 2002, a 30 year old Vietnamese of Chinese parentage presented in the early hours of one morning complaining of sudden generalised weakness. The previous day he had worked hard in his “take away” restaurant. He

then had a carbohydrate rich meal at around 8 pm and went to bed at around 12 midnight. He had no significant medical or family history.

He had a tachycardia of 100/min and his blood pressure was 119/85. He was fully alert and coherent. His motor power was reduced. He was totally areflexic with both plantars downgoing. He was clinically euthyroid.

His initial serum K^+ was 1.2 mmol/l (3.5–4.8) with normal renal function, calcium, glucose, and liver function tests. His serum creatinine phosphokinase concentration was 1099 IU/l (38–148). His blood pH was 7.44. His Hb%, white blood cell count, platelets, erythrocyte sedimentation rate, and C reactive proteins were all normal.

Treatment was started with intravenous administration of K^+ of 80 mmol over 90 minutes. Once the serum K^+ reached 4.2 his weakness disappeared. His subsequent serum K^+ remained normal. His urine electrolytes in 24 hours' urine were within normal range. His free thyroxine (FT4) level was 73 pmol/l (7–17), and the TSH level was 0.02 mu/l (0.35–2.75). Anti-thyroid peroxidase antibodies were 439 IU/ml (0.0–35) on admission, which later was increased to 723 IU/ml. Electrocardiogram showed mild tachycardia of 100/min and U-waves in lead II, III, VI–V6.

Ultrasound examination of the thyroid gland showed mild enlargement of the thyroid gland with diffuse parenchymal disease. He was found to have the following human leucocyte antigen (HLA) sub-type: A11, A33; B15, B58; DR3; DQ2, DQ1.

Table 1 Diagnostic criteria in two most common forms of hypokalaemic periodic paralysis

	Thyrotoxic hypokalaemic periodic paralysis	Hypokalaemic periodic paralysis (non-familial)
Episodic	Yes	Yes
Acute episodes	Yes	Yes
Hypokalaemia	Yes	Yes
Family history of paralysis	No	No
Large carbohydrate meal	Often	Rare
Adrenergic stress	Often	Rare
Rapid pulse rate	Yes	No
Blood pressure	Sometimes high	Normal
Features of thyrotoxicosis	May be absent	No
Family history of hyperthyroidism	Some	No

He was given a carbimazole and β blocker combination. In three weeks his FT4 came down to 35 pmol/l. He became biochemically euthyroid by the end of the 14th week. He has remained hypokalaemic weakness free ever since.

DISCUSSION

This patient has Graves' disease with hypokalaemic paralysis. He recovered quickly with K^+ replacement. He was given a much larger dose of intravenous K^+ , instead of the recommended 10 mmol/h,⁵ but fortunately did not show any rebound rise.⁴

Many reported cases of TPP did not have clinical features of thyrotoxicosis.⁵ However, in the Asian patients hyperthyroidism were always obvious, although this has been disputed recently in another study.⁹ A total of 2% of thyrotoxic patients in China and Japan were reported to have TPP,² but it has also been recognised in the Thai, Vietnamese, Korean, and Malay populations.⁹

TPP is characterised by sudden transient recurrent episodes of painless paralysis usually occurring after heavy exertion or a high carbohydrate meal, followed by a prolonged rest.⁵ The diurnal variation in K^+ movement where there is nocturnal K^+ influx into the muscles would explain the tendency for TPP to occur at night.⁷ Deep tendon reflexes are depressed in most patients.⁵ Respiratory muscles are usually, but not always spared.³

Recovery can be spontaneous after 3 to 36 hours, but must be speeded up by K^+ administration. The onset of paralytic attacks in familial periodic paralysis is often during adolescence, but in most published reports, TPP were found to be between the ages of 20 and 40 years.⁴ The male:female ratio in TPP has been estimated to be 20:1.² Recently, a case of TPP with hypokalaemia has been described in a white woman.⁸ Table 1 shows the differential diagnostic criteria for the two most common forms of hypokalaemic periodic paralysis.

Ober reported seven cases of TPP and provided an excellent review of the literature.¹ It has been postulated that the thyroid hormone directly stimulates Na-K-ATPase dependent K^+ channel and increases the catecholamine mediated intracellular potassium shift, which is also stimulated by insulin.⁵ Therefore, β blockers are a very useful treatment in TPP, associated with hypokalaemia.⁵

Male dominance and the racial difference in TPP with hypokalaemia are very puzzling.⁸ Difference in HLA subtypes

in different racial groups have been reported in the past. Japanese (DRw 8), Singapore Chinese (A2BW22, AW19B17), and Hong Kong Chinese (B5, BW46), all have different HLA subtypes.⁵ HLA DRw8 makes the Japanese susceptible to TPP, and yet the same HLA subtype in the white population makes them susceptible to Graves' disease.¹⁰ A genetic mutation in the control of Na-K-ATPase activity within the same HLA subtype may explain the racial difference. Interestingly, a genetic mutation in TPP has recently been described in the K^+ -ionic channel gene KCNE3.¹¹ Anti-TPO autoantibody is present in 85%–95% of Graves' disease cases,¹² which is diagnostic of the condition.

This case shows that among the Chinese and other Oriental population living in Britain, hypokalaemic periodic paralysis may be caused by thyrotoxicosis and that thyroid function studies are mandatory in these people.

Correspondence to: Dr R Sinharay, Department of Medicine, Tameside General Hospital, Fountain Street, Ashton Under Lyne, Lancashire OL6 9RW, UK; ranjitsinharay@hotmail.com

Accepted for publication 9 April 2003

REFERENCES

- Ober KP. Thyrotoxic periodic paralysis in the United States: report of 7 cases and review of the literature. *Medicine* 1992;**71**:109–20.
- Mcfadzean AJS, Yeung R. Periodic paralysis complicating thyrotoxicosis in Chinese. *BMJ* 1967;**1**:451–5.
- Ahlawat SK, Sachdev A. Hypokalaemic paralysis. *Postgrad Med J* 1999;**75**:193–7.
- Manoukain MA, Foote JA, Crapo LM. Clinical and metabolic features of thyrotoxic periodic paralysis in 24 episodes. *Arch Intern Med* 1999;**159**:601–6.
- Salifu MO, Ota K, Carroll HJ, et al. Thyrotoxic hypokalaemic paralysis in a Blackman. *Q J Med* 2001;**94**:659–60.
- Engel AG. Thyroid function and periodic paralysis. *Am J Med* 1961;**30**:327–33.
- Charness ME, Johns RJ. Hypokalaemic periodic paralysis. *Johns Hopkins Med J* 1978;**143**:148–53.
- Dixon AN, Jones R. Thyrotoxic paralysis in a white woman. *Postgrad Med J* 2002;**78**:687–8.
- Goh SH. Thyrotoxic periodic paralysis: reports of seven patients presenting with weakness in an Asian emergency department. *Emerg Med J* 2002;**19**:78–9.
- Lazarus JH. Hyperthyroidism. *Lancet* 1997;**349**:339–43.
- Da Silva MRD, Cerutti JM, Arnaldi LAT, et al. A mutation in the KCNE3 potassium channel gene is associated with susceptibility to thyrotoxic hypokalaemic periodic paralysis. *J Clin Endocrinol Metab* 2002;**87**:4881–4.
- Kotani T. anti-TPO autoantibodies. *Rinsho Byori* 1998;**46**:324–30.

Fatal descending necrotising mediastinitis

M Bulut, V Balci, Ş Akköse, E Armağan

Emerg Med J 2004;21:122–123

Descending necrotising mediastinitis rarely develops and this variety of mediastinitis is a highly lethal disease. A case is reported of descending necrotising mediastinitis caused by an odontogenic infection. The importance is emphasised of prompt diagnosis and aggressive surgical mediastinal drainage for the survival of these patients. Most acute mediastinal infections result from oesophageal perforation, either secondary to oesophagoscopy or tumour erosion. Mediastinitis occasionally develops as descending necrotising mediastinitis originating from the complications of cervical or odontogenic infections. Descending necrotising mediastinitis usually has a fulminant course, leading commonly to sepsis and death.

A 16 year old woman presented to our accident and emergency (A&E) department complaining of dyspnea, fever, pleuritic chest pain, and swollen throat. Her initial complaints had been a right sided premolar toothache and swollen cheek. Initially she had been treated as an outpatient with parenteral antibiotics for five days in a local medical facility. Upon worsening of her condition (orthopnea, high fever, chest pain), the patient was brought by ambulance to our A&E department.

On admission, blood pressure was 110/70 mm Hg, pulse rate was 110 beat/min, respiration rate was 30 breath/min, and her body temperature was 38.9°C. She appeared to be sick and orthopneic. Physical examination revealed dyspnea, orthopnea, diffuse painless cervical swelling, and decreased breath sounds on the right side of the chest.

Laboratory tests were as follows: white blood cells 17.600/mm³, haemoglobin 9.6 g/dl, platelets 481000/mm³. Blood glucose, urea, sodium, potassium, creatinine, and liver function tests were in the normal range.

Chest roentgenogram showed a pneumomediastinum and pneumohydrothorax on the right side. Computed tomography (CT) of the neck and the chest revealed gas bubbles in the neck region, marked inflammation in the mediastinal soft tissues and azygo-oesophageal recessus suggesting a mediastinal abscess and a pleural effusion and pneumothorax on the right side were noted.

In the A&E department, 1300 ml empyema was drained from the right hemithorax via a chest tube. She was then taken to the operating room and her mediastinal abscess was drained. Intravenous triple antibiotic treatment (ceftazidime 2×1 g IV, amikacin 1×1 g IV, and metronidazole 3×500 mg IV) was started. The following day, repeat neck CT was done and revealed a submandibular abscess, which was also drained. Bacteriological results from samples obtained from the neck, pleura, mediastinum, and blood did not reveal any micro-organism. On hospital day five, another cervicothoracic CT was done and revealed residual pleural empyema and effusion around the major vessels, gas in the soft tissues of the neck and the mediastinum, and a pneumonic infiltration in the posterior lobe of the left lung.

Although she continued to receive intravenous antibiotics, she remained febrile for nine days. At this time her general condition and arterial blood gas pressures deteriorated. She was in significant respiratory distress and she, once again, needed endotracheal intubation. But this was not possible because of massive laryngeal oedema. A tracheostomy was attempted but meanwhile she developed cardiorespiratory arrest. CPR was not successful.

DISCUSSION

DNM is an uncommon complication of oral and pharyngeal infections,^{1–3} spreading caudally through the retropharyngeal space, which is limited by the middle and deep layers of the deep cervical fascia.⁴

The criteria for the diagnosis of DNM were accurately defined by Estrera and colleagues.⁵ These criteria include: 1—clinical manifestation of severe oropharyngeal infection (odontogenic, peritonsillar, or retropharyngeal abscesses, Ludwig's angina, or infection secondary to traumatic pharyngeal perforations); 2—demonstration of characteristic roentgenographic features of mediastinitis; 3—documentation of mediastinitis at operation or postmortem examination, or both; and 4—establishment of relation between oropharyngeal infection and development of the necrotising process in the mediastinum.^{1 3 5 6} Our patient met all these criteria.

Mediastinitis occurring secondary to descending odontogenic or deep cervical infections is polymicrobial and mixed, caused by a combination of aerobic and anaerobic organisms reflecting the oropharyngeal flora.^{4 6 7 8} Review of the literature shows that although DNM is quite rare, this variety of mediastinitis is a highly lethal disease. The mortality rate of this disease is between 40% and 50%.^{1 3 4} Marty-Ane and associates reported that aggressive surgery provides improved results with an 83% survival rate.^{1 3 4}

Delayed diagnosis and inadequate drainage are the main causes of the high mortality rate in DNM.^{5 7 9} Because of the absence of early clinical or radiological signs, diagnosis is usually delayed. The diagnosis of acute mediastinitis from conventional radiological studies may be difficult. CT is an extremely useful tool in the treatment of cervical infections and the diagnosis of mediastinitis.^{1 3 7 10} CT is also extremely useful in the postoperative assessment of the results of surgical drainage and the timing of the re-operations in patients with continued sepsis.^{1 3 5 7}

The mainstay of treatment in patients with DNM is aggressive and adequate surgical drainage of the cervical and mediastinal collections and intravenous broad spectrum antibiotic treatment.^{1 3 4 11}

Mediastinitis secondary to descending odontogenic and deep cervical infections is rare. It should be suspected in patients whose conditions are disproportionately severe to be justified by a trivial pharyngeal or odontogenic infection, or conversely, in whom a mediastinitis develops in the absence of known risk factors, such as previous surgical and diagnostic procedures involving the oesophagus or the upper airways or pharyngeal foreign bodies. At present, CT is a

valuable tool for early diagnosis of DNM in all patients with deep neck infections to determine the presence of mediastinal involvement.

Authors' affiliations

M Bulut, V Balci, Ş Akköse, E Armağan, Department of Emergency Medicine, Uludag University Medical School, Bursa, Turkey

Correspondence to: Dr M Bulut, Department of Emergency Medicine, Uludag University Medical School, 16049 Bursa, Turkey; mbulut94@yahoo.com

Accepted for publication 14 January 2003

REFERENCES

- Marty-Ane CH, Berthel JP, Alric P, et al. Management of descending necrotizing mediastinitis: An aggressive treatment for an aggressive disease. *Ann Thorac Surg* 1999;**68**:212–17.
- Brunelli A, Sabbatini A, Catalini G, et al. Descending necrotizing mediastinitis. Surgical drainage and tracheostomy. *Arch Otolaryngol Head Neck Surg* 1996;**122**:1326–9.
- Wheatley MJ, Stirling MC, Kirsh MM, et al. Descending necrotizing mediastinitis: transcervical drainage is not enough. *Ann Thorac Surg* 1990;**49**:780–4.
- Marty-Ane CH, Alauzen M, Alric P, et al. Descending necrotizing mediastinitis. Advantage of mediastinal drainage with thoracotomy. *J Thorac Cardiovasc Surg* 1994;**107**:55–61.
- Estrera AS, Landay MJ, Grisham JM, et al. Descending necrotizing mediastinitis. *Surg Gynecol Obstet* 1983;**157**:545–52.
- Berlot G, Tomasini A, Cioffi V, et al. Fatal streptococcus viridans descending mediastinitis: case report and review of the literature. *Eur J Emerg Med* 1997;**4**:111–14.
- Levine TM, Wurster CF, Kresp YP. Mediastinitis occurring as a complication of odontogenic infections. *Laryngoscope* 1986;**96**:747–50.
- Moncada R, Warpeha R, Pickleman J, et al. Mediastinitis from odontogenic and deep cervical infection. Anatomic pathways of propagation. *Chest* 1978;**73**:497–500.
- Morey-Mas M, Caubet-Biayna J, Iriarte-Ortobe JL. Mediastinitis as a rare complication of an odontogenic infection. Report of a case. *Acta Stomatol Belg* 1996;**93**:125–8.
- Kim HJ, Park ED, Kim JH, et al. Odontogenic versus nonodontogenic deep neck space infections: CT manifestations. *J Comput Assist Tomogr* 1997;**21**:202–8.
- Payne WS, Larson RH. Acute mediastinitis. *Surg Clin North Am* 1969;**49**:999–1009.

Troponin testing: beware pulmonary embolus

S Conroy, I Kamal, J Cooper

Emerg Med J 2004;**21**:123–124

Serum troponin estimation is widely used in the diagnosis and management of coronary syndromes, but it is possible to be misled by a positive result unless it is put carefully into clinical context. The serum troponin can be positive in pulmonary embolus and carries prognostic significance. A case report is presented and a review of the relevant literature.

Troponin assays have had a considerable impact in the diagnosis and risk stratification of patients presenting with an acute coronary syndrome. A correctly timed troponin assay has 95%–100% sensitivity and 85%–90% specificity for myocardial damage.^{1–4} However, there are many other causes of a positive troponin test that should be remembered when assessing these patients in order to avoid misdiagnosis.

CASE REPORT

A 77 year old woman was admitted to hospital with an exacerbation of chronic obstructive pulmonary disease. She had a background of controlled hypertension, paroxysmal atrial fibrillation, and had been a heavy smoker for 30 years. She responded well to initial treatment with antibiotics, bronchodilators, and corticosteroids.

Five days after her admission, she deteriorated acutely with tight central chest pain and dyspnoea. On examination, she was tachycardic (rate 110, sinus rhythm), hypotensive (BP 90/50), but heart sounds were normal and the jugular venous pressure was not elevated. Her ECG showed ST segment depression and T-wave inversion antero-laterally, suggestive of myocardial ischaemia. The chest radiograph showed hyperinflated lung fields but was otherwise unremarkable; her peripheral oxygen saturations were 85%. She was treated with diamorphine, low molecular weight

Joint European and American guidelines on the diagnosis of myocardial infarction⁸

- A typical rise and fall in troponin or CK-MB
- Plus at least one of the following:
 - ischaemic symptoms
 - development of pathological Q waves on the ECG
 - ECG changes suggestive of ischaemia—ST elevation or ST depression
- Coronary artery intervention, for example, coronary angioplasty

heparin, and aspirin. There was a small rise in creatine kinase (52 to 209 IU/l (normal range 24–195 IU/l)) and the troponin T 12 hours after the pain was significantly raised at 2.1 µg/l (normal <0.01 µg/l), consistent with significant myocardial damage. Two days later she was still limited by significant dyspnoea and hypoxia (pO₂ 8.3 on air) and the diagnosis of pulmonary embolus was suspected. An echocardiogram was normal in particular demonstrating normal left and right ventricular dimensions and no regional wall motion abnormalities. There was no significant tricuspid regurgitation to permit an estimation of the pulmonary artery pressure. A D-dimer assay was strongly positive at 3872 ng/ml (normal <500 ng/ml) and a ventilation-perfusion scan showed multiple unmatched defects, indicating a high probability of pulmonary embolus.

DISCUSSION

An increased serum troponin concentration is found not only in acute coronary syndromes but in several other conditions: myocarditis, stroke, myocardial trauma, post-CABG, chronic renal failure, ischaemic heart failure, ischaemic arrhythmia,

and pulmonary embolism.⁵⁻⁷ The combination of the ischaemic ECG and positive troponin test made pulmonary embolus a difficult initial diagnosis to make in this patient. Indeed, our patient met the revised joint guidelines⁸ for the diagnosis of myocardial infarction. Whether there was concomitant pulmonary embolus and myocardial infarction or pulmonary embolus revealing underlying myocardial ischaemia, is difficult to determine absolutely as further investigation (for example, coronary angiography) was not performed.

Typically, the ECG changes of a pulmonary embolus, if any, are those of either a sinus tachycardia or atrial fibrillation with right heart strain, though many electrocardiographic abnormalities have been reported—rhythm disturbances, axis change, ST-T changes and QT interval prolongation.⁹⁻¹¹ However, such changes are often transient and the value of the 12-lead ECG in the diagnosis and risk stratification of suspected pulmonary embolus is limited.¹² ST elevation on the ECG in pulmonary embolus is recognised but an unusual finding.¹³ Given the cardiovascular risk factors present in our patient, it seems probable that there was significant coronary artery disease, which was revealed by the hypoxia and acute cardiac insult of a pulmonary embolus.

A positive troponin test is an important prognostic marker in pulmonary embolus, and predicts right ventricular dilatation and an increased number of segmental perfusion defects on lung scintigraphy.¹⁴ In a series of 56 patients presenting with pulmonary embolism, 18 (32%) had an increased troponin T concentration (>0.1 µg/l) taken at 12 hours after presentation. These patients had higher risk of developing complications including in-hospital death (odds ratio 29.6:1, 95% CI 3.3 to 265.3), prolonged hypotension and cardiogenic shock (odds ratio 11.4:1, 95% CI 2.1 to 63.4), the need for inotropic support (odds ratio 37.6:1, 95% CI 5.8 to 245.6), and mechanical ventilation (odds ratio 78.8:1, 95% CI 9.5 to 653.2).¹⁵ Moreover, after adjustment, an increased troponin T value was an independent predictor of 30 day mortality in pulmonary embolus (odds ratio 15.2:1, 95% CI 1.22 to 190.4). It is suggested that troponin estimation may be used to help risk stratify patients with pulmonary embolus who warrant more aggressive treatment.

The recent Joint American and European guidelines⁸ on the diagnosis of myocardial infarction rely heavily upon troponin testing (box). An awareness of other conditions that cause an increased troponin concentration is needed to avoid misdiagnoses. This is particularly the case for patients with pulmonary emboli that may clinically mimic myocardial ischaemia but for which the management is completely different.

Contributors

Simon Conroy: original idea and literature review. Imad Kamal: advice on content and style. John Cooper: advice on cardiological aspects, style, and content.

Consent for publication was not sought in view of the anonymity of the people concerned.

Authors' affiliations

S Conroy, I Kamal, J Cooper, Bedford Hospital, Bedford, UK

Correspondence to: Dr S Conroy, Bedford Hospital, Kempston Road, Bedford MK42 9DJ, UK; spcon@doctors.org.uk

Accepted for publication 9 January 2003

REFERENCES

- Gerhardt W, Ljungdahl L. Detection of myocardial damage by serial measurements of cardiac troponin T, CK MBmass, and TROP T rapid test. *Cardiovasc Drugs Ther* 1997;**11**(suppl 1):227-40.
- Gerhardt W, Ljungdahl L. Troponin T. A sensitive and specific diagnostic and prognostic marker of myocardial damage. *Clin Chim Acta* 1998;**272**:47-57.
- Gerhardt W, Ljungdahl L, Herbert AK. Troponin-T and CK MB (mass) in early diagnosis of ischemic myocardial injury. The Helsingborg Study, 1992. *Clin Biochem* 1993;**26**:231-40.
- Stubbs P, Collinson P, Moseley D, et al. Prospective study of the role of cardiac troponin T in patients admitted with unstable angina. *BMJ* 1996;**313**:262-4.
- Douketis JD, Crowther MA, Ginsberg JS. Elevated cardiac troponin levels in patients with submassive pulmonary embolism. *Arch Intern Med* 2002;**162**:79-81.
- Pacouret G, Schellenberg F, Hamel E, et al. Troponin I in massive acute pulmonary embolism; results of a prospective series. *Presse Med* 1998;**27**:1627.
- Wong CB, Tang IT, Tiu A. Pulmonary embolism mimicking acute myocardial infarction. *Tex Med* 1999;**95**:10.
- Joint European Society of Cardiology/American College of Cardiology Committee. Myocardial infarction redefined—a consensus document of the Joint European Society of Cardiology/American College of Cardiology Committee for the redefinition of myocardial infarction. *Eur Heart J* 2000;**21**:1502-3.
- Sreeram N, Cheirix EC, Smeets JLRM, et al. Value of the 12-lead ECG at hospital admission in the diagnosis of pulmonary embolus. *Am J Cardiol*, 1994;**73**:298.
- Ullman E, Brady WJ, Perron AD, et al. Electrocardiographic manifestations of pulmonary embolism. *Am J Emerg Med* 2001;**19**:514-19.
- Watanabe T, Kikushima S, Tanno K, et al. Uncommon electrocardiographic changes corresponding to symptoms during recurrent pulmonary embolism as documented by computed tomography scans. *Clin Cardiol* 1998;**21**:858-61.
- Chan TC, Vilke GM, Pollack M, et al. Electrocardiographic manifestations: pulmonary embolism. *J Emerg Med* 2001;**21**:263-70.
- Falterman TJ, Martinez JA, Daberkow D, et al. Pulmonary embolism with ST segment elevation in leads V1-V4: case report and review of the literature regarding electrocardiographic changes in acute pulmonary embolism. *J Emerg Med* 2001;**21**:255-61.
- Meyer T, Binder L, Hruska N, et al. Cardiac troponin I elevation in acute pulmonary embolism is associated with right ventricular dysfunction. *J Am Coll Cardiol* 2000;**36**:632-6.
- Giannitsis E, Muller-Bardorff M, Kurowski V, et al. Independent prognostic value of cardiac troponin T in patients with confirmed pulmonary embolism. *Circulation* 2000;**102**:211-17.