IMAGE CHALLENGE

Splashed by a clear liquid

CLINICAL INTRODUCTION

A 35-year-old male presented with painful skin rash for 2 hours after a clear liquid splashed onto his legs when a car drove past him and squelched a roadside container on a street populated by small-sized factories. Burning sensation and pain developed

in 10 min. He called on the emergency room after irrigation. His trousers were intact, wet but not slippery, and had a pungent sour odour.

QUESTION

What is the best answer for the clear liquid?

- A. It is a strong alkaline fluid
- B. It is concentrated sulfuric acid
- C. It is concentrated hydrofluoric acid
- D. It is 50% of hydrogen peroxide

For the answer see page 475

Figure 1 Erythematous, centrally greyish changes can be seen over both thighs and knees. The patient felt a sharp pain, deep into the bone, leading him to tightly grasp his thighs to try to relieve it.

Figure 2 Necrotic changes with depression of the skin over the left knee.

Splashed by a clear liquid

From the question on page 453

ANSWER: C

The affected skin had a blue-greyish depression with surrounding erythema and blisters (figures 1 and 2), a picture of liquefactive necrosis. This was suggestive of an alkaline injury; however, the pH of liquid on the trouser, measured by litmus paper, was 1, suggesting an acid.

The free H^+ ion in strong sulfuric acid solution would result in not only coagulum formation on skin but also break down fabrics, causing burnt holes in clothing,¹ but the patient's trousers were intact. Concentrated hydrogen peroxide, a strong oxidant, could cause severe burns of mucosa but should not damage intact skin.² Hydrofluoric acid can cause similar skin changes to those seen, but because it is a weak acid, will not affect clothing. Fluoride ion was detected by ion exchange chromatography from a 5 cm×5 cm wet fabric cut from left knee area.

Hydrofluoric acid is widely used for etching, polishing or cleaning glass, ceramics and walls and rust removing. Its dissociated H^+ ion causes skin damage and leaves fluoride ion that penetrates deep into the tissues, binding to and sequestering Ca^{2+} and Mg^{2+} ions, producing extensive tissue destruction. Irrigation with copious amount of water and treatment with calcium or magnesium compounds intravenously or locally are the conventional therapy for hydrofluoric acid burns.³ In this case, the affected skin was then covered with calcium gluconate-soaked gauze, which drastically reduced the pain. Systemic toxicity like hyperkalaemia, hypocalcaemia or hypomagnesaemia was not observed, but the wound required skin graft because of extensive tissue necrosis.

Kai-Wei Yang,¹ Dong-Zong Hung,² Shih-Yu Chang³

¹Emergency Department, China Medical University Hospital, China Medical University, Taichung, Taiwan

²Division of Toxicology, China Medical University Hospital; Graduate Institute of Clinical Medical Science, College of Medicine, China Medical University, Taichung, Taiwan

³Department of Public Health, Chung Shan Medical University, Taichung, Taiwan

Correspondence to Dr Dong-Zong Hung, Division of Toxicology, China Medical University Hospital; 2 Yu-Der Road, Taichung City 40402, Taiwan; dzhung0224@gmail.com

Contributors K-WY was a trainee of toxicology, cared for this case and prepared the manuscript. S-YC performed the fluoride ion identification test. D-ZH cared for this case and prepared the manuscript.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; internally peer reviewed.

OPEN ACCESS

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

CrossMark

To cite Yang K-W, Hung D-Z, Chang S-Y. *Emerg Med J* 2017;**34**:475.

Accepted 27 July 2016

Published Online First 22 March 2017

Emerg Med J 2017;**34**:475. doi:10.1136/emered-2016-206063

REFERENCES

- 1 Dinis-Oliveira RJ, Carvalho F, Moreira R, *et al*. Clinical and forensic signs related to chemical burns: a mechanistic approach. *Burns* 2015;**41**:658–79.
- 2 Agency for Toxic Substances and Disease Registry. Medical management guidelines for hydrogen peroxide. Atlanta: Centers for Disease Control and Prevention, 2014. 20 p.
- 3 Burgher F, Mathieu L, Lati E, *et al*. Experimental 70% hydrofluoric acid burns: histological observations in an established human skin explants ex vivo model. *Cutan Ocul Toxicol* 2011;**30**:100–7.