

Comparison of qSOFA with current emergency department tools for screening of patients with sepsis for critical illness

Robert M Rodriguez,¹ John C Greenwood,² Thomas J Nuckton,³ Bryan Darger,¹ Frances S Shofer,⁴ Dawn Troeger,³ Soo Y Jung,⁴ Kelly G Speich,⁵ Joel Valencia,¹ J Hope Kilgannon,⁵ Danny Fernandez,¹ Brigitte M Baumann⁵

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/emmermed-2017-207383>).

¹Department of Emergency Medicine, University of California, San Francisco, San Francisco, California, USA

²Department of Emergency Medicine, Perelman School of Medicine, University of Pennsylvania, Philadelphia, Pennsylvania, USA

³Department of Medicine, Sutter Eden Medical Center, San Francisco, California, USA

⁴Department of Emergency Medicine, Hospital of the University of Pennsylvania, Philadelphia, Pennsylvania, USA

⁵Department of Emergency Medicine, Cooper Medical School of Rowan University, Camden, New Jersey, USA

Correspondence to

Dr Robert M Rodriguez, Department of Emergency Medicine, San Francisco General Hospital, San Francisco, CA 94110, USA; robert.rodriguez@ucsf.edu

Received 8 December 2017

Revised 6 March 2018

Accepted 28 March 2018

Published Online First

2 May 2018

ABSTRACT

Objective We sought to compare the quick sequential organ failure assessment (qSOFA) to systemic inflammatory response syndrome (SIRS), severe sepsis criteria and lactate levels for their ability to identify ED patients with sepsis with critical illness.

Methods We conducted this multicenter retrospective cohort study at five US hospitals, enrolling all adult patients admitted to these hospitals from their EDs with infectious disease-related illnesses from 1 January 2016 to 30 April 2016. We abstracted clinical variables for SIRS, severe sepsis and qSOFA scores, using values in the first 6 hours of ED stay. Our primary outcome was critical illness, defined as one or more of the composite outcomes of death, vasopressor use or intensive care unit (ICU) admission within 72 hours of presentation. We determined diagnostic test characteristics for qSOFA scores, SIRS, severe sepsis criteria and lactate level thresholds.

Main results Of 3743 enrolled patients, 512 (13.7%) had the primary composite outcome. The qSOFA scores were ≥ 1 , > 2 and 3 in 1839 (49.1%), 626 (16.7%) and 146 (3.9%) patients, respectively; 2202 (58.8%) met SIRS criteria and 1085 (29.0%) met severe sepsis criteria. qSOFA ≥ 1 and SIRS had similarly high sensitivity [86.1% (95% CI 82.8% to 89.0%) vs 86.7% (95% CI 83.5% to 89.5%)], but qSOFA ≥ 1 had higher specificity [56.7% (95% CI 55.0% to 58.5%) vs 45.6% (43.9% to 47.3%); mean difference 11.1% (95% CI 8.7% to 13.6%)]. qSOFA ≥ 2 had higher specificity than severe sepsis criteria [89.1% (88.0% to 90.2%) vs 77.5% (76.0% to 78.9%); mean difference 11.6% (9.8% to 13.4%)]. qSOFA ≥ 1 had greater sensitivity than a lactate level ≥ 2 (mean difference 24.6% (19.2% to 29.9%)).

Conclusion For patients admitted from the ED with infectious disease diagnoses, qSOFA criteria performed as well or better than SIRS criteria, severe sepsis criteria and lactate levels in predicting critical illness.

BACKGROUND

Early identification of patients who are critically ill or may have poor outcomes is a cornerstone of sepsis patient management. For nearly two decades, clinicians have used systemic inflammatory response syndrome (SIRS), sepsis, severe sepsis and septic shock as the primary terms (with corresponding criteria) to gauge severity of illness and guide patient management. Using these terms and criteria, most EDs have developed screening

Key messages

What is already known on this subject

- The Society of Critical Care Medicine and the European Society of Intensive Care Medicine have recently proposed the quick sequential organ failure assessment (qSOFA) criteria as replacements for traditional sepsis and severe sepsis criteria to identify critically ill patients outside of the intensive care unit.
- Prior examinations of the utility of qSOFA have focused primarily on 30-day mortality, which is not a pragmatic, ED-centred outcome.

What this study adds

- In this multicenter study, we evaluated qSOFA from a pragmatic, ED-centred vantage point.
- For patients admitted from the ED with infectious disease diagnoses, qSOFA criteria performed as well or better than SIRS criteria, severe sepsis criteria and lactate levels in predicting critical illness.

protocols designed to identify patients with sepsis, especially those who are critically ill and could benefit from prompt therapies, at triage and during early stages of their presentations.

Investigators and experts from the Society of Critical Care Medicine and the European Society of Intensive Care Medicine have recently proposed new criteria to define sepsis.¹ They propose that the quick sequential organ failure assessment (qSOFA) criteria, consisting of systolic blood pressure ≤ 100 mm Hg, respiratory rate ≥ 22 breaths/min and altered mental status (GCS < 15), can be used outside of the intensive care unit (ICU) to identify patients critically ill and in need of intensive care.² Patients are given one point for each criterion and patients with a qSOFA score of ≥ 2 are thought to be at high risk for poor outcomes.

Although qSOFA was not expressly developed to replace current ED sepsis screening tools, it nevertheless has been proposed to be used as such.^{3,4} Several concerns arise when attempting use qSOFA in this manner.⁵ First, given that qSOFA was developed in a heterogeneous group of patients (including many patients outside of the ED), it may not accurately reflect the ED population. Second, the primary



► <http://dx.doi.org/10.1136/emmermed-2018-207668>



To cite: Rodriguez RM, Greenwood JC, Nuckton TJ, et al. *Emerg Med J* 2018;**35**:350–356.



outcome driving the derivation and validation of qSOFA was mortality, which is not the only outcome measure under consideration in ED sepsis screening and disposition decisions. ED clinicians use sepsis screening tools for broader purposes, including identifying the sickest patients among the many they see, initiating critical, time-sensitive interventions and determining the need for ICU care. Finally, the primary outcome of mortality of 30 days is much longer than typical ED-centred outcomes. Patients who may not be critically ill on ED presentation, but subsequently become critically ill after days to weeks in the hospital and die later in the hospital course (or even as outpatients) would be considered to have this outcome, as would patients who receive comfort measures only. In general, triage decisions of ED patient with sepsis are expected to anticipate the need for ICU care and mortality over a much shorter time period after presentation—within approximately 72 hours of presentation.

The objective of this study was to assess the ability of qSOFA to detect patients with critical illness from a pragmatic, ED-centred vantage point. We therefore chose to compare it with the performance of the most commonly used current ED sepsis identification tools of SIRS, sepsis and severe sepsis criteria (former criteria) and lactate levels for predicting the need for critical care and mortality within the first 72 hours of admission to the hospital.

METHODS

This retrospective study was conducted at five US hospitals (three urban, county academic medical centres and two community hospitals) between 1 January and 30 April 2016. The mean number of beds of the EDs at these hospitals was 45 (range 24–70) and the mean number of annual visits was 58 700 (range 45 000–72 000). We included adult (age >17 years) patients admitted to an observation unit, inpatient ward or intensive care unit from the ED with a presumed infectious disease-related illness. We excluded patients who (1) were transferred from an outside hospital, (2) admitted to the hospital primarily for other reasons beyond their infectious disease illness (ie, a patient with cellulitis and severe trauma who was primarily admitted for the trauma would be excluded) and (3) left against medical advice (AMA) prior to hospital admission. None of the five hospitals had explicit ICU admission criteria.

Participants

Prior to abstraction of data, we generated an inclusive list of infectious disease-related admission diagnoses (see online Supplementary appendix). Each ED maintains an admission log that lists all admitted patients along with their admission diagnoses. During the study time period, investigators at each site reviewed these admission logs to identify potentially eligible subjects and then reviewed the ED charts of these patients to confirm that they were admitted with diagnoses on this list. As our goal was to test the performance of tools currently used for determining the severity of illness in patients presumed to have sepsis, we included patients who received an infectious disease-related diagnosis in the ED, even if those patients were ultimately found to not have infections during their hospital admission. Ambiguous inclusions or exclusions were adjudicated by consensus review of two additional investigators.

We defined our primary outcome of critical illness as a composite of ICU stay, receipt of vasopressor support or hospital death (excluding hospice or ‘do not resuscitate’ patients) within 72 hours of presentation (any of the three occurring within 72 hours qualified as a positive outcome). Vasopressor support included any intravenous infusions of norepinephrine,

Box 1 Systemic inflammatory response (SIRS), sepsis, severe sepsis and quick sequential organ failure assessment (qSOFA) criteria definitions

SIRS

Two or more of:

- ▶ Temperature >38°C or <36°C.
- ▶ Heart rate >90/min.
- ▶ Respiratory rate >20/min or arterial partial pressure of carbon dioxide <32 mm Hg (4.3 kPa).
- ▶ White cell count >12×10⁹/L or <4×10⁹/L or >10% immature bands.

Sepsis

- ▶ Criteria meeting SIRS.
- ▶ Source of infection.

Severe sepsis=sepsis induced tissue hypoperfusion or organ dysfunction

- ▶ Criteria meeting sepsis (SIRS+source of infection) and any of the following:
 - ▶ Lactic acidosis ≥2.2 mmol/L.
 - ▶ Urine output <0.5 mL/kg/hour for more than 2 hours despite adequate fluid resuscitation.
 - ▶ Acute lung injury with PaO₂/fractional inspired oxygen (FIO₂) <250 in the absence of pneumonia as infection source.
 - ▶ Acute lung injury with PaO₂/FIO₂ <250 in the presence of pneumonia as infection source.
 - ▶ Creatinine >2.0 mg/dL.
 - ▶ Bilirubin >2 mg/dL.
 - ▶ Platelet count <100 000 μL.
 - ▶ Coagulopathy (international normalised ratio>1.5).

qSOFA

- ▶ Systolic blood pressure <100 mm Hg.
- ▶ Respiratory rate ≥22 breaths per min.
- ▶ Altered mental status (GCS <15).

epinephrine, dopamine, phenylephrine, dobutamine and vasopressin. Single push-doses of vasopressors were not included.

We adhered to STAndards for the Reporting of Diagnostic accuracy studies (STARD) guidelines for diagnostic test studies and used standardised abstraction methods to record qSOFA, SIRS and severe sepsis criteria from patients’ ED charts and electronic health records.^{6,7} Specifically, we generated a structured abstraction data collection form, held abstractor orientation meetings and conducted preliminary abstraction of 10 charts with abstractors to ensure abstractor consistency. We also held regular meetings to review progress and conducted frequent audits of abstractor data.

See **box 1** for criteria comprising SIRS, sepsis, severe sepsis and qSOFA.^{8–11} Given that infectious disease-related illness was an inclusion criterion for the study, all patients meeting SIRS criteria also met sepsis criteria and were considered one group: SIRS/sepsis. We used the most abnormal values in the first 6 hours of ED stay for all data points, that is, highest respiratory rate, lowest systolic blood pressure, lowest GCS and highest or lowest temperatures. Because most decisions regarding disposition are made within 6 hours, we did not include vital signs beyond this time point. Respiratory rates at triage were recorded from automated pulse oximetry devices. GCS were obtained from both nursing and physician assessments—at four sites all patients were given GCS scores at triage by nursing. We also recorded first venous lactate levels if performed within 6 hours

of ED presentation. Missing data elements were recorded as missing, but assumed to be normal (non-aberrant) in analyses.

To minimise bias, we abstracted outcomes blinded to qSOFA, SIRS and sepsis criteria. To measure interabstractor reliability, we conducted dual, independent abstraction of 200 subjects and calculated a kappa statistic of agreement.

Sample size considerations

The sample size to achieve our objectives was governed by the precision (width of (CIs)) around the point estimate of sensitivity of sepsis criteria and qSOFA for the primary composite outcome. From prior studies, we estimated that these instruments would have approximately 85% sensitivity for this combined outcome.^{12–14} We sought 3% CIs around this point estimate and expected that approximately 15% of enrolled patients would have our primary outcome; we therefore estimated the need to enrol approximately 3533 patients in this study. From pilot data, we estimated that we would enrol approximately 1000 patients per month at the five sites and therefore set our study time period from January to 30 April 2016.

Data management

We managed data using Research Electronic Data Capture hosted by the University of California San Francisco and transferred de-identified data to Excel spreadsheets (Microsoft Office Professional Plus 2010) for sorting and analysis.

Primary outcome analysis

We determined the sensitivity, specificity, positive and negative predictive value and positive and negative likelihood ratios for our composite outcome for qSOFA scores of ≥ 1 , > 2 and 3, SIRS/sepsis criteria, severe sepsis criteria and lactate levels of ≥ 2 and > 4 . We also generated three receiver operating characteristic (ROC) curves and calculated the area under the ROC curve (AUROC) for these same qSOFA cut-off points, SIRS/

sepsis and severe sepsis criteria and lactate level cut-off points. For lactate, we used cut points of 2 and 4, because these cut points are commonly used as thresholds for sepsis interventions. Because the hypotension element in qSOFA could inflate its performance with vasopressor as an outcome and because the qSOFA score could affect decisions about ICU disposition (incorporation bias), we also calculated qSOFA performance for each of the individual elements (ICU stay, mortality, and vasopressor use) as single (non-composite) outcomes.

RESULTS

Of 21038 patients admitted from the EDs of the study sites, 3869 were deemed to have infectious disease-related illnesses as one of their primary reasons for admission. Subsequent review of 223 ambiguous cases by two authors excluded 126 patients, leaving 3743 patients for full analysis (figure 1). Median age of the cohort was 58 years (IQR (44–72)) and their hospital mortality was 1.9% with a median of 5 days (IQR 2–9) until death. Other patient characteristics are summarised in table 1.

Missing qSOFA, SIRS and severe sepsis data elements comprised less than 0.05% of the total; GCS was the most commonly absent criterion of the qSOFA—missing in four (0.1%) of cases. The kappa statistics of agreement with 95% CIs were 0.987 (95% CI 0.97 to 1.00) for qSOFA criteria and 0.972 (95% CI 0.944 to 0.999) for the composite outcome, indicating extremely high and reliable interabstractor agreement.

Among the cohort, 2202 (58.8%) and 1085 (29.0%) met SIRS/sepsis and severe sepsis criteria, respectively. The qSOFA scores were ≥ 1 , > 2 and 3 in 1839 (49.1%), 626 (16.7%) and 146 (3.9%) patients, respectively. Of the 2584 patients who had lactate levels measured in the ED, 888 (34.4%) had a lactate level ≥ 2 and 205 (7.9%) had a lactate level ≥ 4 .

Main analysis

The primary outcome occurred in 512 (13.7%) patients. Among this group, 493 (96.3%) were admitted to an ICU, 219 (42.8%) patients received a vasopressor and 45 (8.8%) patients died within 72 hours of ED presentation; 216 (42.2%) patients had more than one of these outcomes. Mechanical ventilation was delivered to 271 (55.0%) ICU patients in the first 72 hours.

For the primary composite outcome, sensitivity and negative likelihood ratios were highest and nearly identical for qSOFA ≥ 1 and SIRS/sepsis: sensitivity 86.1% (95% CI 82.8% to 89.0%) versus 86.7% (95% CI 83.5% to 89.5%) and likelihood ratio negative 0.24 (95% CI 0.20 to 0.30) versus 0.29 (95% CI 0.23 to 0.36). However, qSOFA ≥ 1 had greater specificity than SIRS/sepsis [56.7% (95% CI 55.0 to 58.5%) vs 45.6% (95% CI 43.9% to 47.3%); mean difference 11.1% (95% CI 8.7% to 13.6%)]. A qSOFA ≥ 2 had higher specificity than severe sepsis criteria [89.1% (95% CI 88.0% to 90.2%) vs 77.5% (95% CI 76.0% to 78.9%); mean difference 11.6% (95% CI 9.8% to 13.4%)] at the expense of lower sensitivity [mean difference 16.2% (95% CI 10.3% to 22.0%)]. A qSOFA=3 had the highest specificity but lower sensitivity than SIRS/sepsis or severe sepsis. A lactate level ≥ 2 had much lower sensitivity than qSOFA ≥ 1 (mean difference 24.6% (95% CI 19.2% to 29.9%)). Like qSOFA=3, a lactate level ≥ 4 had high specificity but very low sensitivity (see table 2).

ROC curves are presented in figure 2; qSOFA had significantly higher AUROC (0.788) than SIRS/sepsis or severe sepsis criteria (0.754) or lactate level thresholds (0.763), $p=0.0026$ and $p<0.001$, respectively.




Figure 1 Screening and enrollment.

Table 1 Patient characteristics, n=3743

Median age (IQR)	58 (44–72)
Male sex, number (%)	1915 (51.2)
Race, number (%)	
Caucasian (non-hispanic)	1544 (41.3)
African American	1143 (30.5)
Hispanic	554 (14.8)
Asian American	175 (4.7)
Native American	18 (0.5)
Other	209 (5.6)
Unknown	100 (2.7)
Source of infection, number (%)	
Pneumonia or pulmonary	1081 (28.9)
Skin or soft-tissue infection	948 (25.3)
Urinary tract infection	548 (14.6)
Intra-abdominal infection	529 (14.1)
Unknown source of infection	507 (13.5)
Other (otorhinolaryngological, gyne, etc)	84 (2.2)
Central nervous system infection	45 (1.2)

Sensitivity analyses

Removal of the vasopressor criterion from the composite outcome had very little effect on qSOFA performance. A qSOFA score of ≥ 1 had a sensitivity of 86.6% (95% CI 83.2% to

89.4%) and a specificity of 56.6% (95% CI 54.9% to 58.3%) for the composite outcome of ICU admission or mortality. A qSOFA score of ≥ 2 had a sensitivity of 54.0% (95% CI 49.5% to 58.4%) and a specificity of 89.0% (95% CI 87.9% to 90.0%).

In tables 3–5, we present the performance characteristics of qSOFA for predicting the single outcomes of mortality, vasopressors and ICU admission within 72 hours, respectively. For all of these outcomes, a qSOFA score of 1 had high sensitivity ($>84\%$) and low specificity ($<57\%$), and a qSOFA score of 3 had low sensitivity ($<43\%$) and high specificity ($<96\%$).

DISCUSSION

Identifying patients with critical illness is paramount in the ED care of patients with infectious disease processes. Highly sensitive prediction tools will allow clinicians to detect all (or nearly all) critically ill patients early in their course to provide time-sensitive interventions such as fluid resuscitation, early antibiotics and timely vasopressor administration.^{15–18} Highly specific predictive tools allow for more judicious disposition decisions and expenditure of resources, particularly ICU beds. In this multicenter study assessing the utility of qSOFA to predict the need for ICU care, vasopressors or mortality within 72 hours of admission from the ED, we found that qSOFA criteria had slightly better performance than traditional SIRS and sepsis criteria. For ruling out critical illness, qSOFA ≥ 1 had equivalent sensitivity and negative predictive value to SIRS criteria,

Table 2 qSOFA, SIRS, severe sepsis and lactate level: screening performance characteristics for composite outcome

Criteria	Sensitivity, % (95% CI)	Specificity, % (95% CI)	Negative predictive value, % (95% CI)	Negative likelihood ratio (95% CI)	Positive predictive value, % (95% CI)	Positive likelihood ratio (95% CI)
SIRS TP=444 TN=1473 FP=1758 FN=68	86.7 (83.5 to 89.5)	45.6 (43.9 to 47.3)	95.6 (94.5 to 96.4)	0.29 (0.23 to 0.36)	20.2 (19.4 to 20.9)	1.6 (1.5 to 1.7)
Severe sepsis TP=357 TN=2503 FP=728 FN=155	69.7 (65.5 to 73.7)	77.5 (76.0 to 78.9)	94.2 (93.4 to 94.9)	0.39 (0.34 to 0.45)	32.9 (31.0 to 34.8)	3.1 (2.8 to 3.4)
qSOFA ≥ 1 TP=441 TN=1833 FP=1398 FN=71	86.1 (82.8 to 89.0)	56.7 (55.0 to 58.5)	96.3 (95.4 to 97.0)	0.24 (0.20 to 0.30)	24.0 (23.0 to 25.0)	2.0 (1.9 to 2.1)
qSOFA ≥ 2 TP = 274 TN=2879 FP=352 FN=238	53.5 (49.0 to 57.9)	89.1 (88.0 to 90.2)	92.4 (91.7 to 93.0)	0.52 (0.48 to 0.57)	43.8 (40.7 to 46.9)	4.9 (4.3 to 5.6)
qSOFA ≥ 3 TP=90 TN=3175 FP=56 FN=422	17.6 (14.4 to 21.2)	98.3 (97.8 to 98.7)	88.3 (87.8 to 88.7)	0.84 (0.81 to 0.87)	61.6 (53.9 to 68.9)	10.1 (7.4 to 14.0)
Lactate ≥ 2 TP=286 TN=1517 FP=602 FN=179	61.5 (56.9 to 66.0)	71.6 (69.6 to 73.5)	89.5 (88.3 to 90.5)	0.54 (0.48 to 0.61)	32.2 (30.1 to 34.4)	2.2 (2.0 to 2.4)
Lactate ≥ 4 TP=121 TN=2035 FP=84 FN=344	26.0 (22.1 to 30.3)	96.0 (95.1 to 96.3)	85.5 (84.9 to 86.2)	0.77 (0.73 to 0.81)	59.0 (52.6 to 65.1)	6.6 (5.1 to 8.5)

qSOFA, quick sequential organ failure assessment; SIRS, systemic inflammatory response syndrome; FN, false negative; FP, false positive; TN, true negative; TP, true positive.




Figure 2 ROC for qSOFA, lactate and SIRS/Severe Sepsis Criteria.

with slightly higher specificity and positive predictive value. For ruling in critical illness, we found that a qSOFA ≥ 2 offers superior discriminative performance (higher specificity and positive likelihood ratio) for short-term critical illness than formerly used SIRS and severe sepsis criteria. When combining overall prognostic accuracy in terms of ROC curve comparisons and AUROC, we found that qSOFA had better discriminative capacity than SIRS/severe sepsis criteria and lactate levels.

Complicated formulas and instruments requiring extensive laboratory testing are less useful than simple ones from a practical implementation standpoint in the ED. With only three very straightforward clinical criteria that are easily assessed in all ED patients and no need for laboratory testing, qSOFA as a risk-stratification tool has substantial pragmatic appeal compared with the criteria for SIRS, sepsis and severe sepsis criteria.

Evaluations of qSOFA have not specifically addressed the utility of qSOFA for identifying patients who will have critical outcomes

in the first 72 hours after admission from the ED.^{21-14 19 20} These studies have nevertheless yielded mixed results. Most investigators have found that qSOFA was a better predictor of 30-day mortality than SIRS or severe sepsis criteria.^{21-14 19} However, other investigators found that although qSOFA and traditional SIRS criteria had similar discrimination for predicting organ dysfunction, qSOFA ≥ 2 was much less sensitive for 30-day mortality than SIRS.²⁰

Given recent concerns of inadequate sensitivity (87.9%) of SIRS criteria for critical illness, the 86.1% sensitivity of a qSOFA ≥ 1 may seem insufficient.²¹ In this regard, other risk stratification tools for critical illness have been proposed as useful alternatives to current standard critical illness screening tools.²¹ The National Early Warning Score (NEWS) and Modified Early Warning Score (MEWS) criteria, which also consist of readily available physiological parameters, have performed well when used to identify critical illness in heterogeneous patient populations.²²⁻²⁴ In a single-site validation study, Churpek *et al*, found that both NEWS and MEWS were more accurate than qSOFA ≥ 2 at predicting early mortality and need for ICU transfer.¹³

Despite the initial reluctance of emergency physicians to embrace qSOFA as a substitute for SIRS, sepsis and severe sepsis criteria, we believe our findings are in line with those presented by the Sepsis-3 task force and that qSOFA may be a superior indicator of critical illness in these patients. As with all scoring systems, we believe that qSOFA should not replace clinical judgment, but should aid clinicians in both risk stratification and clinical decision-making. Given the superior performance over previous methods of identifying patients with infectious disease who are critically ill, the early use of qSOFA may help improve the identification of those who need timely interventions, further diagnostic testing and potentially higher intensity care. Future investigations should assess these tools prospectively.

Limitations

Although our study design was retrospective, less than 1% of our eligible subjects had missing qSOFA or outcome data and we had excellent high interabstractor agreement. The study only included patients admitted to the hospital and not those discharged, and thus did not assess qSOFA's ability to detect critical illness among all potentially infected patients. Although our use of admission logs and ED diagnoses to screen for eligible patients could lead to selection bias, this method is ED centred and practical. Use of blood cultures as a required inclusion

Table 3 qSOFA: screening performance characteristics for mortality at 72 hours

Criteria	Sensitivity, % (95% CI)	Specificity, % (95% CI)	Negative predictive value, % (95% CI)	Negative likelihood ratio (95% CI)	Positive predictive value, % (95% CI)	Positive likelihood ratio (95% CI)
qSOFA ≥ 1 TP=38 TN=1896 FP=1802 FN=7	84.4 (69.9 to 93.0)	51.3 (49.6 to 52.9)	99.6 (99.2 to 99.8)	0.30 (0.15 to 0.60)	2.1 (1.5 to 2.9)	1.7 (1.5 to 2.0)
qSOFA ≥ 2 TP=29 TN=3101 FP=597 FN=16	64.4 (48.7 to 77.7)	83.9 (82.6 to 85.0)	99.5 (99.1 to 99.7)	0.42 (0.29 to 0.63)	4.6 (3.2 to 6.7)	4.0 (3.2 to 5.0)
qSOFA ≥ 3 TP=17 TN=3569 FP=129 FN=28	37.8 (24.2 to 53.5)	96.5 (95.9 to 97.1)	99.2 (98.9 to 99.5)	0.64 (0.51 to 0.81)	11.6 (7.1 to 18.3)	10.8 (7.2 to 16.3)

qSOFA, quick sequential organ failure assessment; TP, true positive; TN, true negative; FP, false positive; FN, false negative.

Table 4 qSOFA: screening performance characteristics for vasopressors at 72 hours

Criteria	Sensitivity, % (95% CI)	Specificity, % (95% CI)	Negative predictive value, % (95% CI)	Negative likelihood ratio (95% CI)	Positive predictive value, % (95% CI)	Positive likelihood ratio (95% CI)
qSOFA \geq 1 TP=206 TN=1889 FP=1634 FN=14	93.6 (89.3 to 96.3)	53.6 (52.0 to 55.3)	99.3 (98.7 to 99.6)	0.12 (0.07 to 0.20)	11.2 (9.8 to 12.7)	2.0 (1.9 to 2.1)
qSOFA \geq 2 TP=158 TN=3055 FP=468 FN=62	71.8 (65.3 to 77.6)	86.7 (85.5 to 87.8)	98.0 (97.4 to 98.5)	0.32 (0.26 to 0.40)	25.2 (21.9 to 28.9)	5.4 (4.8 to 6.1)
qSOFA \geq 3 TP=67 TN=3506 FP=79 FN=91	42.4 (34.7 to 50.5)	97.8 (97.2 to 98.2)	97.5 (96.9 to 97.9)	0.59 (0.52 to 0.67)	45.9 (37.7 to 54.3)	19.2 (14.5 to 25.6)

qSOFA, quick sequential organ failure assessment; TP, true positive; TN, true negative; FP, false positive; FN, false negative.

criterion would unnecessarily exclude many patients who may still be septic. For example, current Infectious Disease Society of America guidelines recommend against routine blood cultures in patients with skin and soft-tissue infections.²⁵

Incorporation bias may have affected the predictive performance of SIRS and sepsis criteria, as well as lactate levels, for the composite outcome by virtue of the fact that these criteria may have been used by clinicians in real time to determine the need for ICU care. Because we examined data from patients before the publication of the qSOFA score, SIRS and severe sepsis screening performance would thereby likely be inflated to a greater extent than qSOFA. Given that one of the qSOFA criteria is hypotension (systolic blood pressure <100), our inclusion of vasopressor use as part of our composite outcome may have biased our results in favour of qSOFA. However, when removing this outcome, qSOFA retained superior predictive performance. Sensitivity analyses of individual outcomes (mortality alone, vasopressor alone) demonstrated similar test characteristics to the composite outcome. Although the original qSOFA cut-off point was 2, our intent was to analyse outcomes according to the full spectrum of qSOFA scores. Dichotomisation of lactate may have also decreased its discriminatory ability. We found lower overall hospital mortality than that seen in the qSOFA study cohorts^{1,2};

however, this was an expected finding given that we included all patients admitted from the ED rather than only patients who had ICU stays.

CONCLUSIONS

In this multicenter study of patients admitted with infectious disease diagnoses, we found that qSOFA criteria had better performance for predicting early critical illness than SIRS, sepsis or severe sepsis criteria or lactate levels. Given that qSOFA criteria are easy to perform without the need for blood tests, they may be preferred over current ED risk stratification tools in the ED.

Contributors RMR, JCG, TJN, BD and BMB contributed to the study design. RMR, JCG, BD, TJN, BMB and FSS were involved in the data analysis. All authors contributed to the study implementation/data acquisition, manuscript preparation and revision.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent Not required.

Ethics approval IRBs of UCSF, University of Pennsylvania and Sutter.

Table 5 qSOFA: screening performance characteristics for intensive care unit at 72 hours

Criteria	Sensitivity % (95% CI)	Specificity % (95% CI)	Negative predictive value % (95% CI)	Negative likelihood ratio (95% CI)	Positive predictive value % (95% CI)	Positive likelihood ratio (95% CI)
qSOFA \geq 1 TP=430 TN=1840 FP=1410 FN=63	87.2 (83.9 to 90.0)	56.6 (54.9 to 58.3)	96.7 (95.8 to 97.4)	0.23 (0.18 to 0.28)	23.4 (21.5 to 25.4)	2.0 (1.9 to 2.1)
qSOFA \geq 2 TP=268 TN=2892 FP=358 FN=225	54.4 (49.8 to 58.8)	89.0 (87.8 to 90.0)	92.8 (91.8 to 93.7)	0.51 (0.47 to 0.56)	42.8 (38.9 to 46.8)	4.9 (4.3 to 5.6)
qSOFA \geq 3 TP=87 TN=3191 FP=59 FN=406	17.6 (14.4 to 21.4)	98.2 (97.6 to 98.6)	88.7 (87.6 to 89.7)	0.84 (0.81 to 0.87)	59.6 (51.1 to 67.5)	9.7 (7.1 to 13.3)

qSOFA, quick sequential organ failure assessment; TP, true positive; TN, true negative; FP, false positive; FN, false negative.

Provenance and peer review Not commissioned; externally peer reviewed.

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2018. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- Singer M, Deutschman CS, Seymour CW, *et al.* The third international consensus definitions for sepsis and septic shock (Sepsis-3). *JAMA* 2016;315:801–10.
- Seymour CW, Liu VX, Iwashyna TJ, *et al.* Assessment of clinical criteria for sepsis. *JAMA* 2016;315:762.
- Franchini S, Duca A. qSOFA should replace SIRS as the screening tool for sepsis. *Crit Care* 2016;20:409.
- April MD, Aguirre J, Tannenbaum LI, *et al.* Sepsis clinical criteria in emergency department patients admitted to an intensive care unit: an external validation study of quick sequential organ failure assessment. *J Emerg Med* 2017;52.
- Simpson SQ. New sepsis criteria: a change we should not make. *Chest* 2016;149:1117–8.
- Bossuyt PM, Reitsma JB, Bruns DE, *et al.* Towards complete and accurate reporting of studies of diagnostic accuracy: The STARD Initiative. *Ann Intern Med* 2003;138:40–4.
- Gilbert EH, Lowenstein SR, Koziol-McLain J, *et al.* Chart reviews in emergency medicine research: Where are the methods? *Ann Emerg Med* 1996;27:305–8.
- Anon. American College of Chest Physicians/Society of Critical Care Medicine Consensus Conference: definitions for sepsis and organ failure and guidelines for the use of innovative therapies in sepsis. *Crit Care Med* 1992;20:864–74.
- Soong J, Soni N. Sepsis: recognition and treatment. *Clin Med* 2012;12:276–80.
- Dellinger RP, Levy MM, Rhodes A, *et al.* Surviving sepsis campaign: international guidelines for management of severe sepsis and septic shock: 2012. *Crit Care Med* 2013;41:580–637.
- Vincent JL, de Mendonça A, Cantraine F, *et al.* Use of the SOFA score to assess the incidence of organ dysfunction/failure in intensive care units: results of a multicenter, prospective study. Working group on “sepsis-related problems” of the European Society of Intensive Care Medicine. *Crit Care Med* 1998;26:1793–800.
- Wang JY, Chen YX, Guo SB, *et al.* Predictive performance of quick sepsis-related organ failure assessment for mortality and ICU admission in patients with infection at the ED. *Am J Emerg Med* 2016;34:1788–93.
- Churpek MM, Snyder A, Han X, *et al.* qSOFA, SIRS, and early warning scores for detecting clinical deterioration in infected patients outside the ICU. *Am J Respir Crit Care Med* 2016.
- Williams JM, Greenslade JH, McKenzie JV, *et al.* SIRS, qSOFA and organ dysfunction: insights from a prospective database of emergency department patients with infection. *Chest* 2016.
- Yealy DM, Kellum JA, Huang DT, *et al.* A randomized trial of protocol-based care for early septic shock. *N Engl J Med* 2014;370:1683–93.
- Peake SL, Delaney A, Bailey M, *et al.* Goal-directed resuscitation for patients with early septic shock. *N Engl J Med* 2014;371:1496–506.
- Mouncey PR, Osborn TM, Power GS, *et al.* Protocolised Management In Sepsis (ProMISe): a multicentre randomised controlled trial of the clinical effectiveness and cost-effectiveness of early, goal-directed, protocolised resuscitation for emerging septic shock. *Health Technol Assess* 2015;19:1–150.
- Chen YX, Wang JY, Guo SB. Use of CRB-65 and quick sepsis-related organ failure assessment to predict site of care and mortality in pneumonia patients in the emergency department: a retrospective study. *Crit Care* 2016;20:167.
- Freund Y, Lemachatti N, Krastinova E, *et al.* Prognostic accuracy of sepsis-3 criteria for in-hospital mortality among patients with suspected infection presenting to the emergency department. *JAMA* 2017;317:301–8.
- Innocenti F, Tozzi C, Donnini C, *et al.* SOFA score in septic patients: incremental prognostic value over age, comorbidities, and parameters of sepsis severity. *Intern Emerg Med*. In Press. 2018;13.
- Kaukonen KM, Bailey M, Pilcher D, *et al.* Systemic inflammatory response syndrome criteria in defining severe sepsis. *N Engl J Med* 2015;372:1629–38.
- National Early Warning Score (NEWS). Standardizing the assessment of acute-illness severity in the NHS. <https://www.rcplondon.ac.uk/projects/outputs/national-early-warning-score-news> (accessed 14 Feb 2017).
- Gardner-Thorpe J, Love N, Wrightson J, *et al.* The value of Modified Early Warning Score (MEWS) in surgical in-patients: a prospective observational study. *Ann R Coll Surg Engl* 2006;88:571–5.
- Subbe CP, Kruger M, Rutherford P, *et al.* Validation of a modified Early Warning Score in medical admissions. *QJM* 2001;94:521–6.
- Stevens DL, Bisno AL, Chambers HF, *et al.* Practice guidelines for the diagnosis and management of skin and soft tissue infections: 2014 update by the Infectious Diseases Society of America. *Clin Infect Dis* 2014;59:e10–52.