



Abstract 015 Figure 1 Where care provider emotion can impact

was used to recruit participants, with each group containing members of the same or similar grade and occupation. Consultant, higher specialty trainee, junior doctor and nursing focus groups were undertaken. Following transcription, data were coded and thematically analysed to arrive at key concepts.

Conclusions Six main themes were identified: ‘ED team rapport’ referring to the personalities and attitudes of those working that day; ‘achievement’ with successful task completion, skill use, being thanked and constructive educational opportunities generating positive emotions and lack of this feeling generating the converse; ‘interpersonal interactions’ where the negative impact of incivility from staff or patients was highlighted; ‘equipment/infrastructure’ showcasing how frustration manifests when either fails; ‘the open and the close’ representing the impact of how the tone set in hand-over influences affect and finally, the self-explanatory ‘a bad day outside work can influence that inside.’

This study illustrates the importance of recognising personal ‘wins’ whether that be a well-managed patient or successfully recognised teaching opportunity. It adds impetus to the campaign against incivility - reaffirming the negative effect rudeness has on affect. On a personal level, it highlights that we can take responsibility to ‘choose our own weather’ as a team member or leader in being a colleague that others enjoy working with. This may improve outcomes for all.


016

AN EVALUATION OF THE IMPACT OF THE NICE HEAD INJURY GUIDELINES ON INPATIENT MORTALITY FROM TRAUMATIC BRAIN INJURY: AN INTERRUPTED TIME SERIES ANALYSIS


^{1,2}Carl Marincowitz, ³Fiona Lecky, ²Victoria Allgar, ⁴Trevor Sheldon. ¹Hull and East Yorkshire Hospitals NHS Trust; ²Hull York Medical School; ³University of Sheffield; ⁴University of York

10.1136/emered-2019-RCEM.16

Background Traumatic brain injury (TBI) is the commonest cause of death and disability in UK Citizens aged 1–40. In England three (National Institute of Health and Care Excellence - NICE) guidelines have been implemented to improve



Abstract 016 Figure 1 The impact of the NICE head injury guidelines on monthly TBI mortality rate per 100 000 population



Abstract 016 Figure 2 The impact of the NICE head guidelines on monthly TBI hospital admissions per 100,000 population

TBI outcomes. All guidelines recommended increased CT imaging. The second guideline recommended the management of patients with severe TBI in specialist neuroscience centres.

This study uses national data and interrupted time series analysis to assess the impact of the NICE guidelines.

Individual level Office of National Statistics (ONS) cause of death data linked to Hospital Episode Statistics for inpatient admissions in England between 1998–2017 were used to estimate the monthly population mortality and admission rate for TBI.

An interrupted time series analysis was conducted with intervention points when each guideline was introduced. The analysis was stratified by guideline recommendation specific age groups (0–15, 16–64 and 65+).

The monthly TBI mortality and admission rate in the 65+ age group increased from 0.5 to 1.5 and 10 to 30 per 100,000 population respectively. The increasing mortality rate was unaffected by the introduction any of the guidelines.

The introduction of the 2nd NICE Head Injury guideline was associated with a significant reduction in the monthly TBI mortality rate in 16–64 age group (–0.005; 95% CI: –0.002 to –0.007).

In the 0–15 age group the TBI mortality rate fell from around 0.05 to 0.01 per 100 000 population, the trend was unaffected by the guidelines.

Conclusion The introduction of NICE head injury guidelines was associated with reduced population based mortality rates after specialist care was recommended for severe TBI. The improvement was solely observed in 16–64 year olds.

The cause of the observed increased admission and mortality rate in those 65+ and potential treatments for TBI in this age group requires further investigation.

017 BYPASSING THE NEAREST EMERGENCY DEPARTMENT FOR A MORE DISTANT NEUROSURGICAL CENTRE IN TRAUMATIC BRAIN INJURY PATIENTS

¹Callum Prosser, ²David Edwards, ¹Fiona Lecky, ³Omar Bouamra. ¹University of Sheffield; ²National Health Service; ³Faculty of Biology, Medicine and Health The University of Manchester

10.1136/emered-2019-RCEM.17

Background The recent introduction of major trauma networks throughout England in 2012 has changed how patients with suspected traumatic brain injury (TBI) are managed at the scene of injury. Selecting certain head trauma patients with suspected TBI for bypass to a more distant specialist neurological centre (SNC) is the networks function but may delay resuscitation whilst expediting neurosurgical/critical care. This comparative effectiveness research study analysed the impact of this strategy on the risk adjusted survival rates of patients confirmed to have a TBI on brain CT scan.

Method and results The study employed data from the Trauma Audit and Research Network. Adult patients with a TBI on CT scan were included if they presented between June 2015 to February 2016 to SNCs or non-specialist acute hospitals (NSAH) in the North of England (South Cumbria, Lancashire and the North East Region). Patients were identified as having bypassed a nearer NSAH emergency department (ED) to a SNC using google maps enabling exclusion of patients whose nearest ED was within a SNC. Their risk adjusted survival was compared to TBI patients who received primary treatment at a NSAH with subsequent secondary transfer to a SNC or

who remained at the NSAH until discharge or death. A multivariate logistic regression model predicting survival after TBI (Ps14ⁿ) was utilised to adjust for variation in casemix between the cohorts.

Conclusions 84 of 339 (25%) of TBI patients bypassed a nearer NSAH to a SNC, whilst 75% received primary treatment at an NSAH (n=255). There was no significant difference in the standardised excess survival rate between the two cohorts; shown as +2.55% for bypass (–5.09% to +10.20%) versus –1.49% for non-bypass (–5.34% to +2.36%).

No significant survival benefit was identified for TBI patients who bypassed the nearest ED compared to those receiving treatment at the nearest NSAH.


018 CLINICIAN PREDICTION OF CARDIAC ARRHYTHMIA IN PATIENTS PRESENTING TO THE ED WITH PALPITATION OR PRE-SYNOPE

¹Matt Reed, ²Neil Grubb, ²Chris Lang, ³Rachel O'Brien, ⁴Kirsty Simpson, ³Mia Padarenga, ¹Alison Grant, ⁵Sharon Tuck, ⁶Liza Keating, ⁷Frank Coffey, ⁸Lucy Jones, ⁹Tim Harris, ¹⁰Gavin Lloyd, ¹¹James Gagg, ¹²Jason Smith. ¹Emergency Medicine Research Group Edinburgh (EMERGE), Royal Infirmary of Edinburgh; ²Department of Cardiology, Royal Infirmary of Edinburgh; ³Emergency Medicine Research Group Edinburgh (EMERGE), Department of Emergency Medicine, Royal Infirmary of Edinburgh; ⁴Emergency Medicine Research Group Edinburgh (EMERGE); ⁵Edinburgh Clinical Research Facility, Epidemiology and Statistics Core, University of Edinburgh; ⁶Emergency Department, Royal Berkshire NHS Foundation Trust; ⁷DREEAM – Department of Research and Education in Emergency Medicine, Acute Medicine and Major Trauma; ⁸Chesterfield Royal Hospital NHS Foundation Trust; ⁹Queen Mary's University; ¹⁰Royal Devon and Exeter Hospital; ¹¹Musgrove Park Hospital; ¹²Emergency Department, University Hospitals Plymouth NHS Trust

10.1136/emered-2019-RCEM.18

The IPED study showed that use of a smartphone-based event recorder in ED patients presenting with palpitation or pre-syncope, increased the number of patients in whom an ECG was captured during symptoms over five-fold to more than 55% at 90 days (Reed MJ *et al.* Lancet eClinical Medicine 2019; 8: 37–46).

This pre-planned analysis looked at the ability of ED clinicians to predict cardiac arrhythmia in patients presenting to the ED with palpitation or pre-syncope.



Abstract 018 Figure 1 ROC analysis of ED clinician likelihood rating for symptomatic cardiac dysrhythmia at 90 days