Occupational exposures and programmatic response to COVID-19 pandemic: an emergency medical services experience

David L Murphy ¹, Leslie M Barnard,² Christopher J Drucker,² Betty Y Yang,¹ Jamie M Emert,² Leilani Schwarcz,² Catherine R Counts,¹ Tracie Y Jacinto,² Andrew M McCoy,^{1,3} Tyler A Morgan,⁴ Jim E Whitney,⁵ Joel V Bodenman,⁶ Jeffrey S Duchin,^{7,8} Michael R Sayre,^{1,9} Thomas D Rea^{2,8}

ABSTRACT

Rigorous assessment of occupational COVID-19 risk and personal protective equipment (PPE) use is not well-described. We evaluated 9-1-1 emergency medical services (EMS) encounters for patients with COVID-19 to assess occupational exposure, programmatic strategies to reduce exposure and PPE use. We conducted a retrospective cohort investigation of laboratory-confirmed patients with COVID-19 in King County, Washington, USA, who received 9-1-1 EMS responses from 14 February 2020 to 26 March 2020. We reviewed dispatch, EMS and public health surveillance records to evaluate the temporal relationship between exposure and programmatic changes to EMS operations designed to identify high-risk patients, protect the workforce and conserve PPE. There were 274 EMS encounters for 220 unique COVID-19 patients involving 700 unique EMS providers with 988 EMS person-encounters. Use of 'full' PPE including mask (surgical or N95), eye protection, gown and gloves (MEGG) was 67%. There were 151 person-exposures among 129 individuals, who required 981 quarantine days. Of the 700 EMS providers, 3 (0.4%) tested positive within 14 days of encounter, though these positive tests were

not attributed to occupational exposure from inadequate PPE. Programmatic changes were associated with a temporal reduction in exposures. When stratified at the study encounters midpoint, 94% (142/151) of exposures occurred during the first 137 EMS encounters compared with 6% (9/151) during the second 137 EMS encounters ($p < 0.01$). By the investigation's final week, EMS deployed MEGG PPE in 34% (3579/10 468) of all EMS person-encounters. Less than 0.5% of EMS providers experienced COVID-19 illness within 14 days of occupational encounter. Programmatic strategies were associated with a reduction in exposures, while achieving a measured use of PPE.

INTRODUCTION

The first case of COVID-19 in King County, Washington, USA, was reported on 28 February 2020. Incidence rose exponentially in subsequent weeks.¹ Emergency medical services (EMS) are the front line of the healthcare system, responding with incomplete information to provide care in heterogeneous, often uncontrolled, circumstances. The COVID-19 pandemic challenges healthcare worker (HCW) safety in part because of limited supplies of personal protective equipment (PPE). Ideally, EMS strategies would incorporate COVID-19 risk assessment and target use of the limited PPE resource in order to achieve EMS provider safety, extend the supply of PPE and support high-quality patient care.

The US Centers for Disease Control and Prevention (CDC) established criteria for COVID-19 testing and case management based on history and recent travel to a high-risk area, contact with known or suspected COVID-19 cases and presence of fever and signs/symptoms of lower respiratory illness.² Based on national guidelines, our regional EMS system initially adopted

a screening framework based on travel, exposure to known cases and specific symptoms. During the initial days and weeks of the outbreak, we identified long-term care facilities (LTCFs) as high-risk locales and observed the atypical presentations involving COVID-19 illness.^{3–5} As a consequence, we implemented a series of iterative protocol changes with regard to COVID-19 risk assessment and PPE use based on the patient's clinical profile and response location.

We evaluated all 9-1-1 EMS responses to patients with COVID-19 to (1) determine occupational exposure, related workforce quarantine and potential transmission, and (2) understand how programmatic changes influenced occupational exposure, workforce quarantine and PPE use amidst the COVID-19 outbreak in Seattle and King County.

METHODS

Study design and setting

The study is a retrospective cohort investigation of EMS providers responding to 9-1-1 calls for laboratory-confirmed COVID-19-positive patients in King County, Washington, USA between 14 February 2020 and 26 March 2020. The first US case was documented in neighbouring Snohomish County on 20 January, with unrecognised transmission of COVID-19 until clinical diagnosis within King County in late February 2020.^{6,7} EMS providers who cared for patients with COVID-19 were monitored through 9 April 2020 to complete a 14-day surveillance after the final patient encounter date. During this time, COVID-19 disease was defined by the State of Washington as positive reverse transcriptase-PCR (RT-PCR) testing for SARS-CoV-2.

King County is a metropolitan region, covering 2132 square miles, with 2.2 million persons who reside in urban, suburban and rural areas. The primary 9-1-1 medical response in King County is two-tiered. The first tier is provided by firefighter emergency medical technicians. Paramedics comprise the second tier and are dispatched in cases of more severe illness. There are 28 first-tier fire departments and five overarching second-tier paramedic agencies that collectively provide primary emergency response to all 9-1-1 medical calls. In general, stable patients are transported via fire department or private ambulance basic life support units, and more acute patients are transported by advanced life support paramedic units. All EMS, fire and private

¹Emergency Medicine, University of Washington, Seattle, Washington, USA

²Public Health Seattle and King County, Emergency Medical Services Division, Seattle, Washington, USA

³American Medical Response Puget Sound, Seattle, Washington, USA

⁴Tri-Med Ambulance, Kent, Washington, USA

⁵Redmond Fire Department, Redmond, Washington, USA

⁶Kirkland Fire Department, Kirkland, Washington, USA

⁷Public Health Seattle and King County, Seattle, Washington, USA

⁸Department of Medicine, University of Washington, Seattle, Washington, USA

⁹Seattle Fire Department, Seattle, Washington, USA

Correspondence to Dr David L Murphy, Emergency Medicine, University of Washington, Seattle, WA 98104, USA; dlmurphy@u.washington.edu

ambulance agencies in King County participated in this study. Collectively, there are approximately 4000 EMS providers in King County.

Study population

Case identification and COVID-19 encounters

The study population consisted of EMS providers who cared for patients with confirmed COVID-19 by RT-PCR tests. EMS is administered by Public Health–Seattle and King County, enabling direct engagement between EMS and Public Health to undertake COVID-19 surveillance. To identify EMS encounters with patients with COVID-19, we linked local and state COVID-19 surveillance systems with EMS electronic records using the patient's name and date of birth. Patient encounters were included if they occurred within a transmission window of 3 days prior to symptom onset (if known) or 14 days prior to or after the diagnosis date. The median interval from EMS encounter to diagnosis date was 4 days (IQR 2–6). Each match was independently verified by an epidemiologist and physician.

Case review process

A physician reviewed each matched encounter for potential EMS exposure in the electronic health record. If the documented PPE was not a complete ensemble of appropriate mask, eye protection, gown and gloves (MEGG), the case was further investigated by the EMS agency's appointed health officer (figure 1). Health officers contacted individuals with possible exposure to understand the specific circumstances of patient involvement and clarify PPE use. The health officer in consultation with physician leadership then made the final determination of exposure and whether quarantine or isolation was indicated according to the CDC risk assessment matrix.⁸

Definitions of encounter and occupational exposure

An encounter was defined as a 9-1-1 EMS response to a patient confirmed to have COVID-19. An occupational exposure to COVID-19 was defined as a provider-level encounter with inadequate PPE for the patient contact.⁸ In addition to eye

protection and gloves, a surgical mask was judged to be sufficient for routine patient encounters. However, an N95 mask was required PPE for aerosol-generating procedures. For any physical contact with the patient, a gown was required.

Employee screening, quarantine/isolation and testing

By the second week of March, most EMS agencies had implemented regular employee symptom screening on arrival at work and during the shift. Anyone who felt unwell for any reason returned home until they were asymptomatic and fit for duty per their agency return to work guidelines. EMS providers who became ill regardless of exposure status were deemed symptomatic, placed on isolation and prioritised for COVID-19 RT-PCR testing through dedicated first responder testing sites. These RT-PCR tests were performed by the University of Washington Virology Laboratory using an assay shown to have a low false negative rate.⁹ Each EMS agency assessed quarantined providers daily. The current investigation used information from both the health officer monitoring programme and the Public Health surveillance to ascertain any COVID-19 tests performed among the EMS provider cohort.

Interventions

Initial high-risk criteria

Prior to the first laboratory-confirmed case of COVID-19 in King County on 28 February 2020, EMS medical direction issued directives for COVID-19 screening and patient care on 6 February and 27 February 2020. Beginning 4 March, EMS providers were advised to don full MEGG PPE if COVID-19 screening included (1) a person with febrile respiratory illness AND travel from an endemic area (initially Wuhan, then broadened to China, South Korea, Iran or Italy) OR (2) febrile respiratory illness AND known contact with a patient with confirmed COVID-19.

Dispatch PPE advised

After 28 February, EMS updated the high-risk criteria to include the first LTCF where initial cases were identified, with dispatch to alert 'PPE advised' for any response to the address. After additional cases were identified at a second LTCF and a dialysis centre, these sites were added as high-risk locations for dispatch. A growing list of LTCFs and congregate living centres soon followed. Beginning 7 March, EMS began to treat all LTCFs (skilled nursing facilities, assisted living

Figure 1 Flow diagram. EMS, emergency medical services; PPE, personal protective equipment.

facilities and adult family homes) as high-risk requiring full MEGG PPE, regardless of clinical illness profile.

Clinical criteria profile

With evidence of community transmission, the requirements for travel history or COVID-19 contact were eliminated as criteria to don MEGG PPE during the first week of March. Medical record review determined that EMS COVID-19 patients did not consistently demonstrate a febrile respiratory illness; criteria were expanded to include any respiratory or fever symptoms beginning 11 March.⁵

Scout programme

Case review indicated that initial symptom classification—often derived from dispatch reporting—did not adequately characterise illness and the potential for COVID-19 illness. In response, EMS was using large quantities of PPE to address this uncertainty, though the prevalence of confirmed COVID-19 EMS encounters was estimated to be less than 5%.¹ Hence, EMS leadership implemented a 'scout programme' beginning 14 March in which one or two EMS providers donned full MEGG PPE and entered the 'hot zone' to perform the initial in-person evaluation while additional crew remained in the 'cold zone', maintaining sight or voice contact, with scout responder(s). The scout evaluation informed the need for remaining EMS crew to don PPE to assist. Conversely, risk assessment was often not feasible in high-acuity, time-sensitive cases. All cardiac arrest cases and cases requiring aerosol-generating therapies required full MEGG PPE with N95 masks.

Data collection and measurements

We used a uniform methodology to review the narrative and formatted data fields from dispatch and EMS records. Dispatch records were abstracted to characterise 9-1-1 patient concern and pre-arrival notifications. EMS records were abstracted to describe patient characteristics, location, initial vital signs, disposition, clinician impression and PPE use. PPE use was assessed through review of the EMS report narrative and discrete data fields. Following the first recognised case of COVID-19 in King County, the EMS leadership directed reporting of full PPE use in the electronic record by responding EMS personnel. Beginning 20 March, mandatory, item-specific PPE reporting became available through the electronic health record (ESO Solutions, Austin, Texas, USA) for all EMS responses. EMS

provider quarantine dates and results from COVID-19 testing were recorded.

Outcomes

We evaluated the number of patient with COVID-19 encounters, PPE use, consequent exposures due to inadequate PPE, resulting quarantine and positive COVID-19 tests among EMS providers.

Analysis

Descriptive analyses were performed at the EMS encounter and EMS provider levels. EMS encounters were stratified by level of transport, while provider-level assessments were stratified at the chronologic midpoint of EMS encounters. Due to a subset of providers with multiple patient encounters, we report provider-level assessments as both total EMS provider encounters and unique EMS providers. We used χ^2 test for trend to evaluate whether adequate PPE use and EMS provider exposure changed over time, where calendar time was the independent variable and EMS provider exposure (or adequate PPE use) was the dependent variable. We used a χ^2 test to compare the proportion of encounters with occupational exposures in the first and second half of EMS encounters. To estimate the potential conservation of PPE relative to an indiscriminate MEGG PPE deployment strategy (MEGG for all EMS personnel for all calls), we determined the actual PPE use during the week of 20–26 March among the total number of EMS providers involved on 9-1-1 responses. SAS (V9.4; SAS Institute) was used to conduct analyses.

RESULTS

Characteristics of patients with COVID-19

There were 220 unique patients with confirmed COVID-19 in Seattle and King County with 9-1-1 EMS encounters in the 14 days prior to, and first 28 days after, the sentinel laboratory-confirmed case in King County. Of these 220 individuals, 54 had two EMS encounters for a total of 274 distinct EMS encounters. Half were female (53%), and the mean age was 74 years. The dispatch complaints were heterogeneous; difficulty breathing was the most common complaint, accounting for about 25% (table 1). The median initial pulse oximetry reading was 93%. The most common EMS impressions included suspected COVID-19 illness (26%), flu-like symptoms (17%), respiratory distress (17%) and weakness (14%).

Main results

Among the 274 EMS encounters with patients with COVID-19, there were 429 responding units, involving 700 unique EMS providers with a total of 988 EMS provider encounters (table 2). Based on initial EMS record review, use of PPE during patient contact was full MEGG (66.9%), basic gloves and eye protection (29.3%), delayed application or partial MEGG (3.1%), or unknown (0.7%), resulting in 327 possible EMS provider exposures. After health officer investigation and physician consultation, 151 EMS provider encounters were determined to have an exposure. As a result, there were 129 unique EMS providers placed on quarantine: 107 after a single exposure and 22 with two exposures. Of the 700 unique EMS providers caring for patients with confirmed COVID-19, 3 (0.4%) tested positive during the 14 days following an encounter (table 3), yet none of these three had a documented occupational exposure.

The series of practice changes involving dispatch advisement, patient COVID-19 risk criteria and initial EMS scene deployment were associated with a temporal increase in adequate PPE use and conversely a decrease in EMS provider exposures (figure 2, $p < 0.01$). When stratified at the encounters midpoint, 94% (142/151) of exposures occurred during the first 137 EMS encounters compared with 6% (9/151) during the second 137 EMS encounters (table 2, $p < 0.01$). The number of EMS providers quarantined each day increased to a peak of 69 on 13 March and then declined (figure 3).

During the final week of the study (20–26 March), there were a total of 3704 EMS incidents involving 10468 EMS providers. Of the 10468 opportunities for PPE deployment, MEGG PPE was used in 3579 (34%) EMS provider encounters.

DISCUSSION

In this population-based observational investigation of 274 EMS encounters for patients with COVID-19 involving nearly 1000 EMS provider encounters, three EMS providers subsequently tested positive for COVID-19 during the 14 days following the patient encounter. Iterative dispatch and operational EMS responses to COVID-19 risk identification and PPE use were associated with both a temporal decrease in EMS provider COVID-19 exposure and conservation of PPE. Based on these programmatic efforts, full MEGG PPE was deployed in about one-third of all

Report from the front

Table 1 Characteristics of 274 EMS encounters with 220 patients with confirmed COVID-19 from 14 February to 26 March 2020

	Total EMS encounters (n=274)	BLS transport (n=180)	ALS transport (n=31)	Not transported (n=63)
Initial dispatch complaint, n (%) of column				
Difficulty breathing	68 (25)	44 (24)	10 (32)	14 (22)
Sick (unknown)	68 (25)	52 (29)	6 (19)	10 (16)
Infectious diseases	51 (19)	35 (19)	4 (13)	12 (19)
Trauma	32 (12)	17 (9)	2 (7)	13 (21)
Other*	55 (20)	32 (18)	9 (29)	14 (22)
Dispatched as 'PPE advised', n (%) of column	196 (72)	136 (76)	16 (52)	44 (70)
Origin of 911 response, n (%) of column				
Home/private residence	128 (47)	76 (42)	16 (51)	36 (57)
Long-term care facility	118 (43)	86 (48)	12 (39)	20 (32)
Outpatient clinic	20 (7)	14 (8)	3 (10)	3 (5)
Public/street	8 (3)	4 (2)	0 (0)	4 (6)
Initial vital signs, median (IQR) of column				
Systolic blood pressure, mm Hg (n=238)	130 (110–146)	130 (110–142)	132 (97–155)	130 (110–140)
Heart rate, beats per minute (n=230)	90 (80–108)	92 (80–110)	97 (72–115)	88 (76–96)
Respiratory rate, breaths per minute (n=238)	21 (16–24)	20 (16–24)	24 (16–30)	18 (16–20)
Peripheral oxygen saturation, % (n=197)	93 (89–96)	92 (89–95)	92 (83–96)	95 (92–97)
Temperature, Celsius (n=172)	37.7 (36.9–38.8)	37.8 (37.1–38.8)	37.5 (36.4–38.2)	37.3 (36.7–38.3)
EMS provider impression, n (%) of column				
COVID†	72 (26)	53 (29)	4 (13)	15 (24)
Flu-like symptoms	47 (17)	37 (21)	4 (13)	6 (10)
Respiratory distress	46 (17)	27 (15)	11 (36)	8 (13)
Weakness	37 (14)	25 (14)	0 (0)	12 (19)
Altered mental status	16 (6)	8 (4)	4 (13)	4 (6)
Injury/pain	16 (6)	13 (7)	1 (3)	2 (3)
Cardiac	15 (6)	6 (3)	6 (19)	3 (5)
Other‡	25 (9)	11 (6)	1 (3)	13 (21)
Any mention of COVID-19 in EMS record, n (%) of column	169 (62)	117 (65)	17 (55)	35 (56)

*Bleeding/pain, cardiac and stroke/headache.

†COVID-19 impressions were added to the electronic health record as an option on 8 March 2020.

‡Vaginal haemorrhage, seizures, obvious death, gastrointestinal haemorrhage, epistaxis, dehydration, urinary tract infection, diabetic hypoglycaemic, unspecified convulsions and skin infection.

ALS, advanced life support; BLS, basic life support; EMS, emergency medical services.

potential EMS provider uses by the end of the study period.

Although HCWs seem to be at higher risk to contract COVID-19, rigorous assessment of exposure and transmission is largely lacking. Epidemiological reports from China and Italy highlight the

substantial burden of illness in HCWs.^{10–12}

Locally, in Washington State, a large portion of LTCF staff tested positive for COVID-19.³ A preliminary report from CDC regarding the burden of COVID-19 infection among US healthcare personnel suggest HCWs account for 11%–19% of

national case burden, but did not discern specific type of employment or evaluate the potential source of exposure.¹³ Other reports involving high-risk circumstances to include aerosolising procedures however have not observed substantial rates of transmission to HCWs.¹⁴

Table 2 Use of PPE and occupational exposures among EMS provider encounters with patients with confirmed COVID-19 from 14 February to 26 March 2020

	Total	*Initial 137 EMS encounters with COVID-19 patients	*Subsequent 137 EMS encounters with COVID-19 patients
Number of EMS provider encounters, n	988	488	500
Any mention of COVID-19 in EMS record, n (%)	133 (49)	49 (36)	84 (61)
PPE, n (%) of column total			
Full MEGG	661 (67)	265 (54)	396 (79)
Partial or delayed MEGG†	31 (3)	17 (3)	14 (3)
Basic (gloves/eyes)	289 (29)	202 (41)	87 (17)
Missing/unknown*	7 (1)	4 (1)	3 (1)
EMS provider encounters with an exposure, n (%) of column total	151 (15)	142 (29)	9 (2)
EMS provider exposures per 9-1-1 encounter with an exposure, median (IQR)	3 (2–3)	3 (2–3)	2 (2–3)
Interval from EMS provider exposure to patient COVID-19 test result, days, median (IQR)	4 (2–6)	5 (2–8)	3 (1–4)

*Initial encounters occurred between 19 February and 15 March 2020 while subsequent encounters occurred between 16 March and 26 March 2020.

†Partial MEGG is defined as basic PPE (eye protection and gloves) plus either a mask or a gown but not both.

EMS, emergency medical services; MEGG, mask, eye protection, gown and gloves; PPE, personal protective equipment.

Table 3 Occupational exposures, quarantine and testing of unique EMS providers with patients with confirmed COVID-19 contact from 14 February to 26 March 2020

	Total	*Initial 137 EMS encounters with COVID-19 patients	*Subsequent 137 EMS encounters with COVID-19 patients
Unique EMS providers†	700	341	382
Unique EMS providers with patient exposure(s), n (% of column total)	129 (18)	121 (35)	8 (2)
Number of exposure(s) for each unique EMS provider, n (% of column total n=700)			
0	571 (82)	220 (65)	374 (98)
1	110 (16)	103 (30)	7 (2)
2	16 (2)	15 (4)	1 (<1)
≥3	3 (<1)	3 (1)	0 (0)
Interval from exposure to EMS provider quarantine, days, median (IQR)	6 (3–9)	6 (3–9)	3 (2–3)
Total EMS provider quarantine days resulting from exposure(s), n (% of row)	981 (100)	951 (97)	30 (3)
COVID-19 testing of unique EMS providers regardless of exposure status, n (% of column)			
No symptoms reported (not tested)	657 (94)	312 (91)	368 (96)
Symptoms reported	43 (6)	29 (9)	14 (4)
Positive	3 (<1)	1 (<1)	2 (1)
Negative	40 (6)	28 (8)	12 (3)

*Initial encounters occurred between 19 February and 15 March 2020 while subsequent encounters occurred between 16 March and 26 March 2020.

†23 providers were represented in both categories.

EMS, emergency medical services.

Similar to our findings, a Taiwanese study reported a transmission rate of 0.9% among the subset of COVID-19 exposures occurring in the healthcare setting.¹⁵ None of these experiences have reported risk to EMS providers, though EMS care appears to be integral for sicker COVID-19 patients. In the 2009 SARS outbreak, the overall incidence of infection was 1.3% in the Taiwanese EMS workforce, which was >100-fold higher than the general public.¹⁶

In the current investigation, EMS had substantial involvement with COVID-19 illness. The 220 patients represented 14% of all COVID-19 diagnoses in King County, Washington, through 26 March. EMS was typically involved in care for older adults who often presented with heterogeneous

symptoms and a range of clinical presentations. COVID-19 in King County was first detected in a clinical population not considered high-risk according to national guidelines at that time, which accounted in part for the fact that 18% of EMS providers in the study had an exposure. Indeed, 85.4% of patients had not been diagnosed with COVID-19 at the time of their EMS encounter.

The high rate of quarantine early on motivated the EMS system to move quickly to adapt to the evolving clinical features and local epidemiology of the COVID-19 outbreak. EMS leadership engaged dispatch and operations to expand COVID-19 risk criteria and to stage patient assessment. The set of measures was associated with a marked

reduction in the risk of exposure over the course of investigation. Certainly, there was a learning curve that may have also contributed to reduction in exposure. The collective effect appears to be a temporal reduction in EMS worker quarantine, even though the number of provider encounters with COVID-19 increased over time (figure 2).

We observed that 3 of the 700 EMS providers (0.4%) with COVID-19 encounters subsequently tested positive for COVID-19. One case occurred at the outset of the outbreak with onset of provider illness occurring on the same date of COVID-19 encounter. The CDC investigated this case and determined that the 9-1-1 incident that qualified the provider for study inclusion was not responsible

Figure 2 Occupational exposures and PPE use among EMS providers caring for patients with COVID-19, Seattle and King County through 26 March 2020. EMS, emergency medical services; PPE, personal protective equipment.

Figure 3 Number of Seattle and King County emergency medical services providers in quarantine by calendar day.

for disease transmission. Nonetheless, the provider may have had a patient exposure in the days prior to identification of COVID-19 cases in King County, as review of prior encounters involving the provider confirmed care for patients with acute respiratory illness. The providers in all three cases had MEGG PPE during their qualifying encounters. We cannot determine whether transmission occurred during these patient-specific exposures, other occupational activities or community transmission.

Overall, the cumulative laboratory-confirmed prevalence in this EMS cohort of 700 unique providers (0.4%) is comparable with the community prevalence (0.2%) during this time frame.¹ Taken together, these findings suggest that occupational risk can be relatively low and that protective measures can potentially limit disease transmission. The anecdotal experiences in other regions reporting high rates of COVID-19 among EMS providers may be related to the higher prevalence of disease paired with limited availability and use of PPE.

There is an inherent tension between proactive measures to don adequate PPE and conservation efforts due to limited supplies. If PPE were limitless, then indiscriminate use by all providers for every call would help assure EMS provider protection. However, our system had limited supply that was coupled with uncertainty about the severity and duration of the pandemic. Thus, the EMS system strived to target the use of PPE to risk-positive patients. The scout strategy for stable patients enabled more deliberate decisions regarding PPE. In contrast, time-critical events such as cardiac arrest required comprehensive EMS PPE, given the need

for care prior to evaluating COVID-19 risk. The current targeted strategies for MEGG utilisation appear to be a viable means to protect EMS providers and conserve PPE.

The retrospective methodology used to assess PPE is imperfect, relying on documentation and case-specific investigation; the two-stage process however enabled detailed provider interviews to assess potential exposure. The initial stage of screening mandated investigation anytime there was no clear documentation of full PPE in a patient with COVID-19. In the second stage health officer review, EMS providers sometimes clarified that full PPE was in fact in place though not adequately documented in the report. In other instances, individual EMS providers without MEGG PPE were not in proximity of the patient (ie, the scout method that deployed only a subset of the crew for direct patient contact). We acknowledge that provider documentation may introduce bias, although providers were motivated to accurately document PPE. Providers received training and education on best practices of donning and doffing of PPE, but there was not a dedicated observer to document the quality of the process. The study could not report on the temporal use of PPE across the system, but rather the status after implementation of various interventions designed to better assess COVID-19 risk and responsibly use PPE. Ideally, the study would have tracked PPE use across the system from the outset of the COVID-19 pandemic to better understand how programmatic changes influenced PPE deployment. Documentation of quarantine evolved during the study period to use a central monitoring database. Thus, quarantine decisions early

in the outbreak may be an underestimate of quarantine.

We relied on the statewide Washington Disease Reporting System database to identify COVID-19 positive patients. There likely were patients ill with COVID-19 who interfaced with EMS but were not tested. Alternatively, EMS encounters with COVID-19 positive patients may exist that were not captured due to failed linking of identifiers between EMS and surveillance databases. The study relied on EMS agency health officers and the Washington Disease Reporting System database to identify EMS providers tested for COVID-19. Although unlikely, this dual approach may have missed a laboratory-confirmed infection in an EMS provider. EMS providers may also have chosen not to get tested or had asymptomatic infection, though symptomatic providers were motivated to be tested and had prioritised access to testing. We cannot confirm the source of the infectious exposure—patient-specific, other occupational or community transmission—among the few providers with positive tests.

In conclusion, less than 0.5% of EMS providers experienced COVID-19 illness within 14 days of caring for a patient with laboratory-confirmed COVID-19. Programmatic risk mitigation strategies were associated with a reduction in occupational exposures to COVID-19 among EMS providers, while achieving a measured use of PPE.

Acknowledgements We wish to acknowledge Public Health—Seattle and King County, the Washington State Department of Health, the Centers for Disease Control and the telecommunicators and EMS professionals of Seattle and greater King County.

Contributors DLM, BYY, MRS and TDR conceived the study and designed the investigation. LMB, CJD, CRC, JME, LS, TYJ, AMM and TAM supervised data collection. LMB, CJD, CRC and JME managed the data, including quality control. LMB, JME and TDR provided statistical advice on study design and analysed the data. DLM drafted the manuscript, and all authors contributed substantially to its revision. DLM takes responsibility for the paper as a whole.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not required.

Ethics approval The study was approved by the University of Washington Institutional Review Board.

Provenance and peer review Not commissioned; externally peer reviewed.

OPEN ACCESS

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

© Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

Handling editor Roland Merchant

Check for updates

To cite Murphy DL, Barnard LM, Drucker CJ, *et al.* *Emerg Med J* 2020;**37**:707–713.

Received 23 May 2020

Revised 10 July 2020

Accepted 5 August 2020

Published Online First 21 September 2020

Emerg Med J 2020;**37**:707–713.

doi:10.1136/emered-2020-210095

ORCID iD

David L Murphy <http://orcid.org/0000-0002-6237-4150>

REFERENCES

- Public Health-Seattle & King County. COVID-19 data dashboard. Available: <https://kingcounty.gov/depts/health/communicable-diseases/disease-control/novel-coronavirus/data-dashboard.aspx>
- Centers for Disease Control and Prevention. Update and interim guidance on outbreak of coronavirus disease 2019 (COVID-19), 2020. Available: <https://emergency.cdc.gov/han/2020/han00428.asp>
- Guan W-J, Ni Z-Y, Hu Y, *et al.* Clinical characteristics of coronavirus disease 2019 in China. *N Engl J Med* 2020;382:1708–20.
- McMichael TM, Currie DW, Clark S, *et al.* Epidemiology of Covid-19 in a long-term care facility in King County, Washington. *N Engl J Med* 2020;382:2005–11.
- Yang BY, Barnard LM, Emert JM, *et al.* Clinical characteristics of patients with coronavirus disease 2019 (COVID-19) receiving emergency medical services in King County, Washington. *JAMA Netw Open* 2020;3:e2014549.
- Bedford T, Greninger AL, Roychoudhury P, *et al.* Cryptic transmission of SARS-CoV-2 in Washington state. *medRxiv* 2020. doi:10.1101/2020.04.02.20051417. [Epub ahead of print: 06 Apr 2020].
- Public Health-Seattle & King County. First death due to novel coronavirus (COVID-19) in a resident of King County, 2020. Available: <https://kingcounty.gov/depts/health/news/2020/February/29-covid19.aspx>
- Centers for Disease Control and Prevention. Interim U.S. guidance for risk assessment and public health management of healthcare personnel with potential exposure in a healthcare setting to patients with coronavirus disease (COVID-19), 2020. Available: <https://www.cdc.gov/coronavirus/2019-ncov/hcp/guidance-risk-assessment-hcp.html>
- Long DR, Gombor S, Hogan CA, *et al.* Occurrence and timing of subsequent SARS-CoV-2 RT-PCR positivity among initially negative patients. *medRxiv* 2020.
- Wang D, Hu B, Hu C, *et al.* Clinical characteristics of 138 hospitalized patients with 2019 novel Coronavirus–Infected pneumonia in Wuhan, China. *JAMA* 2020;323:1061–9.
- Wu Z, McGoogan JM. Characteristics of and Important Lessons From the Coronavirus Disease 2019 (COVID-19) Outbreak in China: Summary of a Report of 72 314 Cases From the Chinese Center for Disease Control and Prevention. *JAMA* 2020;323:1239–42.
- Onder G, Rezza G, Brusaferro S. Case-Fatality rate and characteristics of patients dying in relation to COVID-19 in Italy. *JAMA* 2020;323:1775–6.
- CDC COVID-19 Response Team. Characteristics of Health Care Personnel with COVID-19 - United States, February 12–April 9, 2020. *MMWR Morb Mortal Wkly Rep* 2020;69:477–81.
- Ng K, Poon BH, Kiat Puar TH, *et al.* COVID-19 and the risk to health care workers: a case report. *Ann Intern Med* 2020;172:766–7.
- Cheng H-Y, Jian S-W, Liu D-P, *et al.* Contact tracing assessment of COVID-19 transmission dynamics in Taiwan and risk at different exposure periods before and after symptom onset. *JAMA Intern Med* 2020:e202020.
- Ko PC-I, Chen W-J, Ma MH-M, *et al.* Emergency medical services utilization during an outbreak of severe acute respiratory syndrome (SARS) and the incidence of SARS-associated coronavirus infection among emergency medical technicians. *Acad Emerg Med* 2004;11:903–11.