indication. Liver bloods, clotting screen, bone profile tests were performed on 40%, 24% and 27% respectively in the second cycle.

The utility of blood tests may be an under recognised subject in paediatric training which needs addressing. This project typified positive impact of culture change via QI champions and educational sessions. Implementation of such changes is sustainable with an estimated savings of at least £500/month.

451

CHILDREN'S EMERGENCY DEPARTMENT REVIEW CLINIC: WHAT WE HAVE LEARNT IN THE COVID-19 PANDEMIC

Charlotte Munday, Neha Jain, Seamus Harrington, Benjamin Darvill, Sophie Perry, Alice Irwin, Francesca Deibel, Richard Turck, Clarissa Barber. *Bristol Royal Hospital for Children*

10.1136/emj-2020-rcemabstracts.54

Aims/Objectives/Background The Children's Emergency Department (CED) in the Bristol Royal Hospital for Children (BRHC) runs 'Review Clinics' by a Paediatric Emergency Consultant. These were suspended in March 2020 due to the Covid-19 pandemic and new patient pathways were put in place. Our aim was to assess the effectiveness of these pathways and to evaluate if they should continue once the pressures of Covid-19 have eased.

Methods/Design We reviewed all the attendances to the ED Review Clinic over a six-week period in 2019 and identified common conditions that are seen in this clinic – abscesses, cellulitis, scaphoid fracture, toddler fracture, limp, headache, peri-orbital cellulitis, torticollis, and neck lumps (accounting for 66% of all attendances).

We then identified all of the ED attendances with these conditions over the same six-week period during the Covid-19 pandemic and performed a service review using medical notes. **Results/Conclusions** All children presenting with a limp during the Covid-19 pandemic were discharged with a leaflet and advised to return in one week if the limp persisted. None of them re-attended.

Scaphoid and toddler fractures that presented during the Covid-19 pandemic were followed-up by Orthopaedics with a combination of telephone and face-to-face reviews. The toddler fractures were put into soft casts to be removed at home - 93% of parents were happy to do this. Orthopaedic review was quicker than the previous ED Review Clinic (an average of 63 hours compared to 216 for scaphoid fractures and 67 hours compared to 235 for toddler fractures).

The results suggest that ED Review Clinics can and should change. Children with a limp can be sent home safely with clear advice. There is a greater role for telephone reviews, in particular, for injuries such as toddler fractures. This would be quicker, less time-consuming for patients, parents and physicians and more cost effective for the department.

468

PRESENTATION OF VULNERABLE CHILDREN TO ED DURING LOCKDOWN: A CONCERNING CONSEQUENCE OF ISOLATION

Lalarukh Asim, Raeesa Jina, Sreena Das. *King's College Hospital*

10.1136/emj-2020-rcemabstracts.55

Aims/Objectives/Background This study aims to evaluate changes to child safeguarding attendances to the emergency department (ED) during the lockdown period 1st of March till 31st May 2020 compared to the same time period in 2019.

The COVID-19 pandemic is the defining health crisis of our time and the first time firm social restrictions have been imposed since safeguarding practices have become embedded in the NHS. The NSPCC reported a large increase in contacts but there remain concerns that vulnerable children are invisible to agencies during this time.

Methods/Design All children (< 18 years) who attended King's College Hospital's ED, and were reviewed in the weekly ED safeguarding meeting were included. Data was collected from electronic patient records for different parameters.

Results/Conclusions The total number of children presenting via ED for safeguarding review fell from 865 in 2019 to 355 for the same period in 2020. However, the proportion requiring action by the hospital safeguarding team showed a significant increase ($p = 1.5 \times 10^{-4}$) suggesting the severity of cases during COVID-19 is worse.

The percentage of stabbings doubled ($p = 0.04$) despite lockdown measures. This may be a contributor to the significant increase in referral to youth workers ($p = 8 \times 10^{-4}$). The number of children attending who were considered high risk due to previous safeguarding concerns dropped by 75%.

As expected, the proportion of household injuries such as accidental ingestion and burns showed a significant increase ($p = 0.01$ and $p = 0.002$ respectively). The proportion of children from outside of our local boroughs was surprisingly higher in 2020 ($p = 0.03$).

The findings show that the number of cases triggering a safeguarding review has dropped during lockdown and raises concerns about vulnerable children who remain hidden. The findings also suggest an increased severity of safeguarding presentations, supporting our fears that the implications of lockdown on vulnerable children is yet to be realised.

332

THE DIAGNOSTIC ROLE OF SHUNT SERIES RADIOGRAPHS (SSR) IN CHILDREN PRESENTING TO THE CHILDREN'S EMERGENCY DEPARTMENT (CED) WITH SUSPECTED VENTRICULOPERITONEAL (VP) SHUNT FAILURE

¹Angharad Griffiths, ¹Ikechukwu Okafor, ²Thomas Beattie. ¹Children's Health Ireland at Temple Street; ²University of Edinburgh

10.1136/emj-2020-rcemabstracts.56

Aims/Objectives/Background VP shunts are used to drain CSF from the cranial vault because of a wide range of pathologies and, like any piece of hardware, can fail. Traditionally investigations include SSR and CT. This project examines the role of SSR in evaluating children with suspected VP shunt failure.

Primary outcome: Sensitivity and specificity of SSR in children presenting to the CED with concern for shunt failure.

Methods/Design Conducted in a single centre, tertiary CED of the national Irish Neurosurgical(NS) referral centre (ED attendance: >50,000 patients/year). 100 sequential SSR requested by the CED were reviewed. Clinical information was extracted from electronic requests. Shunt failure was defined by the need for NS intervention(Revision).

Abstract 332 Figure 2

	Shunt malfunction	No shunt malfunction
Abnormal SS	6	1
Normal SS	16	77

Sensitivity = 0.27 (27%)

Specificity = 0.99 (99%)

PPV = 0.86 (86%)

NPV = 0.83 (83%)

Abstract 332 Figure 1

Results/Conclusions

- Sensitivity and specificity is presented in figure 1 (two by two table).
- 100 radiographs performed in 84 children.
- 22% shunts revised (see flow diagram).
- 7 SSR's were abnormal.
 - 85% (n=6) shunts revised. [5 following abnormal CT].
- Of the normal SSR's; 16 had abnormal CT and revised.
- 85/100 received CT.
 - 64 of 85 CT's (75%) were normal.
 - 6 of the 64 had focal shunt concern.
- SSR's shouldn't be used in isolation. NPV&PPV, Sensitivity&Specificity is low.
- SSR's are beneficial where there's concern over focal shunt problems (injury/pain/swelling) or following abnormal CT.

- VP shunt failure is not well investigated with SSR alone.
- SSR's could be omitted where there is no focal shunt concern/ after normal CT (without impacting clinical outcome) reducing radiation exposure and reduce impact on CED's.

59 SSR's *could* have been avoided without adverse clinical outcome.

31

HEAD HOME: A PROSPECTIVE COHORT STUDY OF A NURSE LED PAEDIATRIC HEAD INJURY CLINICAL DECISION TOOL AT A DISTRICT GENERAL HOSPITAL

¹Patrick Aldridge, ¹Heather Castle, ¹Emma Russell, ¹Clare Phillips, ²Richard Guerrero-Ludueno, ³Raj Rout. ¹Frimley Park; ²University of Southampton, School of Health Sciences, Centre for Implementation Science, UK; ³Sanofi Genzyme

10.1136/emj-2020-rcemabstracts.57

Aims/Objectives/Background

Objectives To assess if application of a nurse-led paediatric head injury clinical decision tool would be safe compared to current practice.

Background >700,000 children attend UK hospitals' each year with a head injury. Research indicates <1% undergo neurosurgical intervention. No published evidence for nurse-led discharge of paediatric head injuries exists.

Methods/Design Methods – All paediatric (<17 years) patients with head injuries presenting to our Emergency department (ED) 1st May to 31st October 2018 were prospectively screened by a nurse using a mandated electronic 'Head Injury Discharge At Triage' questionnaire (HIDATq). We determined which patients underwent computed tomography (CT) brain and whether there was a clinically important intracranial injury or re-presentation to ED. The negative predictive value of the screening tool was assessed. We determined what