

**Supplementary table 2.** Reported symptoms of patients diagnosed both COVID-19 positive and negative by PCR or CT scanning.

Reported symptoms	PCR or CT COVID-19 negative (n,%)	PCR or CT COVID-19 positive (n,%)	Total missing data
Malaise	14 (61%)	15 (83%)	52
Fatigue	14 (56%)	14 (70%)	48
Fever*	15 (34%)	18 (78%)	26
Headache	7 (27%)	7 (35%)	47
Rhinorrhoea	6 (21%)	1 (6%)	47
Sore throat	4 (14%)	2 (10%)	45
Coughing	33 (57%)	19 (73%)	9
Sputum production	9 (20%)	4 (17%)	25
Dyspnoea	32 (55%)	18 (67%)	8
Pain with breathing	6 (18%)	2 (11%)	40
Haemoptysis	2 (7%)	0	45
Orthopnea	1 (4%)	3 (16%)	46
Palpitations	6 (14%)	0	27
Chest pain*	20 (36%)	3 (12%)	11
Anorexia*	19 (56%)	19 (86%)	37
Nausea	9 (21%)	7 (29%)	26
Vomiting	11 (24%)	4 (18%)	26
Abdominal pain	10 (24%)	3 (13%)	29
Diarrhoea	11 (24%)	11 (46%)	23
Anosmia	0	6 (75%)	84
<b>Total</b>	<b>66 (71%)</b>	<b>27 (29%)</b>	

Percentages are calculated from the total within that group minus the missing data.

CT = computed tomography; n = number; PCR = polymerase chain reaction. \*p <0,05.