

Retrospective single-centre descriptive study of the characteristics, management and outcomes of adult patients with suspected sepsis in the emergency department

Lisa Sabir ¹, Laura Wharton,² Steve Goodacre ¹

Handling editor Ed Benjamin Graham Barnard

► Additional online supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/emmermed-2020-211111>).

¹School of Health and Related Research, The University of Sheffield, Sheffield, UK
²Academic Unit of Reproductive and Developmental Medicine, The University of Sheffield Jessop Hospital for Women, Sheffield, UK

Correspondence to

Dr Lisa Sabir, The University of Sheffield School of Health and Related Research, Sheffield, UK; l.sabir@sheffield.ac.uk

Received 28 December 2020
Accepted 16 July 2021
Published Online First 6 August 2021



► <http://dx.doi.org/10.1136/emmermed-2021-211627>



© Author(s) (or their employer(s)) 2022. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Sabir L, Wharton L, Goodacre S. *Emerg Med J* 2022;**39**:272–278.

ABSTRACT

Background Guidelines for adults presenting to the emergency department (ED) with suspected sepsis recommend protocols and bundles that promote rapid and potentially intensive treatment, but give little consideration of how patient characteristics, such as age, functional status and comorbidities, might influence management. This study aimed to describe the characteristics, management and outcomes of adults attending the ED with suspected sepsis, and specifically describe the prevalence of comorbidities, functional impairment and escalations of care.

Methods We undertook a single-centre retrospective observational study involving medical record review of a random sample of adults admitted to an ED between February 2018 and January 2019 with suspected sepsis. Descriptive statistics were used with 95% confidence intervals (CIs) for key proportions.

Results We included 509 patients (median age 74 years), of whom 49.3% met the Sepsis-3 criteria. Less than half of the patients were living at home independently (42.5%) or could walk independently (41.5%), 19.3% were care home residents and 89.2% of patients had one or more comorbidity. 22% had a pre-existing do not attempt resuscitation order. 6.5% were referred to intensive care, and 34.3% of the 13.2% who died in-hospital had an escalation plan explicitly documented.

Conclusion Adults with suspected sepsis have substantial functional limitations, comorbidities and treatment directives that should be considered in guidelines, especially recommendations for escalation of care.

INTRODUCTION

Sepsis is a life-threatening dysregulated response to infection that can lead to organ dysfunction, causing 52 000 deaths annually in the UK.¹ Diagnosing sepsis is difficult, as reflected in the evolution of its definition. The latest definition, ‘Sepsis-3’, combines the presence of infection with the Sequential Organ Failure Assessment (SOFA) score. The latter requires a change from baseline SOFA score of 2 or more to represent organ dysfunction, and the assumption that the baseline score is zero unless the patient is known to have pre-existing organ dysfunction before the onset of infection.¹ The quick SOFA (qSOFA) was developed for recognising those at a greater risk of poor

Key messages

What is already known on this subject

- Guidelines for suspected sepsis recommend rapid and potentially invasive treatment focused on saving lives.
- There has been limited research investigating when these treatments may not be appropriate or epidemiological studies describing the cohort of patients with suspected sepsis.
- Adults attending the emergency department often have functional limitations and comorbidities that, if reflected in people with suspected sepsis, may limit the potential benefit of intensive treatment.

What this study adds

- This retrospective, single-centre study has demonstrated that patients with suspected sepsis are typically elderly, less than half are living at home independently or can walk independently, almost one-fifth are care home residents, and few have no comorbidities.
- Guidelines for suspected sepsis should include these characteristics in management recommendations, especially treatment escalation decisions.

outcome outside of the intensive care unit (ICU) (see online supplemental material).^{1,2}

The number of patients presenting to emergency departments (ED) is increasing and we have an ageing population; between 2003 and 2015, the number of people aged over 85 years presenting to ED increased by nearly 40%.³ Multiple risk factors for infection exist in the elderly including immunosuppression, malnutrition, hospitalisation and medical procedures. Additionally, the population is increasingly comorbid and as people age they are more likely to live with long-term conditions (LTC) or frailty.⁴ 14% of those aged under 40 report having a LTC, increasing to 58% in the over 60s with 25% having two or more LTCs.⁵ These conditions influence outcomes in sepsis and sepsis also worsens their chronic disease.⁶

Clinical protocols aim to facilitate early recognition and treatment, most prominently the Surviving Sepsis Campaign (SSC) 1-hour bundle which describes the ‘Sepsis-6’: administration of



antibiotics, fluids and oxygen, and measurement of lactate, urine output and blood cultures in suspected sepsis.^{7,8} More invasive treatment options include vasopressors, mechanical ventilation and central lines; these will require ICU admission and should prompt discussions regarding escalation of care.⁷

Earlier disposition decisions, such as admission to ICU, result in lower hospital mortality.⁹ Despite this, guidelines contain limited recommendations about the effect of functional status and comorbidities on escalation decisions and this may fail to create realistic expectations about outcomes.¹⁰ The Surviving Sepsis Guidelines recommend that prognosis and care goals are discussed with patients early, and palliative care initiated if appropriate.⁷ However, these guidelines focus on ICU care. There is a drive for this decision making earlier to improve the quality of care by involving patients promptly in treatment decisions. The SARS-CoV-2 pandemic has brought these issues to the fore. The use of the Clinical Frailty Score has become standard practice and discussions regarding prognosis, escalation and patient wishes are occurring more frequently in the ED.¹¹

The National Institute for Health and Care Excellence (NICE) guidelines are evidence-based recommendations used in England. NICE sepsis research recommendations highlight the need for epidemiological studies to help plan services.¹² This is especially relevant as the population presenting to the ED changes. We aimed to describe the patient characteristics of suspected sepsis in the ED and factors influencing their escalation and treatment decisions.

METHODS

We conducted a retrospective observational study of adults presenting to a type 1 ED, the Northern General Hospital (NGH) in Sheffield, with suspected sepsis. It followed the Strengthening the Reporting of Observational Studies in Epidemiology guideline for cohort studies (online supplemental material).

Setting and study population

The NGH is Sheffield's only adult ED, with approximately 100 000 attendances annually. In 2019, Sheffield had a population of 584 853. Its median age was 35.4 (40.3 for the UK as a whole), there is a noticeably large proportion of 20–24 years old thought to be due to the student population at its two universities, it is an ethnically diverse city with approximately 19% of the population from black or minority ethnic groups.^{13,14}

All patients in whom blood cultures were performed in the ED between 01 January 2018 and 31 January 2019 were identified, as a screening method for suspected sepsis. The sample size was estimated (see below) and then cases were randomised using a computer-generated randomisation sequence and each entry was subsequently looked at in an ascending order of randomisation number for inclusion until the prespecified sample size was achieved.

Two clinicians (LS and LW) reviewed charts and determined whether the case was included based on whether the ED clinician suspected sepsis. This specifically required documentation of suspected sepsis, but also included other wording such as 'Sepsis-6/Red-Flag Sepsis/Sepsis bloods/BUFALO' (BUFALO is an acronym for the components of Sepsis-6; at NGH this is printed on all charts as an aid memoire with a box for a signature for treatments for audit purposes). Any ambiguity was discussed, and additional documentation reviewed such as ED discharge coding. If a patient had multiple presentations within this period, the first eligible event was included. Direct referrals to specialities were excluded in addition to incomplete electronic records.

Definitions

Cases were retrospectively evaluated according to the Sepsis-3 criteria, defined by evidence of infection and a change in baseline SOFA score ≥ 2 . The baseline score was assumed to be zero where it was not known if there was any pre-existing organ dysfunction. The infection site was determined from culture results, raised inflammatory markers or radiological evidence, where this was not clear cases were discussed to agree whether infection was present. Positive blood cultures, in the absence of a clear source, were diagnosed as bacteraemia. Blood cultures documented as 'likely contaminant organism' were not included. The latter are flagged by the microbiologist by the type of bacteria (skin flora suggesting contaminant at venepuncture, and the time it takes for a BC to flag as positive—small numbers of contaminating bacteria take longer to grow compared with a real bacteraemia). All investigation results were reviewed before a decision on excluding these as usually this is a clinical decision made by assessing the patient and the likelihood that the result does represent a contaminant; often a repeat blood culture is sent.

The SOFA score was modified to substitute PaO₂/FiO₂ (partial pressure of arterial oxygen/fraction of inspired oxygen) with




Figure 1 Summary of the identification of the main study participants.

Table 1 Baseline characteristics

Variables	N	Suspected sepsis
No of patients, n (%)	509	509 (100)
Age (years), median (IQR)	509	74.0 (58.0–82.0)
Female gender, n (%)	509	246 (48.3)
Method of arrival, n (%)	509	
Ambulance	509	415 (81.5)
General practitioner referral	509	27 (5.3)
Self-presented	509	62 (12.2)
Community team referral	509	3 (0.6)
Outpatient clinic referral	509	1 (0.2)
Police transport	509	1 (0.2)
Location within the ED, n(%)		
Resuscitation room	509	228 (44.8)
Majors	509	171 (33.6)
Minors	509	4 (0.8)
Majors transferred to resuscitation room	509	21 (4.1)
Resuscitation room transferred to majors	509	2 (0.4)
Not documented	509	83 (16.3)
Initial emergency department observations, mean±SD or median (IQR)		
RR (breaths per minute) (first available RR used)*	507 (509)	22.9±6 (22.4±6)
HR (beats per minute)	509	105.7±23.4
Systolic BP (mm Hg) (first available SBP used)*	506 (509)	126.5±29.9 (126.4±29.9)
Diastolic BP (mm Hg) (first available DBP used)*	506 (509)	70.2±16.2 (70.2±16.3)
Mean arterial pressure (mm Hg) (first available BP used)*	506 (509)	89.0±19.26 (88.9±19.29)
Saturations (%) (first available saturation used)*	501 (509)	95.3±3.9 (95.2±3.9)
FiO ₂ (first available FiO ₂ used)*	506 (508)	0.21 (0.21–0.35) (0.21 (0.21–0.35))
Temperature (°C) (first available temperature used)*	502 (508)	37.7±1.2 (37.7±1.2)
GCS (first available GCS used)*	478 (506)	15(14–15) (15(14–15))
qSOFA ≥2, n(%)		
Missing values considered as 0 points	509	132 (25.9)
First available values used	509	139 (27.3)
NEWS, median (IQR)		
Missing values considered as 0 points	509	6 (3–8)
First available values used	509	6 (3–8)
Change in SOFA score, median (IQR)	509	2 (0–3)
Change in SOFA ≥2, n(%)	509	256 (50.3)
Definitive infection and change of SOFA ≥2, n(%)	509	251 (49.3)

*If the first value was missing, the next recorded value was used.

BP, blood pressure; ED, emergency department; FiO₂, fraction of inspired oxygen; GCS, Glasgow Coma Scale; HR, heart rate; NEWS, National Early Warning Score; qSOFA, quick Sequential Organ Failure Assessment; RR, respiratory rate; SOFA, Sequential Organ Failure Assessment.

SpO₂/ FiO₂, (oxygen saturations/fraction of inspired oxygen) as previously validated,¹⁵ but was otherwise unchanged.¹

Sample size

This is a descriptive study; therefore, the aim was for a study size sufficient to estimate a typical proportion. A sample size of 500 was chosen, which allowed us to estimate a typical proportion of around 20% with a reasonable degree of precision (ie, a 95% CI 16.6% to 23.8%).

Table 2 Social circumstances and mobility

Variables	Suspected sepsis (n=509)
Place of residence, n (%)	
Home—-independent	217 (42.6)
Home with a care package	83 (16.3)
Home with support from relatives	76 (14.9)
Home with district nurse support	9 (1.8)
Home with support from relatives and district nurse	2 (0.4)
Sheltered/supported accommodation	5 (1.0)
Sheltered/supported accommodation with a care package±district nurse support	5 (1.0)
Intermediate care/active recovery	3 (0.6)
Rehabilitation centre (substance misuse)	2 (0.4)
Care home resident	98 (19.3)
No fixed abode	3 (0.6)
Not documented	6 (1.2)
Mobility, n (%)	
Independent	211 (41.5)
Needs walking aid	181 (35.6)
Wheelchair	22 (4.3)
Needs walking aid or wheelchair	4 (0.8)
Bedbound	36 (7.1)
Bed or wheelchair users	2 (0.4)
Hoist transfers*	16 (3.1)
Other†	20 (3.9)
Not documented	17 (3.3)

*No other information available, so recorded as separate category.

†Included comments that were difficult to classify: 'poor mobility', 'housebound', 'dependent'.

Outcome measures

The outcome measures were descriptive: describing the characteristics, management and outcomes of adults attending the ED with suspected sepsis. Specifically, the prevalence of comorbidities, functional impairment and ceilings of care within this group.

Data extraction

Electronic ED records, discharge summaries, laboratory and radiology reporting systems were reviewed to obtain data including the patient demographics, medical and social history, management (including escalation plan) and outcomes. This was recorded using a standardised extraction form, with explicit definitions for study variables, by two data collectors (LS and LW).

The coroner's office was contacted for those patients who had been referred to them. The General Register Office was contacted for patients that had died in the community to obtain the date and cause of death.

qSOFA and National Early Warning Score (NEWS) were calculated from the initial ED observations.^{1,16} For missing values, the next recorded value was taken, and it was documented that the initial value was missing. To convert the Glasgow Coma Scale (GCS) to the Alert, Voice, Pain, Unresponsive (AVPU) scale for NEWS, GCS ≤13 was accepted as being equivalent to VPU.¹⁷

SOFA scores were calculated from the initial observations and blood results; missing values were assumed to score zero. Charlson Comorbidity Index (CCI) was calculated based on the comorbidities recorded on admission.¹⁸

Statistical analysis

Descriptive statistics are used with 95% CI for key proportions. Continuous data, if normally distributed, are presented as

mean±SD and if skewed, as median (IQR). Categorical data are presented as proportions (percentages).

Parametric assumptions for statistical tests were checked. All statistical data analyses were performed using SPSS V.25.0 (IBM).

Patient and public involvement

The design and methodology of this study was presented to the Sheffield Emergency Care Forum who advised on the study concept.¹⁹

RESULTS

A total of 1750 patients were randomly selected from the blood culture list and reviewed, resulting in 509 patients treated as suspected sepsis (figure 1). Extrapolating this to the total number of BCs taken in the year suggests an estimated incidence of suspected sepsis per year of 1798 cases, which is similar to other studies.^{20 21}

Patient characteristics

Table 1 describes characteristics of the study population. The median age was 74 years (IQR 58–82). 81.5% of the cohort arrived by ambulance and 44.8% were seen in the resuscitation room. Almost 50% met the Sepsis-3 criteria (49.3% (45.0%–53.6%)). The most common suspected sources were chest or urinary and there did not appear to be a seasonal pattern in the date of presentations (online supplemental material).

Regarding the calculated qSOFA, 25.9% (22.1%–29.7%) would have met the qSOFA criteria to suggest investigation for sepsis. The median NEWS score was 6 (IQR 3–8).

Table 2 shows patient social circumstances: 42.6% (38.3%–46.9%) were living at home independently; 17.3% (14.0%–20.6%) with a care package and 19.3% (15.9%–22.7%) in a care home. Less than 50% were mobile independently (41.5% (37.2%–45.8%)). Figure 2 graphically demonstrates that care home residents or those with a package of care are less independently mobile than those at home without a care package.

The comorbidity burden was high with 10.8% of the study population having no comorbidities (table 3). The most common comorbidity was pulmonary disease (30.6%, 26.6%–34.6%), the median CCI was 5 (IQR 2–6).

Table 4 shows that 22% (18.4%–25.6%) of the study population had an existing do not attempt resuscitation (DNAR) order. 39 (7.7%) patients had a discussion regarding resuscitation in the ED (11 of which were by the medical or ICU team).

16.5% (13.3%–19.7%) had an escalation plan explicitly documented. 6.5% (4.4%–8.6%) were referred to ICU; patients were seen by the ICU team and decided appropriate for level 3 care (ICU care), 13 patients for level 2 care (high-dependency unit (HDU)). For both groups, the median NEWS was 11 (IQR 10–11); higher than the median NEWS of 6 (IQR 3–8) for the study population.

Regarding treatment, 98.8% (97.9%–99.8%) of the study population received antibiotics, and 93.3% (91.1%–95.5%) had fluids (online supplemental material).

Just over 13% (13.2%) of patients died in-hospital with a median time from admission to death of 5 days (IQR 2–16).

Sepsis-3-defined sepsis

Online supplemental table 5 includes a summary table for the whole cohort and those that meet the Sepsis-3 definitions of sepsis. Online supplemental tables 6–8 describes the characteristics of the Sepsis-3-defined sepsis cohort in more detail. 49.3% met the Sepsis-3 definitions of sepsis, these patients were older with a median age of 77 years (IQR 65.0–85.0), 90.4% arrived by ambulance a greater proportion were treated in the resuscitation room (59.4%). This is reflected in the physiology with the NEWS scores being higher for this group 7 (IQR 5–10).

Generally, these patients were less independent than the whole cohort; 25.1% were care home residents, with 34.3% living independently at home. Median CCI was 5 (IQR 3–7), and more patients were likely to already have a DNAR in place on presentation (27.5%).




Figure 2 Distribution of mobility of the study population by social circumstance.

Table 3 Comorbidities, CCI and medication

Variables	Suspected sepsis (n=509)
Comorbidity, n (%)	
No comorbidities*	55 (10.8)
Pulmonary disease	156 (30.6)
COPD	105 (20.6)
Hypertension	143 (28.1)
Diabetes Mellitus	143 (28.1)
End-organ damage	20 (3.9)
Ischaemic heart disease	98 (19.3)
Chronic kidney disease	98 (19.3)
CCI criteria moderate to severe CKD	21 (4.1)
Solid tumour (all)	91 (17.9)
Metastatic	30 (5.9)
CVA or TIA	80 (15.7)
Atrial fibrillation	75 (14.7)
Dementia	75 (14.7)
Congestive heart failure	61 (12.0)
Connective tissue disease	27 (5.3)
Peripheral vascular disease	25 (4.9)
Haematological malignancy	17 (3.3)
Liver disease (all)	11 (2.2)
Moderate to severe	8 (1.6)
Peptic ulcer disease	10 (2.0)
CCI score, median (IQR)	5 (2–6)
Immunosuppressant medication, n(%)	
LT steroids	47 (9.2)
Chemotherapy	19 (3.7)
Other (DMARDs, antiproliferative medication, calcineurin inhibitors)	23 (4.5)
None	424 (83.3)
Not documented	8 (1.6)

*0 CCI score.

CCI, Charlson Comorbidity Index; CKD, chronic kidney disease; COPD, chronic obstructive pulmonary disease; CVA, cerebrovascular accident; DMARDs, disease-modifying antirheumatic drugs; LT, long term; TIA, transient ischaemic attack.

This group had a larger proportion referred to ICU (10.4%), and higher in-hospital mortality (19.5%, median 5 days, IQR 13). Of the latter, 19 out of 49 patients had an explicit escalation plan (10 patients for ward-level care, 1 for a palliative care bed and 8 for full escalation).

DISCUSSION

Adults attending the ED with suspected sepsis are elderly (median age 74 years) and dependent, with fewer than half living at home independently or walking independently, and almost 20% from a nursing home. Only 10.8% of the population have no comorbidities and over 20% of the study population have an existing DNAR order. Despite this, only 16.5% have an escalation plan explicitly documented in the ED (ie, the level of care escalation for the patient—ward or higher level such as HDU or ICU). Those meeting the Sepsis-3 definition (49.3%) were particularly elderly (median age 77 years), and likely to be dependent, have comorbidities and an existing DNAR order.

A quarter of patients that met the Sepsis-3 criteria were from a care home. Other authors report similar results.²² Similarly, ICU admission (4.4%) and in-hospital mortality (13.2%) were both within the range reported by other studies.^{23–25}

The reporting of comorbidities in the literature is variable. For example, studies looking at sepsis outcomes in specific populations are likely to have a different distribution of comorbidities,

Table 4 Do not attempt resuscitate decisions and escalation

Variables	Suspected sepsis (n=509)
DNAR in place on presentation, n(%)	
Yes	113 (22.2)
No	318 (62.5)
Not documented	78 (15.3)
DNAR discussed in the ED, n(%)	
Yes, and implemented	15 (2.9)
Yes, and already in place	9 (1.8)
Yes, and not implemented	2 (0.4)
Considered by ED clinician	2 (0.4)
Yes, by the medical/ICU team	11 (2.2)
No documented discussion or form	470 (92.3)
Escalation of care (explicit), n(%)	
Ward level care decision by ED team	25 (4.9)
Ward-level care decision by medical team	5 (1.0)
Full escalation	44 (8.6)
For early medical review	2 (0.4)
Under palliative care team	8 (1.6)
Not documented	425 (83.5)
ICU decisions, n(%)	
Considered ICU, but decision for ward level care by team	17 (3.4)
Decision for ICU referral if not improving	3 (0.6)
Discussed with ICU:	
Admitted under medical team	4 (0.8)
For treatment/medical review initially	2 (0.4)
Referred to ICU	33 (6.5)
Seen by ICU team, and decision for level 0 care	8 (1.6)
Seen by ICU team, and decision for level 1 care	1 (0.2)
Seen by ICU team, and decision for level 2 care	13 (2.6)
Seen by ICU team, and decision for level 3 care	9 (1.8)
Referred to ICU by medical team	2 (0.4)
Not needed*	388 (76.2)
Not documented	62 (12.2)

*Included records where it was explicitly documented that the plan was to transfer to ward or specialty referral, with no specific ceiling of care reasons documented. Where this was not written or if there was any ambiguity then the record was not included in this category.
DNAR, do not attempt resuscitation; ED, emergency department; ICU, intensive care unit.

similarly trials are likely to report fewer comorbidities due to exclusion criteria sometimes encompassing advanced directives or clinician decision regarding suitability for aggressive care.

One of the drivers for doing this study was to define the prevalence of comorbidities, baseline functional state and consideration of escalation decisions in patients managed as sepsis in the ED. There is little current literature that combines these factors together or discusses escalation decisions. Some studies report proportions of patients with a DNAR order in place, and these are similar to these study results, although, of note, the additional information of whether these discussions were had in the ED are not reported in these studies.^{26,27}

The majority of those that died in-hospital did not have an explicit escalation plan documented. This suggests that either implicit ceiling of care decisions are being made and not documented or that recognition of those patients that may require ICU is poor.

Studies have demonstrated that acute infections worsen chronic diseases which can lead to poor longer term outcomes,⁶

therefore knowledge of the prevalence of chronic diseases in an ED population will help improve our understanding of management of sepsis in an ageing population. The SSC advise 'setting goals of care' in the acute setting and having these escalation discussions with patients early.⁷ This study suggests that given the characteristics of the population described the guidelines should reflect and create realistic expectations for both patients and clinicians, not simply a 1-hour target.

Feedback from the patient and public involvement group welcomed these discussions in the ED. Some members related to personal experience of being unaware of a relative's poor prognosis and despite the difficulties surrounding these conversations in a busy environment, overall, they welcomed having these candid discussions.

Physiological scoring may be helpful for predicting the need for intervention, but our findings suggest that comorbidities may have an important role in determining prognosis. Treatment and escalation decisions therefore need to take both physiological scoring and comorbidities into account.

This research area has not been well explored therefore the main strength of this descriptive primary research study is to address this gap and address one of the NICE research recommendations—an epidemiological study on presentation and management of sepsis in England. Many of the previous studies have either used selected cohorts or routine data sources so may have underestimated the rate of comorbidities with little reporting of functional status. In this study, we have looked at the proportion of suspected sepsis patients that have risks of poor outcomes and the need for discussions of care and more individualised treatment goals. We hope this will develop the guidelines that are currently focused on rapid intensive treatment, to include decision making in a population described in this study.

Limitations

All scores were calculated retrospectively; if applied prospectively, this may have altered decision making, for example, if scores reached a threshold for action such as senior involvement. Furthermore, both the calculation of the scores and selection of the cases were performed by the researchers who knew both the outcomes and hypothesis of the study; this could lend itself to observer bias.

Second, using blood cultures to identify the cohort may not identify all relevant cases due to variable compliance with blood cultures being taken (29% in a multicentre study),²⁸ either due to an omission or intentional decision that the investigation was not clinically appropriate for example, during end-of-life care.

Finally, with regard to the SOFA score, the adaptation to the respiratory component due to unavailable arterial oxygen results could have resulted in differing results. Similarly, the cardiovascular component did not consider other medication given for treating hypotension.

Conclusions

This study demonstrates that adults with suspected sepsis are elderly, have substantial functional limitations, comorbidities and treatment directives that should be considered in guidelines, especially recommendations for escalation of care. There must be a balance to create realistic treatment expectations and involvement of patients regarding these.

Contributors LS and SG were responsible for the conception and design of the study, data extraction (LS and LW), and analysis and interpretation (LS and SG). LS drafted the article and all other authors have revised it critically.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval The project was registered with the Clinical Research and Innovation Office and was determined to not require NHS Research Ethics Committee review, as it involves analysis of data collected in routine clinical care. An independent scientific review, local costs and approvals were submitted. The project was also registered and approved by the School of Health and Related Research (SchHARR) Research Ethics Committee. The University of Sheffield is the Research Governance Sponsor.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement All data relevant to the study are included in the article or uploaded as online supplemental information.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

ORCID iDs

Lisa Sabir <http://orcid.org/0000-0001-6488-3314>

Steve Goodacre <http://orcid.org/0000-0003-0803-8444>

REFERENCES

- 1 Singer M, Deutschman CS, Seymour CW, *et al.* The third international consensus definitions for sepsis and septic shock (Sepsis-3). *JAMA* 2016;315:801–10.
- 2 Seymour CW, Liu VX, Iwashyna TJ, *et al.* Assessment of clinical criteria for sepsis: for the third International consensus definitions for sepsis and septic shock (Sepsis-3). *JAMA* 2016;315:762–74.
- 3 Maguire D, Dunn P, McKenna H. *How hospital activity in the NHS in England has changed over time.* The King's Fund Report, 2016.
- 4 Oliver D, Foot C, Humphries R. *Making our health and care systems fit for an ageing population.* King's Fund, 2014.
- 5 Department of Health. Long term conditions compendium of information. 3rd edn [online], 2012. Available: <https://www.gov.uk/government/publications/long-term-conditions-compendium-of-information-third-edition> [Accessed 4 Dec 2018].
- 6 Elfeky S, Golabi P, Otgonsuren M, *et al.* The epidemiologic characteristics, temporal trends, predictors of death, and discharge disposition in patients with a diagnosis of sepsis: a cross-sectional retrospective cohort study. *J Crit Care* 2017;39:48–55.
- 7 Dellinger RP, Levy MM, Rhodes A, *et al.* Surviving sepsis campaign: international guidelines for management of severe sepsis and septic shock, 2012. *Intensive Care Med* 2013;39:165–228.
- 8 Levy MM, Evans LE, Rhodes A. The surviving sepsis campaign bundle: 2018 update. *Intensive Care Med* 2018;44:925–8.
- 9 Fernando SM, Rochweg B, Reardon PM, *et al.* Emergency department disposition decisions and associated mortality and costs in ICU patients with suspected infection. *Crit Care* 2018;22:172.
- 10 Singer M, Inada-Kim M, Shankar-Hari M. Sepsis hysteria: excess hype and unrealistic expectations. *Lancet* 2019;394:1513–4.
- 11 NICE. 2 Admission to critical care | COVID-19 rapid guideline: critical care in adults. Guidance [online]. Available: <https://www.nice.org.uk/guidance/ng159/chapter/2-Admission-to-critical-care> [Accessed 9 Aug 2020].
- 12 NICE. Sepsis: recognition, diagnosis and early management | Guidance and guidelines [online], 2016. Available: <https://www.nice.org.uk/guidance/ng51> [Accessed 4 Sep 2018].
- 13 Population and census [online]. Available: <https://www.sheffield.gov.uk/home/your-city-council/population-in-sheffield> [Accessed 24 Jun 2021].
- 14 Office for National Statistics. Estimates of the population for the UK, England and Wales, Scotland and Northern Ireland [online]. Available: <https://www.ons.gov.uk/peoplepopulationandcommunity/populationandmigration/populationestimates/datasets/populationestimatesforukenglandandwales/scotlandandnorthernireland> [Accessed 24 Jun 2021].
- 15 Pandharipande PP, Shintani AK, Hagerman HE, *et al.* Derivation and validation of Spo₂/Fio₂ ratio to impute for Pao₂/Fio₂ ratio in the respiratory component of the sequential organ failure assessment score. *Crit Care Med* 2009;37:1317–21.
- 16 RCP London. National early warning score (NEWS) 2 [online], 2017. Available: <https://www.rcplondon.ac.uk/projects/outputs/national-early-warning-score-news-2> [Accessed 14 Nov 2018].

- 17 McNarry AF, Goldhill DR. Simple bedside assessment of level of consciousness: comparison of two simple assessment scales with the Glasgow coma scale. *Anaesthesia* 2004;59:34–7.
- 18 Charlson ME, Pompei P, Ales KL, *et al.* A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *J Chronic Dis* 1987;40:373–83.
- 19 Hirst E, Irving A, Goodacre S. Patient and public involvement in emergency care research. *Emerg Med J* 2016;33:665–70.
- 20 Henning DJ, Puskarich MA, Self WH, *et al.* An emergency department validation of the SEP-3 sepsis and septic shock definitions and comparison with 1992 consensus definitions. *Ann Emerg Med* 2017;70:544–52.
- 21 Szakmany T, Lundin RM, Sharif B, *et al.* Sepsis prevalence and outcome on the general wards and emergency departments in Wales: results of a multi-centre, observational, point prevalence study. *PLoS One* 2016;11:e0167230.
- 22 Williams JM, Greenslade JH, Dymond CA, *et al.* Characteristics, treatment and outcomes for all emergency department patients fulfilling criteria for septic shock: a prospective observational study. *Eur J Emerg Med* 2018;25:97–104.
- 23 de Groot B, Stolwijk F, Warmerdam M, *et al.* The most commonly used disease severity scores are inappropriate for risk stratification of older emergency department sepsis patients: an observational multi-centre study. *Scand J Trauma Resusc Emerg Med* 2017;25:91.
- 24 Goulden R, Hoyle M-C, Monis J, *et al.* qSOFA, SIRS and NEWS for predicting in-hospital mortality and ICU admission in emergency admissions treated as sepsis. *Emerg Med J* 2018;35:345–9.
- 25 Redfern OC, Smith GB, Prytherch DR, *et al.* A comparison of the quick sequential (sepsis-related) organ failure assessment score and the National early warning score in Non-ICU patients with/without infection. *Crit Care Med* 2018;46:1923–33.
- 26 Drumheller BC, Agarwal A, Mikkelsen ME, *et al.* Risk factors for mortality despite early protocolized resuscitation for severe sepsis and septic shock in the emergency department. *J Crit Care* 2016;31:13–20.
- 27 Warmerdam M, Stolwijk F, Boogert A, *et al.* Initial disease severity and quality of care of emergency department sepsis patients who are older or younger than 70 years of age. *PLoS One* 2017;12:e0185214.
- 28 Gray A, Ward K, Lees F, *et al.* The epidemiology of adults with severe sepsis and septic shock in Scottish emergency departments. *Emerg Med J* 2013;30:397–401.