

# Association of emergency department boarding times on hospital length of stay for patients with psychiatric illness

Daniel J Lane ,<sup>1</sup> Lauren Roberts,<sup>2</sup> Shawn Currie,<sup>3</sup> Rachel Grimminck,<sup>1,4</sup> Eddy Lang<sup>1,5</sup>

**Handling editor** Loren De Freitas

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/emered-2020-210610>).

<sup>1</sup>University of Calgary, Calgary, Alberta, Canada

<sup>2</sup>Emergency Medicine, University of Saskatchewan, Saskatoon, Saskatchewan, Canada

<sup>3</sup>Addiction and Mental Health Services, Alberta Health Services, Calgary, Alberta, Canada

<sup>4</sup>Psychiatry, Alberta Health Services, Calgary, Alberta, Canada

<sup>5</sup>Department of Emergency Medicine, University of Calgary, Calgary, Alberta, Canada

## Correspondence to

Dr Daniel J Lane, University of Calgary, Calgary, AB T2N 1N4, Canada; [dan.lane@ucalgary.ca](mailto:dan.lane@ucalgary.ca)

Received 6 September 2020

Accepted 12 June 2021

Published Online First

29 June 2021



► <http://dx.doi.org/10.1136/emered-2021-212089>



© Author(s) (or their employer(s)) 2022. No commercial re-use. See rights and permissions. Published by BMJ.

**To cite:** Lane DJ, Roberts L, Currie S, et al. *Emerg Med J* 2022;**39**:494–500.

## ABSTRACT

**Background** Extended periods awaiting an inpatient bed in the emergency department (ED) may exacerbate the state of patients with acute psychiatric illness, increasing the time it takes to stabilise their acute problem in hospital. Therefore, we assessed the association between boarding time and hospital length of stay for psychiatric patients.

**Methods** ED clinical records were linked to inpatient administrative records for all patients with a primary psychiatric diagnosis admitted to a Calgary, Alberta hospital between April 2014 and March 2018. The primary exposure was boarding time (admission decision to inpatient bed transfer), and primary outcome was inpatient length of stay. Confounders for this relationship, including indicators of illness severity, were selected a priori then the association was assessed using hierarchical Bayesian Poisson regression, which accounts for repeat observations of the same patient and differences between hospital sites. Changes in length of stay were measured using a rate ratio (ie, expected change in length of stay for each 1 hour increase in boarding time).

**Results** A total of 19 212 admissions (14 261 unique patients) were included in the analysis. The average boarding time was 14 hours (range: 0–186 hours). Patients who were boarded for greater than 14 hours more frequently required a high-observation bed (14% vs 3.5%), received an antipsychotic (44% vs 14%) or received sedation (55% vs 33%) while in the ED. The probability that boarding time increased hospital length of stay (rate ratio: >1) was 92%, with a median increase for a patient boarded for 24 hours of 0.01 days.

**Conclusion** Boarding in the ED was associated with a high probability of increasing the hospital length of stay for psychiatric patients; however, the absolute increase is minimal. Although slight, this signal for longer length of stay may be a sign of increased morbidity for psychiatric patients held in the ED.

## INTRODUCTION

The emergency department (ED) is the initial site of evaluation for many patients experiencing acute mental health concerns. Psychiatric patients account for an estimated 4% of all patients accessing the ED<sup>1</sup>; however, they often wait longer for admission than non-psychiatric patients.<sup>2–4</sup> This extended period waiting in the ED after an admission decision has been made, known in the literature as boarding time, may exacerbate the psychiatric state

## Key messages

### What is already known on this subject

⇒ Patients with acute psychiatric illness are commonly admitted to the hospital through the emergency department (ED) but may be boarded for extended periods of time before being admitted. Extended boarding times may exacerbate the psychiatric state of these vulnerable patients, resulting in a longer hospitalisation to treat their acute needs.

### What this study adds?

⇒ In this large cohort study, boarding in the ED was associated with a high probability of increased hospital length of stay for psychiatric patients; however, the absolute increase is minimal. Efforts to minimise boarding time for patients with acute psychiatric illness should be considered.

of these vulnerable patients,<sup>5</sup> resulting in a longer hospitalisation to treat their acute needs.


Knowledge of the impact that extended boarding times have on a psychiatric patient may be critical for policy decision making, but no randomised trial can be ethically conducted to assess the causal effect of a system-access exposure such as boarding time. Several descriptive studies have identified factors associated with increased boarding time in the ED, including the specific psychiatric diagnosis,<sup>6,7</sup> use of physical or chemical restraints,<sup>8,9</sup> and the day/time a patient access the ED.<sup>6,10–12</sup> Therefore, utilising knowledge from these prior studies, we hypothesised a potential causal relationship, including known confounders, then assessed the association between boarding time and hospital length of stay for psychiatric patients.

## METHODS

### Design/setting

ED clinical records were linked retrospectively to inpatient records for all patients admitted to hospital with a primary psychiatric diagnosis between April 2014 and March 2018 in Calgary, Alberta, Canada. All hospitals included in this study have general adult EDs with the capacity to admit psychiatric patients to inpatient units. A model for the relationship of interest and significant confounders was





**Figure 1** Directed acyclic graph model for the effect of boarding time on hospital length of stay. Note: patient severity is an unmeasurable construct and, therefore, we used the variables indicated to approximate the patient's severity while in the ED. CTAS, Canadian Triage Acuity Scale; ED, emergency department.

developed a priori through consultation between senior ED and psychiatry physicians and is presented in [figure 1](#).

### Participants

Psychiatric diagnosis was identified by examining the most responsible diagnosis field (ie, first diagnosis code listed) in the Discharge Abstract Database for International Classification of Disease Tenth edition (ICD-10CA) codes beginning with F, which represent psychiatric diagnoses.<sup>13</sup> All adult patients with a most responsible diagnosis consistent with a psychiatric disorder admitted to hospital during the study period were linked to their hospital records. Inpatient diagnosis was used instead of ED diagnosis, as it was hypothesised that this would be more accurate as patients would have more time to be thoroughly assessed by psychiatric specialists and would be unlikely to have a new psychiatric illness evolve during the course of the hospital stay. These diagnoses were classified into 15 unique categories as is described in online supplemental Table 1.

### Measures

The primary outcome of interest was hospital length of stay, defined by a count of calendar days where a patient remained

admitted to the hospital. The primary exposure was boarding time, defined as a continuous measure based on the difference between the admission request by the consulting psychiatrist and the actual time a patient leaves the ED for the ward. Confounders for the association of interest were identified from the ED record and are presented in [figure 1](#): (1) ED wait time (defined as the time from arrival to the time of initial assessment by an ED physician) and ED decision time (defined as the time from initial assessment by an ED physician to the time an admission order was placed) were included as continuous measures; (2) Mental Health Act (MHA) detention, patients who meet criteria for certification under the provincial MHA and will be admitted to the hospital involuntarily, was determined from the extracted triage or nursing note of the same; (3) the number of previous admissions were entered as a count representing the number of times each individual patient had been admitted in the previous 1-year period; (4) the use of physical restraints, chemical sedation/hypnotics (eg, lorazepam) or antipsychotic medication (eg, haloperidol) in the ED was identified in ED care record; (5) requests for a high-observation bed on the inpatient unit (locking secure rooms for the purposes of acute containment of agitation or other risky behaviours,

**Table 1** Characteristics of study patients

	Overall	Boarding <14 hours	Boarding ≥14 hours	SMD for boarding
n	22 447	14 357	8 029	
<b>Patient characteristics</b>				
Age (years) (IQR)	47 (32, 65)	49 (33, 66)	45 (30, 63)	0.117
Male sex (%)	12 127 (54)	7688 (54)	4406 (55)	0.027
Admissions in previous year (%)	0.0 (0.0, 0.0)	0.0 (0.0, 0.0)	0.0 (0.0, 0.0)	0.037
No fixed address (%)	1688 (7.5)	1017 (7.1)	666 (8.3)	0.045
Family MD (%)	14 672 (83.4)	9422 (84.1)	5206 (82.2)	0.052
Primary psychiatric diagnosis				0.176
Anxiety (%)	446 (2.0)	321 (2.2)	124 (1.5)	
Bipolar (%)	1251 (5.6)	744 (5.2)	505 (6.3)	
Conduct (%)	103 (0.5)	58 (0.4)	45 (0.6)	
Depression (%)	2176 (9.7)	1500 (10.4)	671 (8.4)	
Dissociative (%)	109 (0.5)	71 (0.5)	38 (0.5)	
Eating (%)	73 (0.3)	57 (0.4)	16 (0.2)	
Neurocognitive (%)	4222 (18.8)	2783 (19.4)	1434 (17.9)	
Neurodevelopmental (%)	132 (0.6)	67 (0.5)	65 (0.8)	
Obsessive (%)	68 (0.3)	45 (0.3)	23 (0.3)	
Other (%)	159 (0.7)	99 (0.7)	59 (0.7)	
Personality (%)	608 (2.7)	350 (2.4)	250 (3.1)	
Psychosis (%)	3803 (16.9)	2296 (16.0)	1491 (18.6)	
Somatic (%)	85 (0.4)	54 (0.4)	31 (0.4)	
Substance (%)	6614 (29.5)	4406 (30.7)	2198 (27.4)	
Trauma (%)	2594 (11.6)	1505 (10.5)	1076 (13.4)	
Secondary substance use (%)	5428 (24.2)	3311 (23.1)	2103 (26.2)	0.073
Alcohol (%)*	6049 (26.9)	4123 (28.7)	1916 (23.9)	0.110
Opioids (%)*	672 (3.0)	395 (2.8)	276 (3.4)	0.040
Cannabinoids (%)*	1549 (6.9)	849 (5.9)	693 (8.6)	0.105
Sedatives (%)*	254 (1.1)	174 (1.2)	79 (1.0)	0.022
Cocaine (%)*	718 (3.2)	413 (2.9)	305 (3.8)	0.051
Stimulant (%)*	1152 (5.1)	595 (4.1)	555 (6.9)	0.121
Hallucinogen (%)*	35 (0.2)	23 (0.2)	12 (0.1)	0.003
<b>Demographic characteristics</b>				
Arrival means				0.136
No ambulance (%)	11 185 (49.8)	7065 (49.2)	4092 (51.0)	
Air ambulance (%)	21 (0.1)	14 (0.1)	7 (0.1)	
Ground ambulance (%)	10 385 (46.3)	6853 (47.7)	3499 (43.6)	
Police (%)	856 (3.8)	425 (3.0)	431 (5.4)	
<b>ED</b>				
CTAS score (IQR)	2(2,3)	2(2,3)	2(2,3)	0.038
Antipsychotic used (%)	5431 (24.2)	1940 (13.5)	3490 (43.5)	0.704
Sedative used (%)	9188 (40.9)	4799 (33.4)	4383 (54.6)	0.436
Restraints used (%)	3577 (15.9)	2091 (14.6)	1477 (18.4)	0.103
Mental health form (%)	6254 (27.9)	3408 (23.7)	2812 (35.0)	0.250
High observation bed (%)	1607 (7.2)	496 (3.5)	1111 (13.8)	0.376
MD assessment to admit (hours) (IQR)	5.2 (3.5, 7.9)	5.2 (3.5, 8.0)	5.2 (3.5, 7.7)	0.019
Boarding time (hours) (IQR)	6.5 (1.9, 18.8)	2.5 (1.4, 5.9)	23 (18, 38)	1.887
Total ED time (hours) (IQR)	16.9 (8.6, 26.4)	10.2 (6.8, 16.2)	31 (25, 46)	1.787
<b>Primary outcome</b>				
Hospital length of stay (days) (IQR)	8.4 (3.5, 25)	8.6 (3.5, 25)	8.1 (3.8, 26)	0.025

SMD comparing differences between ≥14 hours and <14 hours boarding time.

\*Primary or secondary substance use disorders

CTAS, Canadian Triage Acuity Scale, 1 is most urgent, 5 is least urgent; ED, emergency department; SMD, standardised mean difference.

which represent 10% of inpatient beds) were identified in the admission orders from the ED and (6) secondary substance use was identified from the same inpatient diagnosis codes as the primary diagnosis, and was defined based on the presence of one ICD-10CA code for substance use (F09–F19) in any of the secondary diagnosis fields.

## Analysis

Descriptive statistics with standardised mean differences were used to compare two groups stratified at the mean boarding time. Although boarding time is right-skewed, we elected to stratify the group by the mean for this descriptive analysis as it was selected for a group with more extreme boarding time (≥14

hours) and less extreme boarding time (<14 hours), providing the readers with a better comparison of patient characteristics in these groups. The association of all identified confounders with a patient's boarding time was assessed using a linear regression model. Then, the association of boarding time with hospital length of stay was modelled using Poisson regression. Hierarchical Bayesian regression was used for both the linear and Poisson regression analyses to model the repeat visits for some patients during the study timeline, and potential clustering effects by hospital site. This approach explicitly accounts for similarities of outcome that we expect among the same patient who presents multiple times, and simultaneously accounts for similarity of outcome we expect for the same hospital—a practice that improves the accuracy and generalisability of our estimates.<sup>14</sup> An interaction between the primary psychiatric diagnosis and secondary substance use was included to model the effect modification of substance use on psychiatric disorders. Spline functions were used for the patients age, time of admission, wait time and decision time measures to allow for non-linear associations of these predictors with the outcome. Mean value imputation was used for missing predictors as the rate of missing values was low, while patients missing the primary outcome were excluded from the analysis—rates of missing measures were reported.

The interpretation of estimates from a Bayesian analysis differs from frequentist analysis. In a Bayesian analysis, the estimates of association (ie, linear or rate-ratio estimates in our study) are probability distributions representing the range (width) and likelihood (height) of estimates compatible with the study data. The area under a probability distribution is equal to 1 by definition, therefore, the area of this distribution consistent with the effect of interest (in our case the area consistent with a rate-ratio greater than 1 - expected change in length of stay for each 1 hour increase in boarding time) can be reported as a measure of certainty for the effect of interest.<sup>15</sup> For this study, we also describe the median of the probability distribution (ie, the estimate with 50% of the probability distribution above and below) and provide an estimate of the width of the distribution using 87% posterior intervals (PIs). These intervals do not describe 'statistical significance' but rather the uncertainty range compatible with 87% of the patients (87% was selected to represent the majority of patients but be distinct from the 95% CI conventionally used in frequentist hypothesis testing). Visual displays of the entire distribution are also presented.

All statistical analysis was completed in R statistical and computing software. The 'tableone' package was used for descriptive statistics, and 'brms' packages with Stan were used for Bayesian models.<sup>16 17</sup> Study reporting followed the ROBUST checklist for Bayesian analysis.<sup>18</sup>

### Patient and public involvement

Patients or public were not involved in the conduct of this study.

### RESULTS

Of the 22 477 admissions with a primary psychiatric diagnosis during the study time period (n=16 080 unique patients), 19 212 (85%; n=14 261 unique patients) had the primary outcome of length of stay available and were included in our analysis. The average boarding time for all patients was 14 hours (SD: 17.2 hours), while the median boarding time was 6.5 hours (IQR: 1.9–19 hours) (table 1). Patients who were boarded for greater than 14 hours more frequently required a high observation bed (14% vs 3.5%), received an antipsychotic (44% vs 14%) or received sedation (55% vs 33%) while in the ED. In

crude analysis, there was a minimal difference in median hospital length of stay for patients that were boarded for more than 14 hours compared with those boarded for less than 14 hours (8 days vs 9 days; standardised mean difference: 0.025). Median boarding time and median length of stay both varied between the four hospital sites (online supplemental figures 1 and 2) but not over the study timeline (online supplemental figure 3). Mean value imputation was used for three measures, with the frequency of missing data as follows: wait time (n=565; 2.9%), decision time (n=808; 4.2%) and boarding time (n=43; 0.2%).

### Association of confounders with boarding time

The confounders with the strongest association with increased boarding time were requiring a high observation bed (median increase of 16 hours PI: 16–17), antipsychotic use (median increase of 11 hours PI: 10–11), sedative/hypnotic use (median increase of 7 hours PI: 10–11), arrival by police (median increase of 5 hours PI: 4.2–5.8) or MHA detention (2.4 hours PI: 2.1–2.8) (figure 2). Age, wait time and decision time all had non-linear associations with boarding time; boarding time was generally longer for patients who were older or younger than the median age of 47 years, longer for patients with increased wait times and was not influenced by longer decision times. Patients with conduct or neurodevelopmental psychiatric disorders had the highest estimated boarding time while patients with substance use disorder had the lowest (online supplemental figure 4). Secondary substance abuse resulted in the largest increase in boarding time for patients with neurodevelopmental or dissociative disorders but resulted in decreased boarding time for most psychiatric diagnoses (online supplemental figure 5).

### Predictors of length of stay


The predictors with the strongest association with increased hospital length of stay were air ambulance arrival (rate ratio: 1.6; PI: 1.3–1.9), restraint use in the ED (rate ratio: 1.4; PI: 1.3–1.4) and previous admissions in the past year (rate ratio: 1.1; PI: 1.1–1.1). MHA detention and requiring a high observation bed were also associated with increased hospital length of stay (online supplemental figure 6). Patients with eating disorders or psychosis had the longest length of stay (median: 18 days and 16 days, respectively), while patients with substance abuse, trauma, conduct or dissociative disorders had the shortest (online supplemental figure 7). Secondary substance use had the greatest increase in length of stay when used by patients with dissociative disorders but tended to decrease length of stay for patients with somatic or neurocognitive disorders, or psychosis (online supplemental figure 8).

### Effect of boarding time on length of stay

The probability that increased boarding time resulted in longer hospital length of stay was 92% (figure 3). The median rate ratio from the posterior distribution for each 24-hour increase in boarding time was 1.01 (PI: 1.00–1.01). This translates to 0.01 more days in the hospital for a patient who boarded for 24 hours, or 0.02 more days in the hospital for a patient who boarded for 72 hours.

### DISCUSSION

Our results suggest that extended ED boarding time likely results in a minimal increase in hospital length of stay for patients with primary psychiatric disorders. Several key associations of increased boarding time consistent with previous literature were also identified, which could be targets for interventions aimed




**Figure 2** Probability distributions representing the linear estimates for change in boarding time associated with each parameter. Note: probability distributions represent the range (spread along X-axis) and likelihood (height of distribution) of estimates consistent with the study data. The point indicates median of the distribution, the thick line indicates the 87% PI and the thin line indicates 99% PI. Any area of the probability distribution to the right of 0 indicates an increase in boarding time. Reference category for arrival mode is self-admission. CTAS, Canadian Triage Acuity Scale; MHA, Mental Health Act; PI, posterior interval.

at reducing boarding time in the future. Although our results demonstrate this statistical signal for harm, the absolute increase in length of stay was small and, therefore, boarding time may not reflect a clinically significant cause of morbidity in these patients.

Assessing the causal impact of exposures using observational data is challenging but remains the best means available for providing evidence of potentially harmful exposures (such as boarding time) that cannot be ethically tested in a randomised trial. Our study attempted to quantify the potential morbidity associated with boarding time using an a priori causal model of the relationship, a hierarchical Bayesian analysis and a large, multicentre cohort of patients in an effort to address many of the limitations inherent to observational research. Despite these efforts, unmeasured confounding is always a risk and may account for some of the effect we describe if not correlated with the other confounders we included. Post hoc, we also identified that including all patients (including those admitted to a medical ward rather than strictly a psychiatric unit), we may have selected for many patients that had acute medical needs exacerbating their psychiatric disorders who only required short stays in hospital, reducing the potential effect of boarding time. Additionally, our outcome may have also acted to increase the boarding time of subsequent patients requiring admission, as an increase in length of stay may decrease the availability of inpatient beds when the hospital is nearing capacity. This complex relationship between bed capacity and boarding time was outside the scope of the

current study but may be considered in future studies. Nevertheless, our finding of a high probability that boarding time results in longer length of stay may be a signal for morbidity from this exposure. The small absolute effect we found may also be a positive sign of the resilience of healthcare providers finding ways to initiate treatment for patients with acute psychiatric illnesses while they are boarded in the ED.

Our results were consistent with previous studies identifying the specific psychiatric diagnosis,<sup>6,7</sup> use of physical or chemical restraints,<sup>8</sup> and the day/time a patient accesses the ED as factors associated with increased boarding time.<sup>6,10-12</sup> In addition, we found that detention for mental health assessment and the need for a high-observation bed were strongly associated with increased boarding time. This association may have been due to the severity of the patient, or alternatively may have been correlated with other factors associated with longer lengths of stay. For example, many patients requiring detention or a high-observation bed are agitated or have risky behaviour, and often require substitute decision-makers on a temporary basis while in hospital, or mental health review panels in order to start treatment. This process may increase their boarding time and delay the initiation of treatment, making them more vulnerable to extended hospital length of stay. Furthermore, high-observation beds are a limited resource, which cannot be offloaded to other services, further increasing the boarding time. We also found notable differences in the boarding time and length of stay at



**Figure 3** Estimated probability distribution for the increase in hospital length of stay for each 24-hour increase in boarding time. Note: this probability distribution represents the range (spread along X-axis) and likelihood (height of distribution) of this estimate consistent with the study data. The shaded area of the distribution above 1 indicates the probability consistent with an increased hospital length of stay.

different hospitals. Of particular interest is one site (site D) which had the longest boarding times of all four sites but also had the shortest length of stay. This site uniquely has 24-hour psychiatric nursing care and daily psychiatry rounds in the ED, instead of ED nursing and single psychiatrist providing care. These results may reflect the differences in care models available at different hospital sites.

Concerns about the potential harms of extended boarding times in the ED for patients with psychiatric disorders have been raised by advocacy, physician and government organisations.<sup>5 19 20</sup> Although our results suggest that these interventions specifically targeted to boarding time may have little impact on the patient's hospital length of stay, there are likely other benefits of reducing boarding time that our analysis did not consider, including reducing the distress experienced by all patients, families and care givers in the ED, and addressing the underlying disparities that perpetuate boarding time. Psychiatric boarding may be perpetuated through structural stigma, including funding disparities for mental health services, including inpatient beds, and hospital practices, policies and procedures that restrict the rights of people with mental illness.<sup>21</sup> Distress associated with

extended periods of boarding in the ED on prior admissions may also be a barrier to accessing future psychiatric care for patients.<sup>22</sup> Other studies have proposed solutions to reducing boarding time, including structured discharge planning from psychiatric units,<sup>12</sup> psychiatric rounds in the ED<sup>23</sup> and dedicated psychiatric emergency services.<sup>24</sup> Future studies may consider the causal model we propose (figure 1) to assess the impact of these interventions on alternative outcomes that may better reflect these morbidities.

### Limitations

This study has several limitations. First, we selected patients based on their primary diagnosis of a psychiatric disorder, but this approach may miss patients with serious medical problems likely comorbid to their psychiatric disorder (eg, acute liver failure) that did not have a psychiatric diagnosis documented as their primary diagnosis for that hospital admission. Second, we excluded patients that were missing the primary outcome of hospital length of stay but were unable to determine why this measure was missing from their records. These patients may

represent a unique subpopulation of patients with psychiatric disorders. Third, we did not explicitly consider the inpatient management of these patients in our analysis as all inpatient management could be influenced by the exposure of interest in the ED (ie, on the causal pathway). Factors including treatment type, legal status, patient's capacity and acceptance of treatment, as well as their disposition destination, availability of community supports and outpatient care, no doubt influenced each patient's length of stay. Fourth, we did not consider insurance status of patients as a predictor of boarding time or confounder for length of stay. Although Canada has a universal healthcare system providing no-fee access to hospital for all citizens, private insurance status may reflect patients with better access to community counselling and, therefore, patients at lower risk of requiring hospitalisation. Finally, although we were able to obtain data representing all potential confounders identified a priori in our causal model, the possibility of unmeasured confounders contributing to a spurious effect is always present when randomisation of patients is not possible.

## CONCLUSIONS

Boarding in the ED is a significant health and safety concern in this Canadian jurisdiction with as many as 25% of patients spending nearly a full day awaiting admission under suboptimal circumstances. Boarding was associated with a high probability of increasing the hospital length of stay for psychiatric patients; however, the absolute increase was minimal. Although this signal for harm was slight, longer length of stay may be a sign of increased morbidity for psychiatric patients boarded in the ED for extended periods.

**Twitter** Daniel J Lane @DanLane911 and Eddy Lang @EddyLang1

**Contributors** LR and EL conceived of this study, obtained data and helped with study design. RG and SC helped design the study and interpret the results. DL conducted the analysis and wrote the primary manuscript. All authors contributed to substantial revisions of the manuscript.

**Funding** The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

**Competing interests** None declared.

**Patient consent for publication** Not required.

**Ethics approval** This study was approved by the Conjoint Health Research Ethics Board at the University of Calgary (REB17-0948).

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data availability statement** Data may be obtained from a third party and are not publicly available. The data used in this study were accessed through Alberta Health Services.

**Supplemental material** This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

## ORCID iD

Daniel J Lane <http://orcid.org/0000-0001-8600-8895>

## REFERENCES

- 1 Barratt H, Rojas-García A, Clarke K, et al. Epidemiology of mental health attendances at emergency departments: systematic review and meta-analysis. *PLoS One* 2016;11:e0154449–14.
- 2 Nicks BA, Manthey DM. The impact of psychiatric patient boarding in emergency departments. *Emerg Med Int* 2012;2012:1–5.
- 3 Chakravarthy B, Tenny M, Anderson CL, et al. Analysis of mental health substance Abuse–Related emergency department visits from 2002 to 2008. *Subst Abuse* 2013;34:292–7.
- 4 Zhu JM, Singhal A, Hsia RY. Emergency department length-of-stay for psychiatric visits was significantly longer than for Nonpsychiatric visits, 2002–11. *Health Aff* 2016;35:1698–706.
- 5 Nordstrom K, Berlin JS, Nash SS, et al. Boarding of mentally ill patients in emergency departments: American psychiatric association resource document. *West J Emerg Med* 2019;20:690–5.
- 6 Stephens RJ, White SE, Cudnik M, et al. Factors associated with longer length of stay for mental health emergency department patients. *J Emerg Med* 2014;47:412–9.
- 7 Smith JL, De Nadai AS, Storch EA. *Correlates of length of stay and boarding in Florida emergency departments for patients with psychiatric diagnoses*. Psychiatr Serv. VA: American Psychiatric Association Arlington, 2016: 67(11). 1169–74.
- 8 Chang G, Weiss AP, Orav EJ, et al. Bottlenecks in the emergency department: the psychiatric clinicians' perspective. *Gen Hosp Psychiatry* 2012;34:403–9.
- 9 Weiss AP, Chang G, Rauch SL, et al. Patient- and practice-related determinants of emergency department length of stay for patients with psychiatric illness. *Ann Emerg Med* 2012;60:162–71.
- 10 Chang G, Weiss A, Kosowsky JM, et al. Characteristics of adult psychiatric patients with stays of 24 hours or more in the emergency department. *Psychiatric Services* 2012;63:283–6.
- 11 Bastiampillai T, Schrader G, Dhillon R, et al. Impact of a psychiatric unit's daily discharge rates on emergency department flow. *Australas Psychiatry* 2012;20:117–20.
- 12 Stover PR, Harpin S. Decreasing psychiatric admission wait time in the emergency department by facilitating psychiatric discharges. *J Psychosoc Nurs Ment Health Serv* 2015;53:20–7.
- 13 ICD-10-CA International Statistical Classification of Diseases and Related Health Problems. Canadian Institute for health information 2015:1–1074.
- 14 Gelman A, Carlin JB, Stern HS. *Bayesian data analysis*. 3rd. CRC press, 2013: 1.
- 15 McElreath R. *Statistical rethinking: a Bayesian course with examples in R and Stan*, 2018.
- 16 Yoshida K, Bohn J. Tableone: Create "Table 1" to Describe Baseline Characteristics [Internet]. R package version 0.7.3, 2015. Available: <https://CRAN.R-project.org/package=tableone>
- 17 Bürkner P-C. Advanced Bayesian multilevel modeling with the R package brms. *R J* 2018;10:395–17.
- 18 Sung L, Hayden J, Greenberg ML, et al. Seven items were identified for inclusion when reporting a Bayesian analysis of a clinical study. *J Clin Epidemiol* 2005;58:261–8.
- 19 Treatment Advocacy Center. Delayed and deteriorating: serious mental illness and psychiatric boarding in emergency departments 2019.
- 20 Bender D, Pande N, Ludwig M. A literature review: psychiatric boarding. *U.S. Department of Health and Human Services* 2009:1–30.
- 21 Corrigan PW, Markowitz FE, Watson AC. Structural levels of mental illness stigma and discrimination. *Schizophr Bull* 2004;30:481–91.
- 22 Allen MH, Carpenter D, Sheets JL, et al. What do consumers say they want and need during a psychiatric emergency? *J Psychiatr Pract* 2003;9:39–58.
- 23 Blumstein H, Singleton AH, Suttentfield CW, et al. Weekday psychiatry faculty rounds on emergency department psychiatric patients reduces length of stay. In: Gerson L, ed. *Acad Emerg Med*. John Wiley & Sons, Ltd, 2013: 20. 498–502.
- 24 Zeller S, Calma N, Stone A. Effects of a dedicated regional psychiatric emergency service on boarding of psychiatric patients in area emergency departments. *West J Emerg Med* 2014;15:1–6.

## Appendix

eTable 1: Categories of Psychiatric Diagnosis

Category	Description
Anxiety	Agoraphobia, Anxiety disorder, unspecified, Elective mutism, Generalized anxiety disorder, Hypochondriacal disorder, Organic anxiety disorder, Other phobic anxiety disorders, Other specified anxiety disorders, Panic disorder [episodic paroxysmal anxiety], Social anxiety disorder of childhood, Social phobias
Bipolar	Bipolar affective disorder, current episode hypomanic, Bipolar affective disorder, current episode manic with psychotic symptoms, Bipolar affective disorder, current episode manic without psychotic symptoms, Bipolar affective disorder, current episode mild or moderate depression, Bipolar affective disorder, current episode mixed, Bipolar affective disorder, current episode severe depression with psychotic symptoms, Bipolar affective disorder, current episode severe depression without psychotic symptoms, Bipolar affective disorder, currently in remission, Bipolar affective disorder, unspecified, Cyclothymia, Hypomania, Mania with psychotic symptoms, Mania without psychotic symptoms, Manic episode, unspecified, Other bipolar affective disorders, Other manic episodes
Conduct	Conduct disorder, unspecified, Habit and impulse disorder, unspecified, Oppositional defiant disorder, Other conduct disorders, Other habit and impulse disorders, Other mixed disorders of conduct and emotions, Pathological fire-setting [pyromania]
Depression	Depressive conduct disorder, Depressive episode, unspecified, Dysthymia, Mixed anxiety and depressive disorder, Moderate depressive episode, Organic mood [affective] disorders, Other depressive episodes, Other recurrent depressive disorders, Other recurrent mood [affective] disorders, Persistent mood [affective] disorder, unspecified, Recurrent depressive disorder, current episode mild, Recurrent depressive disorder, current episode moderate, Recurrent depressive disorder, current episode severe with psychotic symptoms, Recurrent depressive disorder, current episode severe without psychotic symptoms, Recurrent depressive disorder, currently in remission, Recurrent depressive disorder, unspecified, Severe depressive episode with psychotic symptoms, Severe depressive episode without psychotic symptoms, Unspecified mood [affective] disorder

Category	Description
Dissociative	Depersonalization-derealization syndrome, Dissociative [conversion] disorder, unspecified, Dissociative amnesia, Dissociative convulsions, Dissociative motor disorders, Mixed dissociative [conversion] disorders, Organic amnesic syndrome, not induced by alcohol and other psychoactive substances, Other dissociative [conversion] disorders
Eating	Anorexia nervosa, Atypical anorexia nervosa, Bulimia nervosa, Eating disorder, unspecified, Other eating disorders
Neurocognitive	Delirium not superimposed on dementia, so described, Delirium superimposed on dementia, Delirium, unspecified, Mild cognitive disorder, Mixed cortical and subcortical vascular dementia, Multi-infarct dementia, Other delirium, Other vascular dementia, Subcortical vascular dementia, Unspecified dementia, Vascular dementia, unspecified
Neurodevelopmental	Asperger's syndrome, Atypical autism, Childhood autism, Combined vocal and multiple motor tic disorder [de la tourette], Developmental disorder of scholastic skills, unspecified, Disturbance of activity and attention, Mild mental retardation with the statement of no, or minimal, impairment of behaviour, Mild mental retardation without mention of impairment of behaviour, Mild mental retardation, other impairments of behaviour, Mixed disorder of scholastic skills, Other developmental disorders of scholastic skills, Other disorders of psychological development, Other pervasive developmental disorders, Other specified behavioural and emotional disorders with onset usually occurring in childhood and adolescence, Pervasive developmental disorder, unspecified, Receptive language disorder, Stuttering [stammering], Tic disorder, unspecified, Unspecified disorder of psychological development, Unspecified mental retardation without mention of impairment of behaviour
Obsessive	Obsessive-compulsive disorder, unspecified, Other obsessive-compulsive disorders, Other specified neurotic disorders, Predominantly compulsive acts [obsessional rituals], Predominantly obsessional thoughts or ruminations
Other	OTHER identity disorder, unspecified, Mental disorder, not otherwise specified, Mild mental and behavioural disorders associated with the puerperium, not elsewhere classified, Nightmares, Nonorganic insomnia, Other OTHER identity disorders, Other mental and behavioural disorders associated with the puerperium, not elsewhere classified, Other specified mental disorders due to brain damage and

Category	Description
	dysfunction and to physical disease, Postconcussional syndrome, Sadomasochism, Severe mental and behavioural disorders associated with the puerperium, not elsewhere classified, Transsexualism, Unspecified behavioural syndromes associated with physiological disturbances and physical factors, Unspecified mental disorder due to brain damage and dysfunction and to physical disease, Unspecified organic or symptomatic mental disorder, Unspecified organic personality and behavioural disorder due to brain disease, damage and dysfunction, Vomiting associated with other psychological disturbances
Personality	Anankastic personality disorder, Anxious [avoidant] personality disorder, Dependent personality disorder, Dissocial personality disorder, Emotionally unstable personality disorder, Histrionic personality disorder, Mixed and other personality disorders, Organic personality disorder, Other organic personality and behavioural disorders due to brain disease, damage and dysfunction, Other specific personality disorders, Other specified disorders of adult personality and behaviour, Paranoid personality disorder, Personality disorder, unspecified, Unspecified disorder of adult personality and behaviour
Psychosis	Acute and transient psychotic disorder, unspecified, Acute schizophrenia-like psychotic disorder, Catatonic schizophrenia, Delusional disorder, Hebephrenic schizophrenia, Induced delusional disorder, Organic delusional [schizophrenia-like] disorder, Other acute and transient psychotic disorders, Other acute predominantly delusional psychotic disorders, Other nonorganic psychotic disorders, Other persistent delusional disorders, Other schizoaffective disorders, Other schizophrenia, Paranoid schizophrenia, Residual schizophrenia, Schizoaffective disorder, depressive type, Schizoaffective disorder, manic type, Schizoaffective disorder, mixed type, Schizoaffective disorder, unspecified, Schizoid personality disorder, Schizophrenia, unspecified, Schizotypal disorder, Undifferentiated schizophrenia, Unspecified nonorganic psychosis
Somatic	Intentional production or feigning of symptoms or disabilities, either physical or psychological [factitious disorder], Other somatoform disorders, Persistent somatoform pain disorder, Somatization disorder, Somatoform autonomic dysfunction, Somatoform disorder, unspecified
Substance	Abuse of non-dependence-producing substances, Mental and behavioural disorders due to multiple drug use and use of psychoactive substances, acute intoxication, Mental and behavioural disorders due

Category	Description
	<p>to multiple drug use and use of psychoactive substances, dependence syndrome, Mental and behavioural disorders due to multiple drug use and use of psychoactive substances, harmful use, Mental and behavioural disorders due to multiple drug use and use of psychoactive substances, other mental and behavioural disorders, Mental and behavioural disorders due to multiple drug use and use of psychoactive substances, psychotic disorder, Mental and behavioural disorders due to multiple drug use and use of psychoactive substances, residual and late-onset psychotic disorder, Mental and behavioural disorders due to multiple drug use and use of psychoactive substances, unspecified mental and behavioural disorder, Mental and behavioural disorders due to multiple drug use and use of psychoactive substances, withdrawal state, Mental and behavioural disorders due to multiple drug use and use of psychoactive substances, withdrawal state with delirium, Mental and behavioural disorders due to use of alcohol, acute intoxication, Mental and behavioural disorders due to use of alcohol, amnesic syndrome, Mental and behavioural disorders due to use of alcohol, dependence syndrome, Mental and behavioural disorders due to use of alcohol, harmful use, Mental and behavioural disorders due to use of alcohol, other mental and behavioural disorders, Mental and behavioural disorders due to use of alcohol, psychotic disorder, Mental and behavioural disorders due to use of alcohol, residual and late-onset psychotic disorder, Mental and behavioural disorders due to use of alcohol, unspecified mental and behavioural disorder, Mental and behavioural disorders due to use of alcohol, withdrawal state, Mental and behavioural disorders due to use of alcohol, withdrawal state with delirium, Mental and behavioural disorders due to use of cannabinoids, acute intoxication, Mental and behavioural disorders due to use of cannabinoids, dependence syndrome, Mental and behavioural disorders due to use of cannabinoids, harmful use, Mental and behavioural disorders due to use of cannabinoids, other mental and behavioural disorders, Mental and behavioural disorders due to use of cannabinoids, psychotic disorder, Mental and behavioural disorders due to use of cannabinoids, unspecified mental and behavioural disorder, Mental and behavioural disorders due to use of cannabinoids, withdrawal state, Mental and behavioural disorders due to use of cocaine, acute intoxication, Mental and behavioural disorders due to use of cocaine, dependence syndrome, Mental and behavioural disorders due to use of cocaine, harmful use, Mental and behavioural disorders due to use of cocaine, other mental and behavioural disorders, Mental and behavioural disorders due to use of cocaine, psychotic disorder, Mental and behavioural disorders due to use of cocaine, unspecified mental and behavioural disorder, Mental and behavioural disorders due to use of cocaine, withdrawal state, Mental</p>

Category	Description
	<p>and behavioural disorders due to use of cocaine, withdrawal state with delirium, Mental and behavioural disorders due to use of hallucinogens, acute intoxication, Mental and behavioural disorders due to use of hallucinogens, dependence syndrome, Mental and behavioural disorders due to use of hallucinogens, harmful use, Mental and behavioural disorders due to use of hallucinogens, psychotic disorder, Mental and behavioural disorders due to use of opioids, acute intoxication, Mental and behavioural disorders due to use of opioids, dependence syndrome, Mental and behavioural disorders due to use of opioids, harmful use, Mental and behavioural disorders due to use of opioids, other mental and behavioural disorders, Mental and behavioural disorders due to use of opioids, psychotic disorder, Mental and behavioural disorders due to use of opioids, unspecified mental and behavioural disorder, Mental and behavioural disorders due to use of opioids, withdrawal state, Mental and behavioural disorders due to use of opioids, withdrawal state with delirium, Mental and behavioural disorders due to use of other stimulants including caffeine, acute intoxication, Mental and behavioural disorders due to use of other stimulants including caffeine, dependence syndrome, Mental and behavioural disorders due to use of other stimulants including caffeine, harmful use, Mental and behavioural disorders due to use of other stimulants including caffeine, other mental and behavioural disorders, Mental and behavioural disorders due to use of other stimulants including caffeine, psychotic disorder, Mental and behavioural disorders due to use of other stimulants including caffeine, residual and late-onset psychotic disorder, Mental and behavioural disorders due to use of other stimulants including caffeine, unspecified mental and behavioural disorder, Mental and behavioural disorders due to use of other stimulants including caffeine, withdrawal state, Mental and behavioural disorders due to use of other stimulants including caffeine, withdrawal state with delirium, Mental and behavioural disorders due to use of sedatives or hypnotics, acute intoxication, Mental and behavioural disorders due to use of sedatives or hypnotics, dependence syndrome, Mental and behavioural disorders due to use of sedatives or hypnotics, harmful use, Mental and behavioural disorders due to use of sedatives or hypnotics, other mental and behavioural disorders, Mental and behavioural disorders due to use of sedatives or hypnotics, psychotic disorder, Mental and behavioural disorders due to use of sedatives or hypnotics, unspecified mental and behavioural disorder, Mental and behavioural disorders due to use of sedatives or hypnotics, withdrawal state, Mental and behavioural disorders due to use of sedatives or hypnotics, withdrawal state with delirium, Mental and behavioural disorders due to use of volatile solvents, dependence syndrome, Pathological gambling</p>

Category	Description
Trauma	Acute stress reaction, Adjustment disorders, Other reactions to severe stress, Post-traumatic stress disorder, Reaction to severe stress, unspecified


Bayesian regression requires specification of distributions representing the possible range of effect for each parameter prior to incorporating the data. These “prior distributions” for each of the parameters were defined following prior predictive simulations. All prior values are described in eTable 2. Convergence of models was assessed using trace-plots, the number of effective-samples (i.e. an assessment of correlation between iterations), and the Gelman-Rubin convergence diagnostic for each parameter.

eTable 2: Prior distributions defined for each parameter

Linear model			Poisson model		
Parameter	Distribution	Values	Parameter	Distribution	Values
Intercept	Normal	Mean: 0, SD: 10	Intercept	Normal	Mean: 0, SD: 2
Coefficients	Normal	Mean: 0, SD: 5	Coefficients	Normal	Mean: 0, SD: 1
Hyperprior	Cauchy	Location: 0, Scale: 1	Hyperprior	Cauchy	Location: 0, Scale: 1
Spline	Student T	Df: 3, Mu: 0, Sigma: 10	Spline	Student T	Df: 3, Mu: 0, Sigma: 10


Note: Poisson model priors are defined on the log scale, therefore narrower prior distributions are appropriate. SD, standard deviation; Df, degrees of freedom

eFigure 1: Variation in average boarding time per site




Note: Area of distribution represents probability. Point indicates median, thick line indicates 87% posterior interval, thin line indicates 99% posterior interval.

eFigure 2: Variation in length of stay per site




Note: Area of distribution represents probability. Point indicates median, thick line indicates 87% posterior interval, thin line indicates 99% posterior interval.

eFigure 3: Change in median length of stay over study period




eFigure 4: Estimated boarding time specific to a patient’s psychiatric diagnosis




Note: Area of distribution represents probability. Point indicates median, thick line indicates 87% posterior interval, thin line indicates 99% posterior interval.

eFigure 5: Change in boarding time for each psychiatric diagnosis when patient also had secondary substance use disorder



Note: Area of distribution represents probability. Point indicates median, thick line indicates 87% posterior interval, thin line indicates 99% posterior interval.

eFigure 6: Rate ratios representing a patient's change in hospital length of stay




Note: Area of distribution represents probability. Point indicates median, thick line indicates 87% posterior interval, thin line indicates 99% posterior interval.

Area to the right of 1.0 represent and increased length of stay, area to the left represent a decrease.


MHA = mental health act, CTAS = Canadian Triage Acuity Scale

eFigure 7: Estimated hospital length of stay specific to a patient's psychiatric diagnosis



Note: Area of distribution represents probability. Point indicates median, thick line indicates 87% posterior interval, thin line indicates 99% posterior interval.

eFigure 8: Change in hospital length of stay for each psychiatric diagnosis when patient also had secondary substance use disorder



Note: Area of distribution represents probability. Point indicates median, thick line indicates 87% posterior interval, thin line indicates 99% posterior interval. Area to the right of 1.0 represent and increased length of stay, area to the left represent a decrease.