

	Random sample (N = 66)	Other patients (N=760)
Age, mean (SD)	70.4 (IQR 55.56 - 79.25)	62.53 (IQR 52.73 - 75.36)
Males, N (%)	35 (53%)	509 (67%)
Triage colour, N (%)		
<i>green</i>	36 (54.5%)	373 (49.1%)
<i>yellow</i>	22 (33.3%)	275 (36.2%)
<i>red</i>	8 (12.1%)	111 (14.6%)
Institutionalized, N (%)	4 (6.1%)	50 (6.6%)
Stroke/TIA, N (%)	3 (4.5%)	34 (4.5%)
Parkinson, N (%)	0 (0%)	9 (1.2%)
Alzheimer disease, N (%)	0 (0%)	9 (1.2%)
Other dementias, N (%)	3 (4.5%)	51 (6.7%)
Epilepsy, N (%)	2 (3%)	23 (3%)
Other neurologic diseases, N (%)	3 (4.5%)	59 (7.8%)
Depression, N (%)	2 (3%)	29 (3.8%)
Number of chronic drugs, median (IQR)	2 (IQR 1 - 6.25)	2 (IQR 0 - 5)
Anticholinergic burden score, median (IQR)	0 (IQR 0 - 1)	0 (IQR 0 - 0)
Psychoactive drugs, N (%)	8 (12.1%)	70 (9.2%)
<i>Sedative-hypnotics</i>	4 (6.1%)	46 (6.1%)
<i>Antidepressants</i>	4 (6.1%)	56 (7.4%)
<i>Antipsychotics</i>	2 (3%)	37 (4.9%)
<i>Memantine</i>	0 (0%)	1 (0.1%)
<i>Acetylcholinesterase inhibitors</i>	1 (1.5%)	4 (0.5%)
<i>Dopaminergic drugs</i>	0 (0%)	12 (1.6%)

p	Statistica test
0.10	U Mann Whitney
0.03	Chi square test
0.7	Chi square test
1	Fisher exact test
1	Fisher exact test
1	Fisher exact test
1	Fisher exact test
0.79	Fisher exact test
1	Fisher exact test
0.47	Fisher exact test
1	Fisher exact test
0.14	U Mann Whitney
0.20	U Mann Whitney
0.51	Chi square test
1	Fisher exact test
1	Fisher exact test
0.76	Fisher exact test
1	Fisher exact test
0.34	Fisher exact test
0.61	Fisher exact test