

HCV RISK FACTOR SURVEY

The following is a list of commonly known risk factors associated with the contraction of Hepatitis C. The answers to these questions are solely for statistical data and are not related to your care here today. Please answer the questions truthfully and to the best of your ability with a checkmark next to each relevant response

Have you now, or have you ever in the past:

1. Used injectable drugs Yes No
2. Shared needles with another person for any purpose Yes No
3. Had an accidental needlestick injury from a high-risk individual (Injectable drug user, Hepatitis C infected individual, dialysis patient, hemophiliac) Yes No
4. Had tattoo work done outside of the United States Yes No
5. Had tattoo work done outside of a commercial tattoo parlor Yes No
6. Had a body piercing done outside of a commercial body piercing store Yes No
7. Had surgical or dental work done outside of the United States Yes No
8. Have received blood or blood products outside of the United States Yes No
9. Received blood or blood products before 1992 Yes No
10. Been an organ recipient Yes No
11. Been a hemophilia patient Yes No
12. Been a hemodialysis patient Yes No
13. Had HIV-positive status Yes No
14. Have been born to an HIV infected mother Yes No
15. Have been born to a Hepatitis C infected mother Yes No
16. Have been born to a mother who was known to be using injectable drugs Yes No
17. Shared bathroom products with a Hepatitis C infected individual for more than 1 year Yes No