

## HCV RISK FACTOR SURVEY

The following is a list of commonly known risk factors associated with the contraction of Hepatitis C. The answers to these questions are solely for statistical data and are not related to your care here today. Please answer the questions truthfully and to the best of your ability with a checkmark next to each relevant response

Have you now, or have you ever in the past:

1. Used injectable drugs  Yes  No
2. Shared needles with another person for any purpose  Yes  No
3. Had an accidental needlestick injury from a high-risk individual (Injectable drug user, Hepatitis C infected individual, dialysis patient, hemophiliac)  Yes  No
4. Had tattoo work done outside of the United States  Yes  No
5. Had tattoo work done outside of a commercial tattoo parlor  Yes  No
6. Had a body piercing done outside of a commercial body piercing store  Yes  No
7. Had surgical or dental work done outside of the United States  Yes  No
8. Have received blood or blood products outside of the United States  Yes  No
9. Received blood or blood products before 1992  Yes  No
10. Been an organ recipient  Yes  No
11. Been a hemophilia patient  Yes  No
12. Been a hemodialysis patient  Yes  No
13. Had HIV-positive status  Yes  No
14. Have been born to an HIV infected mother  Yes  No
15. Have been born to a Hepatitis C infected mother  Yes  No
16. Have been born to a mother who was known to be using injectable drugs  Yes  No
17. Shared bathroom products with a Hepatitis C infected individual for more than 1 year  Yes  No