




Adult SATS Chart



EMERGENCY

- Obstructed Airway - not breathing
- Seizure- current
- Burn - facial / inhalation
- Hypoglycaemia - glucose less than 3
- Cardiac arrest

VERY URGENT

- High energy transfer (severe mechanism of injury)
- Shortness of breath - acute
- Level of consciousness reduced / confused
- Coughing blood
- Chest pain
- Stabbed neck
- Haemorrhage - uncontrolled (arterial bleed)
- Seizure- post ictal
- Focal neurology - acute (stroke)
- Aggression
- Threatened limb
- Eye injury
- Dislocation of larger joint (not finger or toe)
- Fracture - compound (with a break in skin)
- Burn over 20%
- Burn - electrical
- Burn - circumferential
- Burn - chemical
- Poisoning / Overdose
- Diabetic - glucose over 11 & ketonuria
- Vomiting fresh blood
- Pregnancy and abdominal trauma
- Pregnancy and abdominal pain
- Severe pain

URGENT

- Haemorrhage - controlled
- Dislocation of finger OR toe
- Fracture - closed (no break in skin)
- Burn - other
- Abdominal pain
- Diabetic- glucose over 17 (no ketonuria)
- Vomiting persistently
- Pregnancy and trauma
- Pregnancy and PV bleed
- Moderate pain

ADULT TEWS

Older than 12 years / taller than 150 cm tall							
	3	2	1	0	1	2	3
Mobility				Walking	With Help	Stretcher/immobility	
RR		less than 9		9 - 14	15 - 20	21 - 29	more than 29
HR		less than 41	41 - 50	51 - 100	101 - 110	111 - 129	more than 129
SBP	Less than 71	71 - 80	81 - 100	101 - 199		more than 199	
Temp		Cold OR Under 35°		35° - 38.4°		Hot OR Over 38.4°	
AVPU		Confused		Alert	Reacts to Voice	Reacts to Pain	Unresponsive
Trauma				No	Yes		

CHECK FOR ADDITIONAL INVESTIGATIONS

If RR scores 1 point or more on TEWS	Check SpO ₂ and hand over to SHCP to give O ₂ Do a finger prick glucofetest if patient is diabetic
Reduced level of consciousness (not alert including confused)	Do a finger prick glucofetest and hand over to SHCP
Diabetes and Hyperglycaemia (glucofetest 11 mmol/L or more)	Urine dipstick to check for ketones
Unable to sit up/ need to lie down	Do a finger prick glucofetest and hand over to SHCP
Chest pain	Immediate ECG and hand over to SHCP
Active seizure / fitting	Do a finger prick glucofetest and hand over to SHCP IV access - NO intramuscular
History of diabetes	Do a finger prick glucofetest and hand over to SHCP
Hypoglycaemia (glucofetest 3 mmol/L or less)	Move to resus hand over to SHCP and give something to eat or drink
Abdominal pain or backache: female	Urine dipsticks and Urine pregnancy test