

Table 1S: comparison of the main characteristics of the included and excluded patients

	Included patients (N = 826)	Excluded patients (N=21)	p	Statistica test
Age, mean (SD)	63.7 (SD 15.16)	70.9 (IQR 42.6 -81.1)	0.67	U Mann Whitney
Males, N (%)	544 (65.9%)	14 (66.7%)	0.94	Chi square test
Triage colour, N (%)				
<i>green</i>	409 (49.5%)	10 (47.6%)	0.61	Chi square test
<i>yellow</i>	297 (36%)	6 (28.6%)		
<i>red</i>	119 (14.4%)	4 (19%)		
Institutionalized, N (%)	54 (6.5%)	0 (0%)	0.63	Fisher exact test
Stroke/TIA, N (%)	37 (4.5)	0 (0%)	1	Fisher exact test
Parkinson, N (%)	9 (1.1%)	0 (0%)	1	Fisher exact test
Alzheimer disease, N (%)	9 (1.1%)	0 (0%)	1	Fisher exact test
Other dementias, N (%)	54 (6.5%)	0 (0%)	0.62	Fisher exact test
Epilepsy, N (%)	25 (3%)	0 (0%)	1	Fisher exact test
Other neurologic diseases, N (%)	62 (7.5%)	0 (0%)	0.63	Fisher exact test
Depression, N (%)	31 (3.8%)	2 (9.5%)	0.14	Fisher exact test
Number of chronic drugs, median (IQR)	2 (IQR 0 - 5)	2 (IQR 0 - 6)	0.79	U Mann Whitney
Anticholinergic burden score, median (IQR)	0 (IQR 0 - 0)	0 (IQR 0 - 0)	0.37	U Mann Whitney
Psychoactive drugs, N (%)	78 (9.4%)	1 (4.8%)	1	Fisher exact test
<i>Sedative-hypnotics</i>	50 (6.1%)	0 (0%)	1	Fisher exact test
<i>Antidepressants</i>	60 (7.3%)	1 (4.8%)	1	Fisher exact test
<i>Antipsychotics</i>	39 (4.7%)	0 (0%)	1	Fisher exact test
<i>Memantine</i>	1 (0.1%)	0 (0%)	1	Fisher exact test
<i>Acetylcholinesterase inhibitors</i>	5 (0.6%)	0 (0%)	1	Fisher exact test
<i>Dopaminergic drugs</i>	12 (1.5%)	0 (0%)	1	Fisher exact test